

received
4/9/07

**PETITION TO LIST FIVE SPECIES OF ROCKFISH IN THEIR PUGET SOUND
PROPER DISTINCT POPULATION SEGMENTS AS ENDANGERED OR
THREATENED SPECIES UNDER THE ENDANGERED SPECIES ACT (ESA)**

TO: SECRETARY OF COMMERCE, UNITED STATES DEPARTMENT OF
COMMERCE, NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION,
NATIONAL MARINE FISHERIES SERVICE.

From: Sam Wright (Petitioner), 1522 Evanston Ct. NE, Olympia, Washington, 98506.
(Tel. 360-943-4424). Petitioner is a fish biologist with 45 years experience in managing
fish populations and fish habitat and is Certified as a Fisheries Professional (CFP) by the
American Fisheries Society.

Subject: Petition the Secretary of Commerce to list as Endangered or Threatened the
Puget Sound Proper Distinct Population Segments (DPSs) of bocaccio (*Sebastes
paucispinis*), canary rockfish (*Sebastes pinniger*), yelloweye rockfish (*Sebastes
ruberrimus*), greenstriped rockfish (*Sebastes elongatus*) and redstripe rockfish (*Sebastes
proriger*) and to designate critical habitat. Brown rockfish (*Sebastes auriculatus*), copper
rockfish (*Sebastes caurinus*) and quillback rockfish (*Sebastes maliger*) were previously
evaluated for ESA listing in the following report: Stout, H.A., B.B. McCain, R.D. Vetter,
T.L. Builder, W.H. Lenarz, L.L. Johnson, and R.D. Methot. 2001. Status review of
Copper Rockfish, Quillback Rockfish, and Brown Rockfish in Puget Sound, Washington.
U.S. Dept. Commer., NOAA Tech. Memo. NMFS-NWFSC-46, 158 p. Stout et al. (2001)
was prepared in response to an earlier effort by the Petitioner (Wright, S. 1999. Petition
to the Secretary of Commerce to list as threatened or endangered 18

“species/populations” of “Puget Sound” marine fishes and to designate critical habitat.
Petition to the U. S. National Marine Fisheries Service, February 1999.). There were 13
species of rockfish originally proposed for status review but only the three most abundant
(and also with the most information) were actually examined. It is logical to assume that
these are the three that would be the least likely to become threatened or endangered. The
inherent flaw with this “data dependent” approach for rockfish is that other species with a
much lower population abundance were rejected for status review. As expected, in late
2004, Washington Department of Fish and Wildlife (WDFW) biologist Greg Bargmann
declared that “Yelloweye, Bocaccio, and Canary have virtually disappeared.” He noted
that “rockfish do not survive being brought to the surface so there is no conservation
benefit to release after incidental take.” (San Juan County Marine Resources Committee,
minutes of November 3, 2004 meeting). Bargmann also reported that “Discarded catch is
substantial and smaller first-caught fish seem to be replaced with larger, later-caught fish
(high grading)” and that “The rockfish encounter rate is very high during salmon fishing”.

Stout et al. (2001) defined DPSs for greater Puget Sound populations of three
rockfish species, including DPSs for a portion of the area defined as Puget Sound Proper.
Due to multiple uses of the term “Puget Sound”, Stout et al. (2001:xiii) “adopted
conventions for geographical regions in the inland waters of Washington State and British
Columbia” and thus defined Puget Sound Proper “as marine waters south of Admiralty
Inlet and east of Deception Pass”. However, the report did not recommend ESA listings

based on selected scientific evidence examined at the time of report preparation. Stout et al. (2001) did concede that the three DPSs met the Musick et al. (2000) criteria to be considered *vulnerable* pending further analysis. The next step that should have been followed from Musick et al. (2000) would be to consider the Puget Sound Proper DPSs for copper and quillback rockfish under the categories of "Small range and endemics" and "Specialized habitat requirements" (Musick et al. 2000:7). Except for their early life history stages, the critical rocky reef habitat for most rockfish species in Puget Sound Proper is mainly non-contiguous and totals only a meager *14 square kilometers* (or less than 2% of total habitat), a small fraction of the habitat available to other Pacific Coast rockfish populations. This group of closely related species must practice habitat partitioning, a basic tenet of evolution. Thus, each individual species can only use part of the available critical habitat. To make matters even worse, "Many vacant habitats have been observed" (Palsson and Pacunski 1998:13). In addition, Puget Sound Proper is, by a wide margin, the poorest and most heavily modified rockfish habitat on the entire Pacific Coast, highlighted recently by frequent water quality problems and fish kills in Hood Canal.

We don't
use this
method

Another critical flaw or omission in Stout et al. (2001) was a failure to recognize the advantages of longevity in rockfish species and the implications to population dynamics of management policies that ignore these evolved advantages. These issues had been recognized in the scientific literature at least as early as 1984 (Leaman, B.M., and R.J. Beamish. 1984. Ecological and management implications of longevity in some northeast Pacific groundfishes. International North Pacific Fisheries Commission Bulletin 42:85-97). The above reference (Leaman and Beamish 1984) did not even appear in the Citations section of Stout et al. (2001). In terms of fisheries resource management, the concept of substantial "yield" is mutually exclusive with maintenance of the age composition structure in rockfish species.

A feature article in the August 2004 issue of *Fisheries* dealt with the same subject (Berkeley, S.A., M.A. Hixon, R.J. Larson, and M.S. Love. 2004. Fisheries sustainability via protection of age structure and spatial distribution of fish populations. *Fisheries* 29(8):23-32.). They believed that fishing truncates the size and age structure of fish populations. Research indicates that old-growth age structure, combined with a broad spatial distribution of spawning and recruitment, is at least as important as spawning biomass in maintaining sustainable populations. Berkeley et al. (2004) found that older, larger female rockfish produce larvae that withstand starvation longer and grow faster than the offspring of younger fish, that stocks may actually consist of several reproductively isolated units, and that recruitment may come from only a small and different fraction of the spawning population each year.

NOAA Fisheries recently disputed the critical value of maintaining longevity in a rockfish population but this challenge was limited to a single component of the multiple arguments advanced by Berkeley et al. (2004) (Varanasi, U. 2006. Memorandum from U. Varanasi (Northwest Fisheries Science Center) to D.R. Lohn (Northwest Regional Administrator) RE: Petition to list Puget Sound quillback and copper rockfish. 10/24/2006. 5pp.). Anyone who disputes the value of maintaining longevity in a rockfish population needs to explain why such a trait evolved if it was not essential to survival of the species.

Dulvy et al. (2004:258) must have had species like Puget Sound Proper rockfish in mind when they concluded that “The intrinsic factors or correlates underlying the major routes to local, regional and global scale extinction in the sea appear to be *large body size, small geographical range and ecological specialization*. The key intrinsic factor appears to be a high degree of exposure to a causal factor, i.e., *high catchability of exploited species*. These patterns are broadly similar to the main correlates of global scale freshwater and terrestrial extinctions.” (emphasis added) (Dulvy, N.K., Ellis, J.R., Goodwin, N.B., Grant, A., Reynolds, J.D., and S. Jennings. 2004. Methods of assessing extinction risk in marine fish. *Fish Fish.* 5:255-276.).

The Washington State Legislature recently stated in House Bill 1076 (Read first time 01/10/2007) that “The department of fish and wildlife has classified certain rockfish species within Puget Sound as critically depressed. These common species of rockfish have *undergone dramatic declines in Puget Sound and the coast during the past three decades*.” (emphasis added). By the mid-1970s, Puget Sound Proper had only a ten species aggregation of rockfish, a sub-set of the large number of rockfish species present in the Pacific Ocean. Now, three decades later, this has been cut in half to a five species aggregation. Canary rockfish undoubtedly suffered the largest decline in abundance ranking relative to other members of the rockfish assemblage, falling from the top three in the early 1960s to “virtually disappeared”.

Basis for the Petition

Section 4 of the ESA contains provisions allowing interested persons to petition the Secretary of Commerce or the Secretary of Interior to add a species to, or remove a species from, the List of Endangered or Threatened Wildlife. Petitioner files this petition under the Endangered Species Act, 16 U.S.C. section 1531-1543 (1982), its implementing regulations, 50 C.F.R. part 424, and the Administrative Procedures Act, 5 U.S.C. section 553 ©. The National Marine Fisheries Service (NMFS) has jurisdiction over this petition under 16 U.S.C. section 1533 (a) and the August 26, 1974 Memorandum of Understanding Between the U.S. Fish and Wildlife Service and National Marine Fisheries Service Regarding Jurisdictional Responsibilities and Listing Procedures Under the Endangered Species Act of 1973.

Basis for the DPS Delineation

Stout et al. (2001:xiv) summarized determinations of three Puget Sound Proper DPSs as follows: “The BRT examined environmental, geological, biogeographic, life history, and genetic information in the process of identifying DPSs. In particular, biogeography, ecological and habitat factors, and genetic population structure were found to be most informative for the species considered in this status review. Based on this examination, the BRT identified a DPS for each of the three rockfish species in Puget Sound Proper *that can be considered a species under the ESA*.” (emphasis added).

The overwhelming evidence presented by Stout et al. (2001) indicates that all of the other rockfish species will also have Puget Sound Proper as a DPS. There does not appear to be any genetic evidence available for bocaccio, canary rockfish, yelloweye rockfish, greenstriped rockfish and redstripe rockfish. However, all of the “environmental, geological, biogeographic and life history” factors fully apply.

Perhaps the most compelling argument is the following from Stout et al. (2001:91-92): "According to Terrie Klinger, out of more than 6,500 releases of drift cards in the San Juans, and more than 2,600 recoveries, 8 were recovered south of Admiralty Inlet. Of these 8, 7 were recovered just south of Admiralty Inlet, and 1 was recovered a year later south of Seattle near Des Moines. Klinger's interpretation is that virtually nothing enters the main basin of Puget Sound proper from the San Juan Islands or Eastern Basin of the Strait of Juan de Fuca on the surface. Thus, it is not likely that North-Puget Sound is a significant source of surface-associated pelagic early life-history stages of rockfish to Puget Sound proper. However, general circulation studies identify some mixing of subsurface waters, so limited exchange is possible."

Assessment in 2007: There does not appear be any critical flaws in the original assessment or any compelling recent information from the past six years that would justify a re-examination of the Puget Sound DPSs previously defined by Stout et al.(2001).

Basis for the Risk Conclusions

Stout et al. (2001:xv,xvii) summarized the risk conclusions for copper rockfish as follows: "The BRT utilized criteria and methods of Wainwright and Kope (1999) and Musick et al. (2000) to assist in organizing the information presented regarding risk to the Puget Sound proper DPS. Bearing the results of the above comparisons in mind, the BRT considered whether the Puget Sound proper DPS was in danger of extinction, likely to become in danger of extinction, or not likely to become in danger of extinction. However, most members expressed concern they could not entirely rule out the possibility that this species at present is likely to become in danger of extinction. The BRT also concluded that this DPS met the International Union for the Conservation of Nature (IUCN) criteria to be considered vulnerable (Musick et al. 2000). The BRT agreed that populations of this species had declined over the last three or four decades, with over-harvesting being a likely factor. Nevertheless, the populations in the DPS had appeared to be stable over the past five years, and the lower population numbers in this DPS compared to the larger numbers in northern Puget Sound are roughly in proportion to the greater amounts of kelp and high-relief habitat in North Puget Sound. The BRT considered the risk to copper rockfish in North Puget Sound to be no greater than the risk to copper rockfish in Puget Sound proper. The BRT also expressed caution that important changes in resource management (e.g., increased harvest levels) and in the ecosystem (e.g., increased numbers of marine mammals or predatory fish species), as well as increased habitat degradation, could result in increased extinction risk for copper rockfish in this DPS."

Stout et al. (2001:xix) summarized the risk conclusions for quillback rockfish as follows: "The BRT, bearing in mind their deliberations regarding risk, using West (1997), Wainwright and Kope (1999), and Musick et al. (2000), considered whether the Puget Sound proper DPS of quillback rockfish was in danger of extinction, likely to become in danger of extinction or not likely to become in danger of extinction. The majority of the BRT concluded that the Puget Sound proper DPS of quillback rockfish are neither at risk of extinction nor likely to become so. However, most members expressed concern that they could not rule out the possibility that this species at present is likely to

become in danger of extinction. The BRT also concluded that this DPS met the IUCN criteria to be considered vulnerable. The BRT agreed that populations of quillback rockfish had, according to a self-contained underwater breathing apparatus (SCUBA) survey, declined to 14% of its 1988 size, with over-harvesting being a likely major factor. Nevertheless, the populations in the DPS had appeared to be stable over the last five years, and the lower population numbers in this DPS compared to the larger numbers in North Puget Sound are roughly in proportion to the greater amounts of kelp and high-relief habitat in North Puget Sound. The BRT considers the risk to quillback rockfish in North Puget Sound to be no greater than the risk to quillback rockfish in Puget Sound proper. The BRT also expressed the same caution as they did with copper rockfish, which is that important changes in resource management practices (e.g., increased harvest levels) and in the ecosystem (e.g., increased numbers of marine mammals or predatory fish species), as well as increased habitat degradation, could result in increased extinction risk for this species in Puget Sound proper and in North Puget Sound.”

Stout et al. (2001:xxi) summarized the risk conclusions for brown rockfish as follows: “The BRT used methods and criteria from Wainwright and Kope (1999) and Musick et al. (2000) to organize information regarding risk to the Puget Sound proper DPS of brown rockfish. They considered whether the species was in danger of extinction, likely to become in danger of extinction or not likely to become in danger of extinction. A majority of the BRT concluded that brown rockfish in Puget Sound proper are neither at risk of extinction nor likely to become so. Factors in this decision included the increasing numbers of brown rockfish observed in SCUBA surveys in central Puget Sound proper during the late-1990s. Moreover, brown rockfish are more habitat generalists than quillback and copper rockfish and consume a wider range of prey species, making them more adaptable to the types of intertidal and subtidal habitats and associated food organisms available in the DPS. However, most members also expressed concern that they could not entirely rule out the possibility that this species is at present likely to become in danger of extinction. The BRT considered brown rockfish in North Puget Sound and the Strait of Juan de Fuca to be associated with the Puget Sound proper DPS and to be vagrants from that DPS. The BRT expressed the same caution as they did with copper and quillback rockfish, that important changes in resource management practices (e.g., increased harvest levels) and in the ecosystem (e.g., increased numbers of marine mammals or predatory fish species) as well as increased habitat degradation, could result in increased risk of extinction for brown rockfish in greater Puget Sound.”

Problems with the Risk Conclusions

Failure to Recognize the Values of Longevity. A serious omission by Stout et al. (2001) was a complete failure to even consider the values of longevity in rockfish species, especially since this was specifically identified as a critical problem in the original ESA petition that included these three rockfish species (Wright 1999). This knowledge was readily available in the scientific literature as early as 1984 (Leaman and Beamish 1984). This failure irretrievably tainted all of the “assessments” described above.

Leaman and Beamish (1984:88) described the importance of longevity as follows: “The most obvious potential benefit of a relatively long life is a long reproductive life.

The extension of the period of reproduction reduces the risk that a long period of unfavourable environmental conditions will result in the loss of a stock. When the period between favourable environmental conditions for a species is relatively long (5-15 yr) it appears that the life span is also relatively long." Leaman and Beamish (1984:87) also determined that "Maintenance of unique age composition among geographically proximate stocks implies little mixing of adult fish among stocks."

Leaman and Beamish (1984:95) conclude their Discussion as follows: "The characteristics of the species considered in this paper typify the problems of managing exploited populations of longer-lived fish. Their extreme life spans appear to be an adaptive feature for ensuring evolutionary persistence under reproductive uncertainty, rather than for maximization of population growth rate. These species have repeated spawning and exploitation has often reduced the number of spawning groups and placed the responsibility for population maintenance on fewer, younger age-groups. There are additional ecological considerations such as the importance of continuously occupying a space, the importance of dispersed spawning, or the selective advantage of specific reproductive products. Extreme longevity in fishes may also allow adults to exist in low productivity environments where the probability of regular recruitment is very low. Exploitation strategies for fishes with long reproductive lives must consider the evolved traits that enable it to meet particular environmental and evolutionary challenges. The production and growth-oriented management strategies derived for shorter-lived species may result in rapid over-exploitation of accumulated biomass and a prolonged period of rehabilitation for the long-lived species. In extreme cases, a competing species may capitalize on vacant space or resources and prevent the over-exploited species from regaining its former abundance."

Stout et al. (2001) did not even consider the Leaman and Beamish (1984) report since it was not listed in their Citations section. The earlier report was based mainly on the simple yet very compelling logic that certain fish species would not be characterized by extreme longevity unless it was essential for ensuring their sustainability. Two decades later, a new report (Berkeley et al. 2004) would provide a great deal more empirical evidence to demonstrate that this logic is indeed correct (see subsequent New Information section).

Problems with Wainwright and Kope (1999) Approach. The first problem, as stated above, is failure to address the longevity issue. In addition, the three risk conclusions from Stout et al.(2001) cited above are incorrect in stating that Musick (2000) was used in conjunction with Wainwright and Kope (1999) for all three species. In fact, the text of Stout et al. (2001) reveals that Wainwright and Kope (1999) was used exclusively to make the three determinations. Musick et al. (2000). was only used to make vulnerable ratings after the ESA determinations had already been finalized. West (1997) did not play any meaningful role in the risk conclusions for quillback rockfish.

The fundamental problem with using Wainwright and Kope (1999) is that scientists must make numerical ratings for population parameters that they are unwilling or unable to quantify directly. This gives a naïve reader the illusion of some sort of quantitative scientific analysis but is merely putting arbitrary numbers on some very subjective and strictly qualitative personal opinions. The recommendations for or against ESA listings become largely dependent upon resource management philosophies and

← not appropriate anyway

areas of expertise of individual BRT members. However, Stout et al. (2001) used this approach as “proof” that three Puget Sound Proper rockfish DPSs were not at-risk. In this particular case, failure of the Wainwright and Kope (1999) process to adequately discriminate is clearly evident in very similar risk conclusions being reached for copper and quillback rockfish in both Puget Sound Proper and North Puget Sound as well as for brown rockfish that are more habitat generalists. Five very different factual situations produced essentially the same end result using Wainwright and Kope (1999).

Problem with Unrecognized High Risk Factors in Puget Sound Proper. The North Puget Sound area has a huge advantage in suitable habitat for rockfish. Pacunski and Palsson (1998) determined that there were over 200 square kilometers of rocky reef habitat in North Puget Sound as opposed to only *14 square kilometers* in all of the remaining Puget Sound Proper basins. In addition, the Puget Sound Proper habitat is typically non-contiguous and a significant part of it is “vacant” in term of the presence of rockfish. There were also wide disparities described in Stout et al. (2001) favoring North Puget Sound with respect to presence of chemical contaminants, degree of human modification of the shoreline, extent of marine vegetation and rockfish growth rates.

The Main Basin of Puget Sound Proper receives about 80% of the amount of waste discharged from urban and industrial point-sources in the entire region (PSWQA 1988). The WDNR (1998) estimates that 52% of the shoreline in the Main Basin has been modified by human activities and it has a relatively small amount of intertidal vegetation, with a point estimate of 28.3% coverage (Bailey et al. 1998). Quillback rockfish from Puget Sound Proper had slower growth rates and were smaller than fish from North Puget Sound (West and O’Neill 1995). Comparable data were not available for copper rockfish and brown rockfish are rare in North Puget Sound.

The only place in Puget Sound Proper where it is even possible to estimate what an unfished or “virgin” population of rockfish might look like is the 27 acre Edmonds Underwater Park (EUP), which has been protected since 1970. Palsson and Pacunski (1998) estimated that the reproductive potential of copper rockfish at EUP exceeded the potential of the average fished site in central Puget Sound by a ratio of 55 to one - and they did not even assign any extra value to production from larger fish on a per egg basis. The difference due to different length frequency distributions between no-take and fished sites accounted for a four-fold increase in estimated egg production while the difference due to densities accounted for almost a fifteen-fold increase in estimated egg production.

Problems with Musick et al. (2000) Approach. In addition to failure with respect to the longevity issue, Musick et al. (2000) was not even employed in determination of the ESA risk assessments. It was only used after the fact to reach vulnerable ratings. Musick et al. (2000) has an inherent methodology advantage over Wainwright and Kope (1999) in that risk ratings are determined directly from quantitative measures of population parameters. In addition, the results in Musick et al. (2000) are simply the application to individual fish species of a rating system already finalized in Musick (1999) and there was no opportunity for other scientists to achieve modifications to it (Musick, J.A. 1999. Criteria to define extinction risk in marine fishes. Fisheries 24(12):6-14.). Note: Petitioner was one of the co-authors of Musick et al. (2000) and did not support the basic approach used (Wright, S. 2002. A critical flaw in the American Fisheries Society initiative to protect marine, estuarine, and diadromous fish stocks:

SO WHAT!

Failure to account for depensation. Prepared for Center for Biological Diversity, Oakland, CA.).

Another critical flaw in Stout et al. (2001) is that they often portray the *vulnerable* in Musick et al. (2000) as being from the "IUCN" (which is incorrect) and as a definite at-risk step below *threatened*. Musick et al. (2000:7) describe the specific intent as follows: "If the decline equals or exceeds the threshold for the appropriate productivity category, the DPS would be automatically listed as *vulnerable* and flagged for further study by expert scientists, who may decide to upgrade the level of threat to *threatened* or *endangered*, or downgrade the status, if appropriate." Musick et al. (2000) is essentially an incomplete process. The intended step described above was never taken for the three Puget Sound Proper rockfish DPSs.

If Stout et al. (2001) had used Musick et al. (2000) as actually written, they might have taken the interim *vulnerable* classification and then focused on the following two sections from Musick et al. (2000:7) to finalize their work: "(2) **Small range and endemics:** Species that are endemic or restricted in range to some relatively small, contiguous geographic entity (i.e., island, archipelago, river system, etc.) in which the habitat is or may be under threat of degradation or destruction should be classified as *vulnerable*. Where significant habitat loss has occurred or is occurring, such species should be classified as *threatened* or *endangered*. Significant habitat loss should be evaluated on a case by case basis in the context of the biology of the DPS under consideration and on both the amount of critical habitat available and the vulnerability of that habitat. (3) **Specialized habitat requirements:** Some species may be relatively widespread but occupy very specific habitats within their range and/or during some specific life history stage. Therefore, their area of occupancy may represent only a small part of that range. When habitats are particularly vulnerable (such as coral reefs or seagrass beds) and subject to degradation, destruction, or fragmentation, habitat loss could be the critical factor leading to population reduction or extirpation. Habitat loss should be examined as a risk factor in the context of the biology of the DPS under consideration."

The critical question *not quantified* by Musick et al. (2000) is how small can the geographical range be before there is a high risk of extinction. The IUCN *Red List* uses thresholds called "extent of occurrence" and "area of occupancy" (IUCN 2001 cited in Dulvy et al. 2004). Extent of occurrence is the area within the shortest continuous imaginary boundary that can be drawn to encompass the taxon. Area of occupancy is the area actually occupied by a taxon within its extent of occurrence (Hilton-Taylor 2000 cited in Dulvy et al. 2004). In order to qualify as threatened, the extent of occurrence must be less than 20,000 square kilometers and the area of occupancy must be less than 2,000 square kilometers. The entire Greater Puget Sound Basin only covers an area of approximately 2,330 square kilometers (Stout et al. 2001). Puget Sound Proper appears to account for less than half of this total and is clearly below either of the IUCN thresholds. The extent of occurrence is only about 5% of the 20,000 square kilometer threshold. The actual area of occupancy - 14 square kilometers of rocky reef habitat - is only 0.7% of the 2,000 square kilometer threshold.

Problems with the Sustainable Fisheries Act. During Petitioner's involvement in the American Fisheries Society initiative described above - which included a number

of NMFS scientists - it became evident that there was a definite policy objective (written or unwritten) within NMFS to handle any problems with at-risk marine fish populations under the Sustainable Fisheries Act (SFA) of 1997, not the Endangered Species Act. This belief was reinforced by extensive discussion of the subject by Stout et al. (2001:126-129). However, "inland waters of Puget Sound proper and North Puget Sound were excluded in the original language of the MSFCMA in 1976." (Stout et al. 2001:126). The NMFS probably has some understandable concerns over the issue of "precedents" between Puget Sound Proper DPSs and marine fish species managed under the SFA. However, the Puget Sound Proper DPSs for rockfish have only a small fraction of suitable habitat as compared to any of the species managed in the ocean. In addition, rockfish species managed under the SFA by the Pacific Fishery Management Council have rebuilding plans that span many decades and reach a high of 92 years for bocaccio, (Shipp, R.L. 2003. A perspective on marine reserves as a fishery management tool. Fisheries 28(12):10-21.). These plans are based on population modeling with numerous tenuous assumptions, including inherent beliefs that the underlying spawner-recruit relationships will be at least stable (i.e., a flat trend line) for most of this century and that they will never have the "multiple domains of attraction" described by Peterman (1977). From the outside perspective of the Petitioner, it seems ludicrous to make these types of assumptions in light of the current state of knowledge on global warming. It will be decades before it is even known if any of these plans can be successful. In any case, it is illogical to champion a similar approach for rockfish DSPs in Puget Sound Proper which only have a tiny fraction of the suitable rockfish habitat available in the ocean but the maximum degree of human-caused environmental stress.

Failure to Recognize Ecological Interactions with Hatchery Chinook. The WDFW professional staff has, for many years, recognized the potential adverse impacts on rockfish populations from their massive hatchery program for Chinook salmon (West 1997). For obvious reasons, this opinion has been effectively muzzled by the WDFW administration in recent years, particularly since the numbers of hatchery Chinook released have been reduced and most marine area net pen rearing programs have been eliminated (WDFW and Puget Sound Treaty Tribes. 2004. Puget Sound Chinook salmon hatcheries. A Component of the Comprehensive Chinook Salmon Management Plan.). However, the reduced potential for adverse impacts on rockfish populations is more illusion than fact.

WDFW and Puget Sound Treaty Tribes (2004:4) state the following under *Ecological Interactions*: "4) releasing fish at a time, size, and physiologically condition that provides a low likelihood of residualization and promotes rapid migration through the estuary to marine waters. Programs typically release subyearling Chinook salmon that are in the 40 to 90 fish per pound (77 to 100mm fork length) during the months of May and June. Fish released at this time are expected to move rapidly through estuarine areas; 5) releasing subyearling fish that are a larger size than natural-origin Chinook salmon of the same brood year to reduce the potential for diet overlap with any co-occurring natural origin fish in marine waters."

The current hatchery Chinook program shows that 45.6 million subyearlings and 2.6 yearlings are released annually in the Greater Puget Sound Basin. Approximately 74% of the subyearlings and 88% of the yearlings are released in Puget Sound Proper

(WDFW and Puget Sound Treaty Tribes 2004). Millions of these fish are now released from rearing sites on small stream systems that did not originally have significant natural Chinook populations and reach marine waters in times ranging from a matter of minutes (e.g., Hoodspout Hatchery on Finch Creek) to a few hours - Tulalip Bay, Grovers Creek, Gorst Creek, Chambers Creek, Minter Creek, Deschutes River (Capitol Lake), Big Beef Creek. The *biomass* of hatchery fish actually reaching Puget Sound marine waters each year is undoubtedly at an all time high. During the massive fin-marking studies of lower Columbia River hatchery fall run chinook in the 1960s, it was estimated that only about 30% of the fish survived to reach the estuary. This was not a dam passage problem since most of the fish were released below Bonneville Dam or in the Bonneville pool. The primary problems were small average fish size, premature (too early) release timing and diseases - problems also common to Puget Sound in the 1960s (WDF. 1992. Salmon 2000 technical report. Phase 2: Puget Sound, Washington Coast and integrated planning.).

New Information on the Status of Puget Sound Rockfish DPSs

Species with Extreme Longevity - A Special Case. In recent years, it has become obvious that status of rockfish populations cannot be judged in terms of traditional resource management paradigms - that each kilogram of spawning stock biomass is identical regardless of adult age, that all larvae have an equal probability of survival regardless of parental age, and that any effects of fishing mortality are reversible. Berkeley et al. (2004:24) provide the proper context for judging rockfish as follows: "we believe that at least part of the explanation for stock collapses is the result of our failure to appreciate the value of both large old fish and fine-scale spatial dynamics of recruitment in the replenishment of fish populations. We discuss recent research that provides what we believe to be a compelling case that the age structure of a stock combined with the spatial distribution of recruitment are as important as spawning biomass in maintaining long-term sustainable population levels. In particular, there are an increasing number of examples of complex population structure in species currently managed as a unit stock, and increasing evidence that only a small fraction of spawners in a stock - those that spawn at the right time and place, which varies annually - successfully contribute to each new cohort. Moreover, it is large, old female fish that produce offspring most likely to recruit successfully to these new cohorts. Based on this evidence, we believe that the best and perhaps only way to ensure old-growth age structure and complex spatial structure in populations of groundfish is through interconnected networks of fully protected marine reserves."

A Single 80cm Fish is Worth Nearly Ten 40cm Fish. An important consideration for Puget Sound Proper rockfish comes from Berkeley et al. (2004:27): "It is well documented that fecundity increases nearly linearly with body mass in adult teleost fishes, and geometrically with body length, which is a decelerating function of age (Weatherly 1972; Wootton 1990). This relationship is due to larger fish not only having a greater body volume for holding eggs, but also devoting a greater proportion of energy stores to egg production. Thus, a 40-cm TL (0.65-kg) bocaccio rockfish produces an average of just over 200,000 eggs per year, whereas an 80-cm (5.41-kg) fish at double the length produces nearly 2 million eggs, nearly 10 times the fecundity (Love et al. 1990).

In other words, considering only fecundity per se - let alone egg or larval quality - a single 80-cm bocaccio is worth nearly ten 40-cm fish.”

Importance of Maternal Age and Larval Quality. Some of the most relevant quantitative information now available regarding status of Puget Sound Proper rockfish DPSs comes from recent studies documenting greater values for older spawners that extend far above their higher fecundity. Berkeley et al. (2004:28) provided the following discussion on the subject: “A variety of studies indicate that egg and larval size and/or viability also increase with female size and age (Chambers et al. 1989; Zastrow et al. 1989; Buckley et al. 1991). Recent experiments with black rockfish by Berkeley et al. (2004), in which gravid females were held until parturition and the larvae reared under controlled conditions, revealed that maternal age was much more predictive of larval growth and survival than either maternal size or condition index. Larvae from the oldest fish (age 17) survived starvation 2.5 times longer than those of the youngest fish (age 5) (Figure 2b), and grew more than 3 times as fast on the same diet (Figure 2a). These differences may be conservative since larvae were reared under constant environmental conditions, with ad libitum rations in fed treatments and no exposure to predators or competitors (other than their siblings). Results of a multiple regression analysis indicated that maternal age accounted for most of the variability in larval survival and growth. Maternal length provided a small but significant increase in goodness-of-fit. These results suggest that older females produce higher quality larvae, and that females that are both old and large produce the highest quality larvae. The mechanism appears to be the volume of the larval oil globule at birth, which is strongly related to maternal age (Berkeley et al. 2004).

It seems likely that such large differences in growth and starvation tolerance have a profound effect on larval survival and subsequent recruitment (review by Heath 1992). The ability of larval fish to survive a period of starvation is often critical due to the spatial and temporal unpredictability of encountering patches of zooplankton prey (Letcher and Rice 1997). Fast growth has clear benefits in allowing larvae to more quickly pass through the most vulnerable life history stages and to develop more rapidly those morphological and physiological capabilities that improve detection and capture of prey, avoidance of predators, and resistance to other environmental challenges (Miller et al. 1988; Bailey and Houde 1989). Indeed, field studies on marine fish larvae have demonstrated that differences in growth rate, especially in the youngest larvae, can have a profound effect on survival. A doubling of the growth rate in larval bluefish (*Pomatomus saltatrix*) and Atlantic cod (*Gadus morhua*) can increase survival by a factor of 5-10 (Meekan and Fortier 1996; Hare and Cowen 1997). Larval quality can also vary in terms of behavioral attributes directly related to survival (Fuiman and Cowan 2003).”

Importance of Adult Body Size and Condition. Berkeley et al. (2004:27) have the following discussion on the subject: “In rockfishes and in other teleosts, body condition and deposition of lipid reserves increase disproportionately with fish length or stage of maturity (Iles 1974; Eliasson and Vahl 1982; Larson 1991). If volumetric variables such as body weight, liver weight, and weight of fat reserves grow proportionately to the rest of the fish, they would increase with the cube of body length. In rockfish, Guillemot et al. (1985) found varying relationships between mesenteric fat weight and fish length (often with low R² values). However, Larson (1991), benefiting

from fish consistently collected in the same areas (in contrast to the fish available to Guillemot), found disproportionate relationships between measures of condition and fish length in kelp rockfish (*Sebastes atrovirens*) and black-and-yellow rockfish (*S. chrysomelas*). Body weight was proportional to slightly greater than the cube of length, liver weight to around the 4th power of length, and mesenteric fat weight to about the 6th power of length. Whether body reserves are utilized directly in reproduction (MacFarlane et al. 1993; Norton and MacFarlane 1994), for overwintering maintenance (Guillemot et al. 1985), or a combination of the two (Larson 1991), the amount of reserves may affect the timing (Lenarz and Wyllie Echeverria 1986) or amount (Eliasson and Vahl 1982; Lambert and Dutil 2000; Blanchard et al. 2003) of reproduction, and the potential for overwintering or post-spawning survival (Guillemot et al. 1985; Lambert and Dutil 2000). The positive allometry of reserves and body condition in fishes indicates that larger (and presumably older) fish may make greater reproductive contributions than smaller fish, and also that larger fish may be able to reproduce and survive reproduction under a greater range of environmental conditions than smaller fish.”

Importance of Maternal Age and Time of Spawning. For Puget Sound Proper rockfish, yet another distinct advantage is described by Berkeley et al. (2004:27): “Older marine teleosts generally spawn earlier during the reproductive season than younger fish (Berkeley and Houde 1978; Pedersen 1984; Lambert 1987). In a recent study of black rockfish (*Sebastes melanops*) off Oregon, Bobko and Berkeley (2004) found that older fish gave birth earlier in the year than progressively younger fish (a trend also noted in yellowtail rockfish, *Sebastes flavidus*, off southern California, Love et al. 1990). Black rockfish, like other rockfishes, are primitive livebearers and normally produce a single batch of larvae annually. Parturition dates were estimated from advanced stage gonads sampled during the pre-parturition period (Bobko 2002). These data were applied to the age distribution of the population to estimate the percentage of larval production by age class for each week. Birthdates of young of the year benthic juveniles were then determined from daily growth rings on their otoliths, from which estimates were made of the relative contribution to recruitment by time period, and thus by implication, age group. By comparing birthdate distributions to adult spawning output, Bobko (2002) determined whether recruitment was proportional to spawning output or whether certain periods during the parturition season were responsible for a greater proportion of recruits. Results indicated that, in 1 of 3 years, significantly greater recruitment came from earlier in the spawning season, a time when few young fish were spawning.

Timing of spawning is likely to be a highly conserved trait in most fishes, as larval survival is highly dependent on larval production coinciding with peak zooplankton production (i.e., the “match-mismatch hypothesis” of Cushing 1969,1975). For fishes that exhibit age-related temporal patterns of spawning, elimination of older age classes through fishing will effectively shorten the spawning season. In those years when successful recruitment is centered early in the season, elimination of older age classes could result in recruitment failure that would otherwise be avoidable if the age structure was intact. Indeed, Marteinsdottir and Thorarinnsson (1998) found that strong year classes of Icelandic cod (*Gadus morhua*) occurred only when the population contained a broad age distribution, suggesting that this relationship may be applicable to a variety of species.”

Importance of Heterogeneity and Recruitment. In view of the factors discussed above, the following from Berkeley et al. (2004:24) is also relevant to Puget Sound Proper rockfish: “Although marine populations are obviously affected by the vagaries of larval survival (Houde 1987), the spatial and temporal features of year-class formation are not yet clearly understood. Hedgecock (1994a,b) proposed the “sweepstakes hypothesis” to explain small-scale genetic heterogeneity observed in some widely distributed marine populations. According to this hypothesis, most spawners fail to produce surviving offspring because their reproductive activity is not matched in space and time to favorable oceanographic conditions for larval survival during a given season. As a result of this mismatch (sensu Cushing 1969, 1975), the surviving year class of new recruits is produced by only a small minority of adults that spawned within those restricted temporal and spatial oceanographic windows that offered good conditions for larval survival and subsequent recruitment.”

Berkeley et al. (2004:24,25) continue: “One testable prediction of Hedgecock’s hypothesis is that a cohort of new recruits would show less genetic diversity than the adult population, reflecting the underlying pattern of only a few adults successfully passing their genome to each new cohort. This hypothesis was tested during 1994, when the National Marine Fisheries Service Tiburon Lab conducted a sampling survey of the entire pelagic stage of shortbelly rockfish (*Sebastes jordani*) combined with juvenile and adult surveys (Larson et al. 1995; Julian 1996). Results supported the Hedgecock hypothesis (Table 2): levels of genetic diversity were lower for later stage (“June” and “Farallon”) pelagic juveniles and their haplotype frequencies were different from both adults and most earlier stages of larvae and juveniles. Results also indicated that these differences were not due to seasonal spawning by a unique portion of the adult population, but were in fact the result of differential survival during the pelagic larval stage. In contrast, Gilbert (2000) found no reduction in genetic diversity within a very strong year class of kelp rockfish (*S. atrovirens*). Burford (2001) also found no reduction in genetic diversity in a year class of blue rockfish (*S. mystinus*) but did find genetic differences among recruits from different locations that were not matched by differences among adult populations.

Gomez-Uchida and Banks (in press), studying the population genetics of darkblotched rockfish (*S. crameri*), found that the breeding population was several orders of magnitude smaller than the spawning stock size indicated by the stock assessment. Only several thousand breeders, rather than the millions of adults in the whole population, would successfully reproduce. While not a direct test of Hedgecock’s hypothesis, these results nevertheless suggest that recruitment is not uniformly distributed throughout the adult population.

Thus, although the generality of Hedgecock’s hypothesis remains in question, indications are that it may be true under at least some circumstances, and that the genetic composition of recruits may otherwise be quite complicated spatially. This suggests that the geographic source of successful recruits may differ from year to year. Based on these observations, management should strive to preserve a minimal spawning biomass throughout the geographic range of the stock (Larson and Julian 1999).”

The Bottom Line - Avoiding Age Class Truncation. In aggregate, all of the previous sections indicate that the age class structure created in Puget Sound Proper

rockfish populations during thousands of years of evolution should have been maintained. However, severe truncation has already occurred due to fishing. The nature of the problem is described by Berkeley et al. (2004:25): “Many marine species, such as those in the north temperate waters of the Pacific Ocean, exhibit long life spans, with the rockfishes being particularly striking in this regard (Table 3). The adaptive value of allocating reproductive output across many years (iteroparity) is generally thought to be a bet-hedging strategy that ensures some individual reproductive success despite long periods of environmental conditions unfavorable for larval survival (Leaman and Beamish 1984). At the population level, longevity ensures that there will be sufficient reproductive output for the population to maintain itself between favorable recruitment events (Longhurst 1999, 2002).

Age truncation - the removal of older age classes via fishing - occurs at even moderate levels of exploitation (Figure 1). Leaman and Beamish (1984) suggest that age truncation will be most detrimental when reproductive success is highly variable, since stock maintenance may be dependent on the relative stability of reproductive output that results from a broad spectrum of age classes. Most, if not all rockfish stocks on the U.S. west coast fall into this category of highly variable and episodic recruitment (Hollowed et al. 1987; Moser et al. 2000).”

Berkeley et al. (2004:26) continue on the same subject: “While the advantage of longevity for persistence of a population in a variable environment is intuitive, a growing body of evidence suggests that a broad age distribution can also reduce recruitment variability (Lambert 1990; Marteinsdottir and Thorarinsson 1998; Secor 2000a,b). There are at least two mechanisms by which stabilization of recruitment could occur: (1) there may be age-related differences in the time and location of spawning (Berkeley and Houde 1978; Lambert 1987; Hutchings and Myers 1993), thereby spreading larval production to cover temporal and spatial variability in favorable environmental conditions, and (2) older fish may produce larger, healthier, or otherwise more fit larvae (Hislop 1988; Marteinsdottir and Steinarsson 1998), which may survive under conditions that are inadequate for the survival of progeny from younger fish. Even slightly enhanced rates of early survival and growth have a cumulative effect that can translate into a greatly increased probability of successful recruitment (Houde 1987).

Long life spans are necessarily associated with low rates of mortality during the adult stage (Stearns 1992). For most teleosts, size-dependent processes, especially lower risk of predation with increasing body size, result in decreasing natural mortality as fish grow older and larger (Hare and Cowan 1997; Sogard 1997). Fishing, however, generally selects for larger, older fish, imposing a selection pressure that works opposite to that of most natural mortality agents. One of the most predictable effects of fishing is the reduction or removal of the older age classes, i.e., age truncation (Figure 1). Prior to reaching a size acceptable to the fishery (either due to market demand or minimum size regulations intended to allow fish to attain reproductive maturity), younger age classes are reduced only through natural mortality, but once vulnerable to fishing gear, a cohort is reduced through both natural and fishing mortality. These sources of mortality are cumulative throughout the life of the cohort.”

Fishing Induced Genetic Change. A subpart of the age class truncation problem described above is the high risk of induced genetic change. Berkeley et al. (2004:29)

describe this risk as follows: “As shown above, fishing generally results in age truncation of the population. It has long been recognized that fishing, by removing the largest and oldest fish inadvertently removes fish that are genetically predisposed to fast growth and late maturation, creating a selection pressure that should theoretically favor early-maturing, slow-growing individuals. Nevertheless, this question has been largely ignored in fisheries management, which tacitly assumes that exploited populations will always retain their inherent rates of productivity and tend to return to their previous levels of abundance. A recent paper by Olsen et al. (2004) challenges these long-held beliefs. Results of this study strongly suggest that heavy and continuous fishing pressure in northern cod resulted in a rapidly-evolved, genetically-based shift in maturation patterns towards earlier and smaller sizes at maturity. The implications of this finding for management are profound. If evolutionary change in response to fishing turns out to be the rule rather than the exception, then, as Hutchings (2004) observed, we must address issues of reversibility of these changes, and their consequences for sustainable harvesting, population recovery and species persistence. At the very least, we believe that these results suggest that some portion of the population should be protected from the impacts of fishing, providing a sanctuary for the genes of fast-growing, late-maturing individuals.”

NOAA Fisheries Challenge to the Importance of Longevity

The importance of maintaining longevity in Puget Sound Proper rockfish populations was recently disputed (Varanasi 2006) but this challenge was limited to a single element of the many facets presented by Berkeley et al. (2004). Instead of correctly focusing on the Berkeley et al. (2004) feature article from the August 2004 issue of *Fisheries*, Varanasi (2006), perhaps inadvertently, addressed the following paper from the same year with the same lead author (S.A. Berkeley, 2004): Berkeley, S.A., Chapman, C., and Sogard, S.M. 2004. Maternal age as a determinant of larval growth and survival in a marine fish, *Sebastes melanops*. *Ecology*, 85:1258-1264. In a recent population modeling exercise, it was found that black rockfish larvae which survived at a low rate during their first 10 days came from a relatively minor proportion of the overall population of mature females (O’Farrell, M.R., and L.W. Botsford. 2006. The fisheries management implications of maternal-age-dependent larval survival. *Can. J. Fish. Aquat. Sci.* 63:2249-2258).

However, O’Farrell and Botsford (2006) also stated the following: “Results from two studies demonstrate that if viable larvae are instead used as the spawning metric in the stock-recruitment relationship, the level to which per-recruit reproduction can be reduced without causing collapse is increased. Murawski et al. (2001), using viable larvae as the spawning metric for the Georges Bank cod stock-recruitment relationship, found that the percentage of unfished viable larvae necessary for population persistence was lower if SSB were used as the spawning metric. Similarly, another study (Spencer et al. 2007) found that resiliency of the stock-recruitment curve based on Gulf of Alaska Pacific ocean perch (*Sebastes alutus*) was increased if viable larvae were used as the spawning metric.”

Varanasi (2006) acknowledged the relationship for cod but dismissed it because black rockfish were much more closely related to Puget Sound Proper rockfish DPSs,

including a possible DPS for black rockfish. The relationship for Pacific ocean perch was not mentioned by Varanasi (2006). In reality, the importance of low survival rates from the progeny of small females is not dependent upon how closely related study species are to populations potentially at-risk. The important parameter is simply the percentage of the spawning population composed of smaller females as demonstrated by O'Farrell and Botsford (2006). When Palsson and Pacunski (1998) estimated that the reproductive potential of rockfish in a marine reserve exceeded the potential of the average fished site in Puget Sound by 55 to one, they did not even try to adjust for any possible relationship between female size and larval survival rate. However, there must have been a substantial component of small females at the fished sites in order to produce the 55 to one disparity. The above comparison was followed-up by a report which dramatically illustrates the differences between protected and fished areas for Puget Sound Proper rockfish in terms of numbers of fish per acre, length frequency distributions and relative number of eggs per acre (Palsson, W. 2001. Marine refuges offer haven for Puget Sound fish. Washington Department of Fish and Wildlife, Olympia, WA.).

Justification for Listing Bocaccio

Both Wright (1999) and Stout et al. (2001) were remiss in that they did not examine one data source that was relevant to the status of Puget Sound rockfish. There are 12 years of catch estimates for the Puget Sound recreational fisheries from 1975 through 1986 that breakdown part of the rockfish catches by species. These estimates were based on relatively low sampling percentages (as compared to coastal fisheries), had large increments of unidentified rockfish, and were dependent on expansion factors from the recreational salmon fishery. The estimates do not include catch by divers, shore and pier anglers. Catch estimates since 1981 bear the following note: "the data for catch areas 5-13 and the freshwater areas were adjusted downward (by multiplying statistics generated by the usual punch card methodology by 0.833) to correct for the assumed bias of 20% in the punch card methodology. Comparisons with sport fishery data before 1981, which have not been corrected, must therefore take this adjustment into account." Nevertheless, the 12 years of sport catch records do provide some valuable quantitative measures of the presence and relative abundance of the individual species. For the five species addressed in this Petition, there are no valid fishery independent abundance estimates, past or present, and there will not be any for the foreseeable future. "Valid" in this case means a lack of any numbers from WDFW surveys that are statistically significant. Trawl surveys simply cannot produce valid population estimates for rockfish species with a low abundance since there are far too many "0" data entries. With rocky reef habitat constituting less than two percent of overall habitat in Puget Sound Proper, any rockfish encounters depend on whether or not the trawl gear happens to pass over or near a "rock pile". The WDFW reported estimated catches of bocaccio taken in Puget Sound Proper (corresponding to Catch Areas 8 through 13) were as follows:

1975 - 675

1976 - 1973

1977 - 9275

1978 - 4915

1979 - 1836

1980 - 1081
1981 - 605
1982 - 1418
1983 - 759
1984 - 9
1985 - 0
1986 - 344

A definite downward trend is evident and bocaccio had been reduced to a very low abundance by the mid-1980s. Of the 22890 total estimated catch of bocaccio, 18742 or 82% were taken from the Tacoma Narrows southward (Catch Area 13). Their abundance coincided with the former location of an intense fishery for walleye pollock (*Theragra chalcogramma*). This extreme southerly abundance concentration in Puget Sound Proper makes it obvious that this was not some small transient "fringe" of a much larger population of bocaccio somewhere to the north. The actual catch during this period probably exceeded 50000 fish due to the catch exclusions and large numbers of unidentified rockfish. WDFW biologist Greg Bargmann recently advised the Petitioner that bocaccio have not been observed in any Puget Sound Proper resource surveys for about 20 years. He also related that all of the individuals he personally observed were large fish, indicating a prolonged period of recruitment failure. Bocaccio were one of 13 rockfish species designated as "important species of bottomfish in Puget Sound" by Palsson et al. (1997). This same "important" status was soon reiterated in the Puget Sound Groundfish Management Plan (Bargmann 1998).

Due to the extreme longevity potential of bocaccio, it is unlikely that the species has been completely extirpated from Puget Sound Proper. The WDFW surveys do not cover many rocky reef habitats, and it is likely that a few bocaccio have persisted in isolated pockets of distribution.

Justification for Listing Canary Rockfish and Yelloweye Rockfish

The twelve years of WDFW Catch Records from 1975 through 1986 prove that canary rockfish and yelloweye rockfish were once significant components of the Puget Sound Proper rockfish community. Catches of canary rockfish were reported in all 12 years, ranging from a low of 288 fish in 1983 to a high of 2571 fish in 1986 and averaging 1301 fish per year. Unlike bocaccio, there was no discernible trend in catches over the time period and catches were spread throughout Catch Areas 8 through 13 (Puget Sound Proper) instead of being concentrated far to the south in Area 13. The total reported canary rockfish catch of 15608 probably equates to an actual catch of about 40000 due to the catch exclusions described previously and large quantities of unidentified rockfish.

Yelloweye rockfish appeared to be somewhat less abundant than canary rockfish in Puget Sound Proper but catches were still reported in 11 of the 12 years of record. No catch was reported in 1979, the high was 1876 fish in 1986 and the average annual reported catch was 730 fish. Like canary rockfish, there was no discernible trend in catches over the 12 year period and the fish were present throughout Catch Areas 8 through 13. The total 12 year reported catch of 8761 yelloweye rockfish in Puget Sound

Proper probably equates to an actual catch of about 25000 due to catch exclusions and unidentified rockfish.

Canary rockfish and yelloweye rockfish were two of the 13 rockfish species described as “important species of bottomfish in Puget Sound” by Palsso et al. (1997). The same “important” rating was carried forward in the Puget Sound Groundfish Management Plan (Bargmann 1998). Between 1986, the last year of catch records (and the 12 year high for both species) and 2003, there was a significant decline in abundance for both species. Early 2003 marked the beginning of the commonly applied expression of “virtually disappeared” to both species. The WDFW justification statement for a proposed ban on the retention of both species in Puget Sound reads as follows: “Canary and yelloweye rockfish are two species of rockfish which are orange in color. Both are commonly called “red snapper” by anglers, and are long-lived species and may reach ages in excess of fifty years. While never common in Puget Sound, both species have virtually disappeared from the sport harvest of sport anglers throughout Puget Sound. The department has no estimates of the abundance of these two species in Puget Sound. However, anecdotal evidence indicates that both canary and yelloweye rockfish have declined in abundance. Both species have declined in abundance in Washington coastal areas and these declines have resulted in severe restrictions on fishing along the coast.” (WDFW. 2003. Concise explanatory statement regarding 2003-2004 sportfishing rule proposals.)

The 12 years of WDFW catch records from 1975 through 1986 provide much more than “anecdotal evidence” of a decline in abundance by 2003 when both species had “virtually disappeared”. It is true that WDFW had no fishery-independent abundance estimates for the two species in 2003, does not have any now and will not have any in the foreseeable future. Unlike bocaccio, small numbers of the two species are still seen during surveys but the numbers are far too low to produce a valid abundance estimate. In addition, canary rockfish sustained the largest decline of any species in their relative abundance ranking within the rockfish assemblage. In the early 1960s, canary rockfish were identified as one of three “principal species” taken by commercial trawling in Puget Sound along with copper and quillback rockfish (Holmberg, E.K., D.Day, N.Pasquale, and B.Pattie. 1967. Research report on the Washington trawl fishery, 1962-1964. Washington Department of Fisheries, Olympia, WA.).

Justification for Listing Greenstriped Rockfish

As was the case for the previous three rockfish species, WDFW has no valid fishery independent abundance assessment of greenstriped rockfish in Puget Sound Proper and the 12 years of catch estimates from 1975 through 1986 are the only quantitative measures available (past, present or future) to prove that they once existed in significant abundance. Greenstriped rockfish are yet another “important species of bottomfish in Puget Sound” (Palsso et al. 1997, Bargmann 1998). The WDFW reported catch estimates of greenstriped rockfish in Catch Areas 8 through 13 were as follows:

1975 - 1046
1976 - 1870
1977 - 720
1978 - 939

1979 - 2416
1980 - 932
1981 - 238
1982 - 1360
1983 - 180
1984 - 261
1985 - 0
1986 - 0

As was the case with bocaccio, a definite downward trend in catches is evident, with no catch being reported in the two most recent years of records, 1985 and 1986. It is also clear that greenstriped rockfish are well isolated from other members of the species since sport catches were only reported from Catch Areas north of Puget Sound Proper in two of the 12 years - 48 fish and 80 fish in 1979 and 1980, respectively. Catches were distributed throughout Areas 8 through 13, unlike bocaccio, which were concentrated to the south in Area 13. The reported catch in 12 years of 9962 greenstriped rockfish falls slightly above the total for yelloweye rockfish (8761) and probably equates to a total catch of about 25000 due to catch exclusions and unidentified rockfish. WDFW biologist Greg Bargmann recently advised the Petitioner that greenstriped rockfish have been observed in deep water surveys off Blake Island in Puget Sound Proper. However, it is very likely that both their abundance and extent of distribution are severely constrained at the present time.

Justification for Listing Redstripe Rockfish

There are no valid fishery independent abundance estimates available from WDFW for redstripe rockfish in Puget Sound Proper - past, present or future. As was the case for the previous four species, the only quantitative measure of their past abundance comes from the 12 years of WDFW Catch Records from 1975 through 1986. In spite of this, redstripe rockfish are yet another "important species of bottomfish in Puget Sound" (Palsson et al. 1997, Bargmann 1998). The WDFW reported catch estimates for Catch Areas 8 through 13 are as follows:

1975 - 211
1976 - 10
1977 - 290
1978 - 15
1979 - 635
1980 - 4314
1981 - 1247
1982 - 5139
1983 - 11451
1984 - 264
1985 - 162
1986 - 0

Catches were small during the first four years (1975-1978), began increasing to reach a peak in 1983, and then declined to low levels in the final three years of catch records. As was the case with bocaccio, redstripe rockfish had their greatest abundance in

the same general area as an intense fishery for walleye pollock. Of the 23738 individuals reported in the 12 year period, 17125 or 72% came from the Tacoma Narrows southward (Catch Area 13). As was the case for greenstriped rockfish, redstripe rockfish in Puget Sound Proper are isolated from other members of their species. Catches were only reported from Catch Areas north of Puget Sound Proper in four of the 12 years of records and all of these were small - 80 fish, 248 fish, 268 fish and 68 fish in 1980, 1982, 1983 and 1984, respectively. The total catch in Puget Sound Proper probably exceeded 50000 fish due to large quantities of unidentified rockfish and catch exclusions. Redstripe rockfish are still seen during WDFW surveys in Puget Sound Proper but their numbers are too small to permit any valid estimate of population abundance.

Resource Management by Washington Department of Fish and Wildlife

In response to declining resource abundance, there have been a number of recent changes in rockfish management for recreational fishing in Puget Sound Proper but only two qualify as partial but legitimate conservation regulations. In any open access fishery where there is no direct control of fishing mortality (fishing effort), the only legitimate conservation regulations are those that apply to *every individual fish* (Hunt, R.L. 1970. A compendium of research on angling regulations for brook trout conducted at Lawrence Creek, Wisconsin. Wisconsin Department of Natural Resources, Research Report 54, Madison.). In 2003, the retention of yelloweye and canary rockfish was banned but hooking mortality continued as a significant cause of fishing mortality. There is no way to estimate impacts since the abundance of both species is unknown. However, a WDFW statement about a comparable situation off the Washington Coast reads as follows: "In an effort to rebuild the population, a recreational harvest limit of three metric tons annually has been set. Fisheries managers estimate that limit is reached just in the number of rockfish that die after being unintentionally harvested and discarded." (WDFW News Release, May 1, 2002). In 2004, spear fishing for rockfish was closed and recreational angling for rockfish was restricted to times when fishing was allowed for lingcod and/or Pacific salmon. However, any possible "savings" from these changes was probably lost because there was no direct control of fishing mortality (effort) during remaining open fishing periods. At best, the rates of population declines have been slowed down.

The main recent change that was touted by WDFW as a conservation regulation was the reduction to a one rockfish daily bag limit in 2000. However, daily bag limit reductions, by themselves, have never been a legitimate conservation regulation in open access fisheries since they violate the fundamental rule of not applying to *every individual fish*. There have been countless failures with this approach but these failures have rarely been documented. The largest documented percentage reduction failure that the Petitioner is aware of (80%) occurred when Idaho reduced their daily bag limit on trout from 15 fish all the way down to three fish but could not detect any positive response in the trout populations (Johnson, T.H., and T.C. Bjornn. 1978. Evaluation of angling regulations in the management of cutthroat trout. Idaho Cooperative Fishery Research Unit, University of Idaho, Job Completion reports F-59-R-7, F-50-R-8, Moscow.) Bag limits have a legitimate place in recreational fishery management, but only as a means of distributing the allowable catch among more participants, thus optimizing the recreation benefits that can be derived from a given fishery.

Two variables that WDFW failed to factor into the restrictions described above are (1) that most anglers will fish for boat or group limits rather than individual limits, and this technically illegal action cannot be controlled; and (2) that temporary harvest reductions are often canceled out by a significant degree of increased fishing effort at a later time, as well as higher catch per unit of effort (both in response to higher-than-expected fish population levels) (Wright, S. 1992. Guidelines for selecting regulations to manage open-access fisheries for natural populations of anadromous and resident trout in stream habitats. *North American Journal of Fisheries Management* 12:517-527).

It is obvious to concerned members of the public that the rockfish resources of Puget Sound Proper are in trouble. The fact of extensive catch sorting and its associated hooking mortality is now common knowledge, caused by declining size of the fish and exacerbated by a one fish daily bag limit. The recreational fishery has become very effective with the widespread use of electronic "fish finders" and heavy "down-rigger" gear and this has further exacerbated an already tenuous situation. These concerns on behalf of the public prompted then-Governor Locke to direct WDFW to produce a resource assessment and management plan for Puget Sound rockfish resources by December 31, 2004. This assignment was never completed. There was apparently no follow-up demands from the new Governor even though the same political party controlled the office and many of the same staff members were retained. Staff work has continued within WDFW but a final product has yet to emerge from the continued recycling of various drafts through multiple layers of review.

It is easy to deduce the cause for this delay. It is common knowledge that WDFW has evolved into a very harvest oriented agency, especially with respect to recreational fisheries. They have stated publicly that they want to prevent any further ESA listings in Puget Sound. They also have a firm policy of not releasing any newer resource assessment data to the public until it is an integral part of a finalized resource management plan. However, as noted previously, there will not be any new resource assessments for the five species that are the subject of this Petition - there is nothing to "wait for". The obvious conflict involved is adjacent or comingled populations of rockfish and Chinook salmon. There is no conceivable successful rockfish management regime that would not have some significant impacts on Chinook salmon fishing opportunities. It is certainly ironic that the main conflict involved is with ESA listed Puget Sound Chinook.

WDFW has requested special funding from the State Legislature to pay for rockfish research, specifically stock assessment. House Bill 1076 describes how this funding will come from short-term (2008-2010) surcharges on commercial fishing licenses that provide for the retention of groundfish species and on recreational saltwater fishing licenses. The justification is that "For many of these stocks there have been no recent stock assessments, or the current assessments are based on poor data." However, it is anticipated that priority will be given to coastal surveys due to presence of a large, economically valuable charter boat fishery and an active fishery management regime by the Pacific Fishery Management Council in which fisheries might be constrained at any time by rockfish conservation concerns. In Puget Sound, there is no outside agency pressure on WDFW to do anything to change the status quo, especially in light of the "poor data".

Assessment of Viability

The viability of Puget Sound Proper rockfish DPSs has been severely compromised by past and current fisheries management practices, especially the resultant severe declines in abundance and truncation of age classes. The key elements that makes management of rockfish different than all other fish populations is *their extreme longevity and their inability to adjust to hydrostatic changes when brought quickly to the surface*. All rockfish have a swimbladder and typically do not survive due to the internal trauma of expansion and rupture of the swimbladder during capture. WDFW has recently announced their intention to increase salmon fishing opportunities in Puget Sound Proper in the near future through the expanded use of selective fisheries. While selective fishing is a viable management alternative for salmon, it is a disaster for rockfish. Fishing mortality will significantly increase due to incidental rockfish catches during salmon fishing and due to the increased targeted fishing opportunities for rockfish that are provided whenever salmon fishing is allowed.

The annual releases of millions of hatchery chinook will continue unabated. Millions of new predators will quickly reach marine waters near small stream systems that never had significant populations of wild spawners. In addition, all of the hatchery chinook fingerlings are now significantly larger than their naturally produced counterparts. This greatly increases the size range of juvenile rockfish that can be consumed at any given point in time.

This is a unique marine area ecosystem with a high retention time, restricted hydrologic exchange and a net outflow of surface waters in which numerous closely related rockfish species must rely upon a meager 14 square kilometers of suitable rocky reef habitat. Due to habitat partitioning, each species will only have available a fraction of this already small number. The actual situation is even worse than it seems since Palsson and Pacunski (1998:13) reported that "Many vacant habitats have been observed throughout Puget Sound during VAT surveys." (VAT is the acronym for Video-Acoustic Technique). Currently, the only relatively abundant species in Puget Sound Proper are brown, quillback and copper rockfish. Even with these species, evidence shows that genetic diversity is relatively low, indicating that Puget Sound Proper rockfish populations experienced a post-glacial founder effect followed by genetic isolation and low effective population sizes. The three common species also appear to have recent mixed ancestry as a result of introgression with each other (Buonaccorsi, V.P., C.A. Kimbrell, E.A. Lynn, and R.D. Vetter. 2005. Limited realized dispersal and introgressive hybridization influence genetic structure and conservation strategies for brown rockfish, *Sebastes auriculatus*. Cons. Genetics 6:697-713.). For the five species addressed in this Petition that are far less abundant at the present time, it is highly likely that their genetic integrity has already been severely compromised by introgressive hybridization.

Sport catch estimates made by WDFW from 1975 through 1986 demonstrate that bocaccio in Puget Sound Proper probably had a minimum population size of at least 50000 individuals in the recent past. Unfortunately, the distribution of bocaccio was mainly from the Tacoma Narrows southward (Catch Area 13), the former site of an intense fishery for walleye pollock - which subsequently disappeared from Puget Sound

Proper. Bocaccio have not been seen during WDFW surveys for the past two decades. They may have been completely extirpated from Puget Sound Proper and, at a minimum, are well past any onset of depensatory mortality factors. At best, one or more viable populations may still exist in the many rocky reef habitats not covered in WDFW surveys.

The same sport catch estimates demonstrate that canary rockfish, yelloweye rockfish, greenstriped rockfish and redstripe rockfish probably had minimum population sizes of at least 40000, 25000, 25000, and 50000 individuals, respectively, in the recent past. All four species are still observed during WDFW surveys but the numbers are far too small to make any type of valid fishery independent abundance estimates. All four species are probably approaching or may have already reached the onset of depensatory mortality factors. Disjunct geographic ranges are an additional critical at-risk factor for all five species of rockfish. Members of each species range from rare to non-existent in the adjacent marine waters of North Puget Sound.

In the early 1960s, canary rockfish were identified as one of three "principal species" taken by trawling in Puget Sound waters along with copper and quillback rockfish (Holmberg et al. 1967). The WDFW survey data and past sport catch statistics demonstrate that Puget Sound Proper had a rockfish assemblage of ten relatively abundant species by the mid-1970s. Brown, copper and quillback rockfish were the most abundant species three decades ago and are still the most common species at the present time. Yellowtail and black rockfish ranked next in order of mid-1970s abundance and still rank at that point today despite marked declines in abundance. These are pelagic, schooling species that are more resistant to extirpation threats in unique "rockfish isolated" Puget Sound Proper because they can move in from other areas as adults. They have wider home ranges and make longer movements than the sedentary species. However, the ten species rockfish assemblage of the recent past is well on its way to becoming cut in half to a five species assemblage. All five sedentary species addressed in this Petition cannot move in from other areas as adults. One species, bocaccio, may already be extinct. Two species, yelloweye and canary, are now commonly referred to as "virtually disappeared". Two other species, greenstriped and redstripe, were already declining in abundance by the mid-1980s in Puget Sound Proper and were rare in the adjacent marine waters of North Puget Sound from at least the mid-1970s.

In view of the facts cited and presented in this Petition, *the Puget Sound Proper populations of bocaccio, canary rockfish, yelloweye rockfish, greenstriped rockfish and redstripe rockfish are in danger of extinction throughout all or a significant portion of their range or are likely to become so in the foreseeable future.*

**REQUEST FOR REEXAMINATION OF YOUR REJECTION DECISION
RELATIVE TO THE APRIL 2007 PETITION TO LIST FIVE SPECIES OF
PUGET SOUND ROCKFISH AS THREATENED OR ENDANGERED**

TO: SECRETARY OF COMMERCE, UNITED STATES DEPARTMENT OF
COMMERCE, NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION,
NATIONAL MARINE FISHERIES SERVICE (NMFS).

From: Sam Wright (Petitioner)

Subject: On October 1, 2007, NMFS rejected the following: Petition to list five species of rockfish in their Puget Sound Proper Distinct Population Segments as Endangered or Threatened under the Endangered Species Act (ESA). Intent of the narrative to follow is provision of additional information that will facilitate a reexamination of that decision.

Special Case for Bocaccio

The current situation for one species, bocaccio, is very different from the other four species covered in the Petition - at least they still probably exist in Puget Sound Proper. NMFS was provided with information demonstrating that published Washington Department of Fisheries (WDF) catch statistics showed that a total landed catch of 22,890 bocaccio was made in the 12 year period, 1975-1986. NMFS was also provided with information showing that this was only part of the actual total catch since catches by shore and pier anglers were excluded as well as the take by divers. In addition, many rockfish were not classified to species since WDF samplers encountered some "hybrids" in Puget Sound and many rockfish were reduced to fillets onboard vessels before catches were examined by samplers. In the Petition, it was estimated that at least 50,000 bocaccio were probably taken in this same period.

NMFS was also provided with information from a qualified expert, WDFW Biologist Greg Bargmann, that bocaccio have not been observed in Puget Sound Proper for the past 20 years. I am sure that he told you exactly the same thing when you interviewed him relative to content of the Petition. This type of information was incorrectly labeled as "anecdotal" and "weak" by NMFS and generally ignored. Zero is an unambiguous *quantitative* expression. Based on this information, I was sure that NMFS would, at a minimum, agree to conduct a status review of bocaccio. A fish population decline from 50,000 to zero should have been more than adequate proof of a legitimate problem. This status review would have included the bocaccio population in the adjacent waters of North Puget Sound. Due to the rockfish population isolation factors recognized by NMFS, these are the only fish that would have even a remote chance of providing progeny to Puget Sound Proper during the rare favorable recruitment event that is characteristic of rockfish populations.

I had envisioned the eventual establishment of one or more marine protected areas if any significant bocaccio populations were located in North Puget Sound, especially if they were relatively close to Puget Sound Proper. I had also envisioned the establishment of several marine protected areas in Puget Sound Proper to protect depleted rockfish

populations from further exploitation. None of this will ever happen under management by the State of Washington.

The Puget Sound Proper Recreation Fishery for Rockfish, 1975-1986.

The primary reason that NMFS rejected the Petition at the policy level was their blanket refusal to use catch data in any manner as measures of population abundance, even though these were the only quantitative fish population data available, fishery-independent or otherwise. Further, no additional usable quantitative information is expected to become available in the foreseeable future. The populations are either gone, as is the case with bocaccio, or persist at very low numbers that cannot be quantified by any known resource assessment techniques. I am fully cognizant of the limitations in using catch data as measures of population abundance and the numerous resource management errors that this has caused in the past. However, the most common error, by far, is that of catch data causing a manager to overestimate population abundance. This is what happened in Puget Sound Proper.

During the 12 year period, 1975-1986, recreational fishery catches in Puget Sound Proper (as reported in published WDF statistics) were dominated by four species. The 12 year totals in order of importance were 406,718 copper rockfish, 354,466 quillback rockfish, 246,225 brown rockfish, and 86,550 black rockfish. None of these top four species showed a discernible trend in catch over time. Yellowtail rockfish ranked as 5th, with an 11 year total of 66,865 (WDF statistics did not have a 1980 entry for this species in the entire state). This species showed a moderate downward trend over time, but yellowtail rockfish are highly migratory and the decline could have been caused by movement out of Puget Sound Proper and/or a decline in population abundance.

The next five species were those addressed in the Petition: 23,738 redstripe rockfish, 22,890 bocaccio, 15,608 canary rockfish, 9,962 greenstriped rockfish, and 8,761 yelloweye rockfish. As noted in the Petition, bocaccio, redstripe rockfish, and greenstriped rockfish all showed pronounced downward trends in catch over time. All three are non-migratory so that there is no possible alternative explanation for a decline in abundance.

To present the complete statistics, there were 12 year totals for the following: 293 widow rockfish, 275 China rockfish, 108 silvergrey rockfish, 39 redbanded rockfish, 19 blue rockfish, and 9 vermilion rockfish (Note: there was no Area 13 in 1975. It was part of Area 11 until 1976).

During the 12 year period, there were no significant regulation changes in the fishery that could have influenced levels of catches. However, there were a number of changes in the fishery itself and all of these should have produced increased catches, both singly and in aggregate. Prior to the Boldt Decision in 1974, the marine area fishery was primarily a salmon fishery, with marine species taken as bycatch and often discarded. After 1974, targeted fishing for marine species was heavily promoted as an alternative to salmon fishing and most of the catch was retained. In addition, the fishery evolved to the widespread use of electronic fish finders and down rigger gear. This allowed anglers to fish at much greater depths and to place their terminal gear at the same depths as the fish. It also allowed effective fishing throughout the tidal cycle instead of being limited to slack tide periods. Thus, large fish populations in deeper waters became available to

anglers. These had been relatively immune to exploitation in the past simply because of the depths they occupied.

The changes described above should have produced a trend of increased catches over time and this seemed to be the case in the first few years of the fishery after 1974. However, the nine non-migratory species showed six stable catch trends over the 12 year period and three declining trends. The reason becomes obvious when the average rockfish catch per trip data over time is examined. These data were provided to NMFS in the 1999 Petition and showed a distinct downward trend over time (see Figure 1. Recreational catch rates of rockfish (catch per trip) of bottomfish anglers in North and South Puget from Palsson and Pacunski (1998). Note that the declining trend continued for over a decade after 1986.). The only possible interpretation is that the increased effectiveness of the fishery (including the elimination of discards by turning them into "catch statistics") actually masked real population declines in the six species showing stable catch trends over time - including the canary rockfish and yelloweye rockfish included in the Petition. These two species simply could not sustain the same catch levels subsequent to 1986. The decline may have been largely in place by 1999 since the 1999 Petition stated (in the context of Greater Puget Sound) that "At the present time, only five species are commonly caught in the commercial and sport fisheries. The others have largely disappeared from catch samples and are rarely seen during fishery-independent stock assessments. The most noticeable declines have been with tiger, canary and yelloweye rockfish." This shows how these two species could easily go from apparently stable catch trends through 1986 to Greg Bargmann's 2004 expert opinion of "virtually disappeared" in the relatively long time span from 1986 to 2004. It also means that there is no plausible alternative explanation for the three species in the Petition showing declining catches. The inferred severe declines were real. The "virtually disappeared" expression is not "anecdotal" and "weak" as described by NMFS. It is a valid *qualitative* statement by a qualified professional and the only type of assessment possible when fish populations become too small for quantitative assessment techniques. I am sure that he told NMFS exactly the same thing when they interviewed him on content of the Petition. Mr. Bargmann has worked with Puget Sound marine fish on a daily basis for many years. He is very familiar with every bit of information available for each species, including results from catch sampling and fishery-independent resource assessment surveys. "Virtually disappeared" ranks only slightly better than "have disappeared" (bocaccio) and should have been more than enough to merit a status review by NMFS.

The North Puget Sound Recreational Fishery (Areas 5, 6 and 7), 1975-1986

This fishery also had no significant regulation changes in the time period shown and had an increased use of electronic fish finders and down rigger fishing gear. However, it did not share the other changes described for Puget Sound Proper. There was a much better balance prior to 1974 between salmon fishing and targeted fishing for marine species. Discards were minimal, at least for rockfish. The salmon present in most of the area were predominately of Canadian origin, thus there was little pressure or interest in curtailing salmon fishing. Active promotion of fishing for marine species as an alternative to salmon fishing was minimal. North Puget Sound was much more of a

seasonal fishery and had fewer population centers immediately adjacent to the fishing areas.

North Puget Sound has a much greater population of rockfish in terms of all species combined, a reflection of the 200 plus square kilometers of rocky reef habitat available (versus 14 in Puget Sound Proper). The multiple population isolating factors of Puget Sound Proper are absent. Total rockfish catch was less than that of Puget Sound Proper. Thus, the recreational fishery in North Puget Sound (at least minus divers) had a much lower exploitation rate on rockfish populations (Note: the catch per trip shows a declining trend but was consistently higher in North Puget Sound. See Figure 1 from Palsson and Pacunski (1998) in the 1999 Petition). However, the Puget Sound commercial trawl fishery has always been centered in North Puget Sound and continued in this area after trawling was banned in Puget Sound Proper. The relatively clear waters on the west side of San Juan Island were once famous for their "meat trips" by divers. Unfortunately, no one ever tried to estimate this harvest in Greater Puget Sound.

The following shows the 12 year total catches of rockfish by species in North Puget Sound, with the comparable Puget Sound Proper catches in parenthesis:

- 308,729 quillback rockfish (354,466) ranks 1st in NPS, 2nd in PSP
- 278,134 copper rockfish (406,718) ranks 2nd in NPS, 1st in PSP
- 160,075 black rockfish (86,550) moves up to 3rd in NPS due to less brown rockfish
- 25,211 yellowtail rockfish (66,865) 11 year totals for both areas, 1980 data missing
- 20,782 canary rockfish (15,608) ranked 3rd in Greater Puget Sound in 1960s
- 19,256 yelloweye rockfish (8,761)
- 7,796 China rockfish (275) present 11 of 12 years in NPS, only 4 of 12 in PSP
- 3,668 brown rockfish (246,225)
- 2,079 bocaccio (22,890) disappointing number in NPS but present 8 of 12 years
- 1,824 tiger rockfish (none) present 11 of 12 years in NPS
- 1,150 blue rockfish (19) present 8 of 12 years in NPS, only 2 of 12 in PSP
- 929 widow rockfish (293) present 8 of 12 years in NPS, 6 of 12 years in PSP
- 664 redstripe rockfish (23,738) low number in NPS isolates species in PSP
- 582 vermilion rockfish (9) present 8 of 12 years in NPS, only 1 of 12 in PSP
- 165 silvergrey rockfish (108) only present 2 of 12 years in NPS, 3 of 12 in PSP
- 128 greenstriped rockfish (9,962) Very low number in NPS isolates species in PSP
- 70 redbanded rockfish (39) only present 2 of 12 years in NPS, 1 of 12 years in PSP

There were four species that had a relatively persistent presence in North Puget Sound but only appeared infrequently or not at all in Puget Sound Proper catch data. These four were China rockfish, tiger rockfish, blue rockfish and vermilion rockfish. These four may have existed in substantial numbers in Puget Sound Proper in the recent past but were lost somewhere along the line during a century plus of heavy exploitation. The fact that they still exist in North Puget Sound in significant numbers means that they have a potential to introduce or reintroduce each of the four species into Puget Sound Proper during the rare favorable recruitment events characteristic of rockfish. Greg Bargmann has advised me that significant numbers of vermilion rockfish were recently observed in Puget Sound Proper (Hood Canal only) and this may be the beginning of just such an introduction or reintroduction event. The fact that this has not yet happened for

the other three species (at least as fish big enough to be identified) is a powerful new confirmation of the net effect of population isolating factors prevailing in Puget Sound Proper for non-migratory rockfish species. As noted for bocaccio, there is a definite need for marine protected areas in North Puget Sound to protect the remaining populations of these four additional species.

The most surprising revelation in the catch statistics was a complete lack of tiger rockfish in Puget Sound Proper. It had been widely assumed that the species was still present in this area. However, as noted previously, the 1999 Petition (in the context of Greater Puget Sound) did identify tiger rockfish, along with canary and yelloweye rockfish, as one of three species that had largely disappeared from catch samples and were rarely seen in fishery-independent stock assessments. The 1999 Petition also stated that "not much is known regarding the tiger rockfish, but they may be the most extreme example of small home ranges. They are extremely cryptic and only inhabit very rocky reefs.". This means that their suitable habitat is limited to only a small fraction of the 14 square kilometers of rocky reef habitat. The specific problem that this poses for tiger rockfish is that anglers with good local knowledge often target the vertical walls or underwater "cliffs" that are characteristic of this type of habitat.

The annual catch records of each species in North Puget Sound were examined for possible trends over time but there were no obvious declining trends comparable to those provided in the Petition for bocaccio, greenstriped rockfish, and redstripe rockfish in Puget Sound Proper. The small catches shown for greenstriped rockfish and redstripe rockfish in North Puget Sound were also given in the Petition, confirming that both species are largely isolated from other members of the same species in Puget Sound Proper. Canary rockfish, which ranked 3rd in abundance for Greater Puget Sound during the 1960s, fell to 5th place in North Puget Sound and 8th place in Puget Sound Proper.

Special Case for Greenstriped Rockfish

Greenstriped rockfish are the only deep water species found in Puget Sound Proper in significant numbers and present some unique at-risk factors. It is surprising that they are even present in this enclosed body of water, especially since the species is almost completely absent from North Puget Sound. The species could not be reintroduced into Puget Sound Proper by a favorable recruitment event in some far distant deep water area. In addition, it is likely that the range occupied by the species is very restricted. There are only 14 square kilometers of rocky reef habitat in the entire area and only a small fraction of that is in deep water. The species was undoubtedly exploited by the commercial trawl fishery in the past since it has been encountered in recent research trawl surveys. However, some of the population may have been protected by being in rough bottom areas that could not be effectively fished by trawl gear. It was probably protected from most sport anglers until the widespread introduction of down rigger gear, and part of the population may still be too deep to be exploited. There are answers to the questions posed and these should be addressed in a NMFS status review. It is probable that the answers will justify the need for a marine protected area.