

Please provide the following information, and submit to the NOAA DM Plan Repository.

Reference to Master DM Plan (if applicable)

As stated in Section IV, Requirement 1.3, DM Plans may be hierarchical. If this DM Plan inherits provisions from a higher-level DM Plan already submitted to the Repository, then this more-specific Plan only needs to provide information that differs from what was provided in the Master DM Plan.

URL of higher-level DM Plan (if any) as submitted to DM Plan Repository:

1. General Description of Data to be Managed

1.1. Name of the Data, data collection Project, or data-producing Program:

National Status and Trends: Bioeffects Program - Biscayne Bay, Florida (1995-1996)
Database

1.2. Summary description of the data:

The toxicity of sediments in Biscayne Bay and many adjoining tributaries was determined as part of a bioeffects assessments program managed by NOAA's National Status and Trends Program. Biscayne Bay was selected by NOAA for this survey because data from the NSandT Mussel Watch Program and data from previous surveys of the bay had shown a potential for toxicity and other adverse biological effects. In addition, no bay-wide information had been generated on the toxicological condition of the bay sediments and several agencies had indicated a need for this type of data and a willingness to assist NOAA in collecting them. The study area was defined as extending from Dumbfoundling Bay at the north end to Little Card Sound at the south end, seaward to the barrier islands or reef, and landward to the shoreline or saltwater control structures. This area was determined to encompass a total of 484 kilometers of the sea floor. During 1995 and 1996, 226 samples were collected from randomly-chosen locations and tested for toxicity and analyzed for chemical concentrations. Data from these tests and analyses are included in the report. Samples for benthic community analyses were collected at one-third of the stations; however, data from those analyses are not included in the report but are available from NOAA's online database <http://nbi.noaa.gov/data/>. The survey was designed to characterize sediment quality throughout the greater Biscayne Bay area. Surficial sediment samples were collected during 1995 and 1996 from 226 randomly-chosen locations throughout nine major regions. Laboratory toxicity tests were performed as indicators of potential ecotoxicological effects in sediments. A battery of tests was performed to generate information from different phases (components) of the sediments. Tests were selected to represent a range in toxicological endpoints from acute to chronic sublethal responses. Toxicological tests were conducted to measure: reduced survival of adult amphipods exposed to solid-phase sediments; impaired fertilization success and abnormal morphological development in gametes and embryos, respectively, of sea urchins exposed to pore waters; reduced metabolic activity of a marine bioluminescent bacteria

exposed to organic solvent extracts; induction of a cytochrome P-450 reporter gene system in exposures to solvent extracts; and reduced reproductive success in marine copepods exposed to solid-phase sediments. The full report is available online at <http://ccma.nos.noaa.gov/about/coast/nsandt/download.aspx>

1.3. Is this a one-time data collection, or an ongoing series of measurements?

One-time data collection

1.4. Actual or planned temporal coverage of the data:

1995 to 1996

1.5. Actual or planned geographic coverage of the data:

W: -80.3758, E: -80.1226, N: 25.9503, S: 25.2906

1.6. Type(s) of data:

(e.g., digital numeric data, imagery, photographs, video, audio, database, tabular data, etc.)

1.7. Data collection method(s):

(e.g., satellite, airplane, unmanned aerial system, radar, weather station, moored buoy, research vessel, autonomous underwater vehicle, animal tagging, manual surveys, enforcement activities, numerical model, etc.)

1.8. If data are from a NOAA Observing System of Record, indicate name of system:

1.8.1. If data are from another observing system, please specify:

2. Point of Contact for this Data Management Plan (author or maintainer)

2.1. Name:

NCCOS Scientific Data Coordinator

2.2. Title:

Metadata Contact

2.3. Affiliation or facility:

2.4. E-mail address:

NCCOS.data@noaa.gov

2.5. Phone number:

3. Responsible Party for Data Management

Program Managers, or their designee, shall be responsible for assuring the proper management of the data produced by their Program. Please indicate the responsible party below.

3.1. Name:

NCCOS Scientific Data Coordinator

3.2. Title:

Data Steward

4. Resources

Programs must identify resources within their own budget for managing the data they produce.

4.1. Have resources for management of these data been identified?**4.2. Approximate percentage of the budget for these data devoted to data management (specify percentage or "unknown"):****5. Data Lineage and Quality**

NOAA has issued Information Quality Guidelines for ensuring and maximizing the quality, objectivity, utility, and integrity of information which it disseminates.

5.1. Processing workflow of the data from collection or acquisition to making it publicly accessible

(describe or provide URL of description):

Process Steps:

- 1996-01-01 00:00:00 - DATA ACQUISITION/FIELD SAMPLING Sediment sampling procedures are described in the report available for download at <http://ccma.nos.noaa.gov/about/coast/nsandt/download.aspx> DATA PREPARATION AND SAMPLE PROCESSING Fields are arranged as follows: Sample Type; Field Holding; Lab Storage; Max 1 Holding SEDIMENT: Organic contaminants; Wet ice (4C); Freezer (-20C); 1 year; Inorganic contaminants; Wet ice (4C); Freezer (-20C); 1 year; Total organic Carbon; Wet ice (4C); Freezer (-20C); 1 year; Grain size; Wet ice (4C); Refrigerated (4C); 1 year; TOXICITY BIOASSAY: Whole sediment and porewater bioassays; Wet ice (4C); Refrigerated (4C); 2 weeks; Organic extract (P450 and Microtox); Wet ice (4C); Freezer (-20C); 1 year; BENTHOS: Taxonomy; 10% buffered formalin; Transfer to 70% ethanol; Indefinitely;
- Chemicals with similar structural properties were summed and reported as "Totals" in addition to their individual measured concentrations. The components of these totals are as follows: Total DDT = sum of concentrations of ortho and para forms of parent and metabolites 2,4'DDE; 4,4'DDE; 2,4'DDD; 4,4'DDD; 2,4'DDT and 4,4'DDT. Total Chlordane = sum of concentrations of four compounds alpha-chlordane, trans-nonachlor, heptachlor, heptachlorepoxide. Total Dieldrin = sum of concentrations of two compounds aldrin and dieldrin. Total Butyl tin = sum of concentrations of parent compound and metabolites monobutyltin, dibutyltin, tributyltin, tetrabutyltin [concentrations are in terms of tin]. Total PCB = the sum of the concentrations of eighteen congeners: PCB8, PCB18, PCB28, PCB44, PCB52,

PCB66, PCB101, PCB105, PCB118, PCB128, PCB138, PCB153, PCB170, PCB180, PCB187, PCB195, PCB206, and PCB209. Total low molecular weight (lmw) PAHs = sum of concentrations of twelve 2- and 3-ring PAH compounds: naphthalene, 2-methylnaphthalene, 1-methylnaphthalene, biphenyl, 2,6-dimethylnaphthalene, acenaphthene, acenaphthylene, 1,6,7-trimehtylnaphthalene, fluorine, phenanthrene, 1-methylphenanthrene, and anthracene. Total high molecular weight (hmw) PAHs = sum of concentrations of twelve 4-and more-ring PAH compounds: fluoranthene, pyrene, benz[a]anthracene, chrysene, benzo[b]fluorantene, benzo[k]fluoranthene, benzo[e]pyrene, benzo[a]pyrene, perylene, dibenzathracene, indeno[1,2,3-cd]pyrene, and benzo[ahi]perylene. Total PAH = low molecular weight PAHs plus high molecular weigh PAHs (sum of 24 PAH compound concentrations). Several numerical indices were chosen for analysis and interpretation of the macroinfaunal data. Infaunal abundance is reported as the total number of individuals per station and the total number of individuals per square meter (= density). Taxa richness is reported as the number of taxa represented in a given site location. Taxa diversity, which is often related to the ecological stability and environmental "quality" of the benthos, was estimated by the Shannon-Weiner Index (Shannon and Weaver, 1949). In order to quantify and compare the equitability in the fauna to the taxa diversity for a given area, Pielou's Evenness Index J' (Pielou, 1966) was calculated as $J' = H'/\ln S$, where $\ln S = H'_{\max}$, or the maximum possible diversity, when all taxa are represented by the same number of individuals; thus, $J' = H' / H'_{\max}$.

- Project Related References:Pielou, E.C. 1966. The Measurement of Diversity in Different Types of Biological Collections. J. Theoretical Biology 13:131-144.Shannon, L.C. and W. Weaver. 1949. The Mathematical Theory of Communication. Univ. of Illinois Press, Urbana, Ill. 117 p.

5.1.1. If data at different stages of the workflow, or products derived from these data, are subject to a separate data management plan, provide reference to other plan:

5.2. Quality control procedures employed (describe or provide URL of description):

6. Data Documentation

The EDMC Data Documentation Procedural Directive requires that NOAA data be well documented, specifies the use of ISO 19115 and related standards for documentation of new data, and provides links to resources and tools for metadata creation and validation.

6.1. Does metadata comply with EDMC Data Documentation directive?

No

6.1.1. If metadata are non-existent or non-compliant, please explain:

Missing/invalid information:

- 1.6. Type(s) of data

- 1.7. Data collection method(s)
- 4.1. Have resources for management of these data been identified?
- 4.2. Approximate percentage of the budget for these data devoted to data management
- 5.2. Quality control procedures employed
- 7.1. Do these data comply with the Data Access directive?
 - 7.1.1. If data are not available or has limitations, has a Waiver been filed?
 - 7.1.2. If there are limitations to data access, describe how data are protected
- 7.2. Name of organization of facility providing data access
 - 7.2.1. If data hosting service is needed, please indicate
- 7.3. Data access methods or services offered
- 7.4. Approximate delay between data collection and dissemination
- 8.1. Actual or planned long-term data archive location
- 8.3. Approximate delay between data collection and submission to an archive facility
- 8.4. How will the data be protected from accidental or malicious modification or deletion prior to receipt by the archive?

6.2. Name of organization or facility providing metadata hosting:

NMFS Office of Science and Technology

6.2.1. If service is needed for metadata hosting, please indicate:

6.3. URL of metadata folder or data catalog, if known:

<https://www.fisheries.noaa.gov/inport/item/38735>

6.4. Process for producing and maintaining metadata

(describe or provide URL of description):

Metadata produced and maintained in accordance with the NOAA Data Documentation Procedural Directive: https://nosc.noaa.gov/EDMC/DAARWG/docs/EDMC_PD-Data_Documentation_v1.pdf

7. Data Access

NAO 212-15 states that access to environmental data may only be restricted when distribution is explicitly limited by law, regulation, policy (such as those applicable to personally identifiable information or protected critical infrastructure information or proprietary trade information) or by security requirements. The EDMC Data Access Procedural Directive contains specific guidance, recommends the use of open-standard, interoperable, non-proprietary web services, provides information about resources and tools to enable data access, and includes a Waiver to be submitted to justify any approach other than full, unrestricted public access.

7.1. Do these data comply with the Data Access directive?

7.1.1. If the data are not to be made available to the public at all, or with limitations, has a Waiver (Appendix A of Data Access directive) been filed?

7.1.2. If there are limitations to public data access, describe how data are protected from unauthorized access or disclosure:

7.2. Name of organization of facility providing data access:

7.2.1. If data hosting service is needed, please indicate:

7.2.2. URL of data access service, if known:

<https://products.coastalscience.noaa.gov/collections/ltmonitoring/nsandt/default.aspx>

<https://products.coastalscience.noaa.gov/collections/ltmonitoring/nsandt/default.aspx>

7.3. Data access methods or services offered:

7.4. Approximate delay between data collection and dissemination:

7.4.1. If delay is longer than latency of automated processing, indicate under what authority data access is delayed:

8. Data Preservation and Protection

The NOAA Procedure for Scientific Records Appraisal and Archive Approval describes how to identify, appraise and decide what scientific records are to be preserved in a NOAA archive.

8.1. Actual or planned long-term data archive location:

(Specify NCEI-MD, NCEI-CO, NCEI-NC, NCEI-MS, World Data Center (WDC) facility, Other, To Be Determined, Unable to Archive, or No Archiving Intended)

8.1.1. If World Data Center or Other, specify:

8.1.2. If To Be Determined, Unable to Archive or No Archiving Intended, explain:

8.2. Data storage facility prior to being sent to an archive facility (if any):

National Centers for Coastal Ocean Science - Silver Spring, MD

8.3. Approximate delay between data collection and submission to an archive facility:

8.4. How will the data be protected from accidental or malicious modification or deletion prior to receipt by the archive?

Discuss data back-up, disaster recovery/contingency planning, and off-site data storage relevant to the data collection

9. Additional Line Office or Staff Office Questions

Line and Staff Offices may extend this template by inserting additional questions in this section.