

**ENVIRONMENTAL ASSESSMENT,
FINDING OF NO SIGNIFICANCE, AND
REGULATORY IMPACT REVIEW
FOR THE 2023 EMERGENCY RULE TO REDUCE RIGHT WHALE INTERACTIONS
WITH TRAP/POT GEAR**

**February 2023
Updated March 31, 2023 (See Errata, Appendix A)**

Contents:

Environmental Assessment	2
Appendices	100
Finding of No Significance	229
Regulatory Impact Review	236

**US DEPARTMENT OF COMMERCE
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
NATIONAL MARINE FISHERIES SERVICE
GREATER ATLANTIC REGIONAL FISHERIES OFFICE**

**ENVIRONMENTAL ASSESSMENT OF A PROPOSAL TO PROMULGATE A 2023
EMERGENCY RULE TO REDUCE RIGHT WHALE INTERACTIONS WITH
TRAP/POT GEAR**

**February 2023
Updated March 31, 2023 (See Errata, Appendix A)**

**US DEPARTMENT OF COMMERCE
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
NATIONAL MARINE FISHERIES SERVICE
GREATER ATLANTIC REGIONAL FISHERIES OFFICE**

1 EXECUTIVE SUMMARY

NMFS prepared an Environmental Assessment (EA) of a 2022 Emergency Final Rule to Reduce Right Whale Interactions with Lobster and Jonah Crab Trap/Pot Gear (described in this document as the 2022 Emergency Rule EA) to analyze the emergency closure of the waters between state and federal portions of the Massachusetts Restricted Area (MRA) to trap/pot buoy lines for a second year for the full length of the MRA seasonal closure (February 1-April 30). As described below, NMFS is proposing to extend that closure and has prepared this EA to analyze the environmental effects of alternative means of promulgating an emergency rule for a similar closure consistent with the stated purpose and need for action. This EA substantially incorporates and relies on the 2022 Emergency Rule EA.

Summary of Purpose and Need and Proposed Action

The Proposed Action, as described in Section 3, would close a small area (the MRA Wedge) in Massachusetts Bay nearly circumscribed by the seasonal MRA to trap/pot fishery buoy lines. This action would extend the Emergency Rule issued last March (87 FR 11590; March 2, 2022) into 2023 and is needed to reduce the risk to North Atlantic right whales (*Eubalaena glacialis*) from potential mortality and serious injury due to entanglement in buoy lines in an area that would otherwise have a high co-occurrence of whales and buoy lines.

At the time of the 2022 emergency action, NMFS anticipated that upcoming modifications to the Atlantic Large Whale Take Reduction Plan (ALWTRP or Plan) would address the risk of entanglement in the MRA Wedge during seasonal restrictions of the MRA. However, as a result of litigation outcomes NMFS was not able to promulgate an amendment to the ALWTRP in time for the 2023 season (see Section 3 for more detail). Accordingly, and based on the new data available from 2021 and 2022 regarding the co-occurrence of whales and buoy lines in this area, the extension of the emergency closure is necessary to mitigate the risk of mortality or serious injury from entanglement in fishing gear to right whales in the area while the MRA is closed in 2023.

Under the No Action Alternative the emergency closure would not be promulgated or implemented, leaving the current Plan intact with no regulatory changes proposed. This includes the restricted areas implemented by the Final Rule on September 17, 2021 (86 FR 51970) that went into effect October 18, 2021 and requirements for minimum traps per trawl and weak inserts throughout the buoy line that went into effect May 1, 2022.

Affected Environment

The Affected Environment for the 2023 closure is described in Section 5. The Affected Environment is described based on the valued ecosystem components (VECs) that may be affected by the three alternatives within the portion of Lobster Management Area 1 adjacent to Massachusetts (i.e. south of the New Hampshire border, MA LMA 1; action area). The proposed action is not expected to have significant impacts on the biological aspects of the fisheries and therefore fish/lobster biology is not included in this analysis. The three major VECs potentially affected by the proposed action are protected species, habitat, and the human community.

Impacts of the Alternatives

The impacts of the alternatives considered on each VEC described in the Affected Environment are in Section 6 and summarized here. Alternative 1 (No Action) would maintain the status quo to the current Plan intact as implemented in 2021. Alternative 2 (Preferred Alternative) would establish a closure that restricts the use of persistent trap/pot gear buoy lines from February 1, 2023 through April 30, 2023. The proposed restricted area within Alternative 2 closes 200 square miles (518 square kilometers) of federal waters adjacent to the existing MRA. Alternative 3 would also establish a closure that restricts the use of persistent trap/pot gear buoy lines from February 1, 2023 through April 30, 2023. The restricted area within Alternative 3 would encompass approximately 1,297 square miles (3,359 square kilometers) and extend the northern MRA boundaries up to the New Hampshire border.

Protected species

The primary difference in biological impacts on protected species between the alternatives relates to the removal of buoy lines within the water column that are directly related to the reduction in right whale entanglement risk within the closure area. Prohibiting the use of buoy lines from February 1, 2023 to April 30, 2023 would reduce entanglement risk for large whales, right whales in particular. Under Alternative 1 (No Action), high negative impacts are expected because there would be a risk of entanglement due to the number of buoy lines that would remain in the water when right whales are abundant in the MRA Wedge. Relative to No Action, Alternative 2 would have a slight positive impact on ESA-listed (right, fin, and sei whales) and MMPA protected species (humpback and minke whales) because large whale entanglement risk in trap/pot gear is reduced. Relative to No Action, Alternative 3 would have a moderate positive impact on ESA-listed and MMPA protected species. Considered alone, ESA-listed and MMPA protected species would be moderately negative to slightly negatively impacted by Alternative 2, and negligible to slightly negatively impacted by Alternative 3, because this proposed emergency action does not eliminate the potential for all interaction risk between fishing gear and marine mammals that could result in takes above PBR.

Habitat

Alternative 1 (No Action) will maintain baseline levels of biological impacts on benthic habitats, negligible to slight negative impacts to habitat due to disturbance to benthic habitat. In comparison to No Action, Alternative 2 and Alternative 3 are expected to have a negligible to slight positive impact on the MA LMA 1 habitat. If ropeless fishing is implemented in closed areas, it is not expected that Alternative 2 or Alternative 3 will significantly change the amount of gear that comes into contact with the seafloor and is unlikely to occur at levels similar to those prior to the closure. Compared to Alternative 2 and Alternative 3, No Action is expected to have negligible impacts on affected fish habitats. Considered on their own, Alternative 2 and Alternative 3 would likely have a negligible to slight negative impact on the environment due to continued disturbance from long trawls outside of the closure period.

Human Community

Alternative 1 (No Action) would maintain the status quo, which has a slight negative to moderate positive impact on fishing communities. Alternative 2, the Preferred Alternative, is expected to have a slight negative impact on the fishing communities impacted by this action. Overall, the economic impacts of the Alternative 2 result in an estimated total cost (including lost revenue) of

\$339,000 to \$608,000 with approximately 30 affected vessels, compared to No Action. Alternative 2 would impact lobster and Jonah crab vessels in Southern Essex County, Suffolk County, Norfolk County, and Northern Plymouth County. Vessels in Plymouth County could be the most vulnerable to the Emergency Rule, while Suffolk County might be the least vulnerable. Alternative 3 is expected to have a moderate negative impact on the human community VEC, as defined here. Alternative 3 is estimated to impact 53 to 66 vessels for a total estimated cost (including lost revenue) of \$898,249 to \$1,452,797, compared to No Action. Alternative 3 has similar social impacts to Alternative 2, except it will affect more vessels in Essex County that fish in the Northern waters offshore and north of Cape Ann.

Cumulative Impacts

We analyzed the impacts of all alternatives on physical habitat, protected species, and human communities. When the proposed action (Alternative 2, the Preferred Alternative) is considered in conjunction with all other impacts from past, present, and reasonably foreseeable future actions, it is not expected to result in any significant impacts, positive or negative; therefore, no significant cumulative effects on the human environment are associated with the proposed action (Subsection 6.5).

Conclusions

A description of the expected environmental impacts and any cumulative impacts of the proposed action are provided in Section 6. The proposed action is not associated with significant impacts to the socioeconomic or physical environment, individually or in conjunction with other actions; therefore, a “Finding of No Significant Impact” is warranted.

2 TABLE OF CONTENTS

Contents

GREATER ATLANTIC REGIONAL FISHERIES OFFICE	i
1 EXECUTIVE SUMMARY	ii
2 TABLE OF CONTENTS	v
2.1 <i>List of Tables</i>	vii
2.2 <i>List of Figures</i>	vii
2.3 <i>List of Acronyms</i>	viii
3 INTRODUCTION AND PURPOSE	1
3.1 <i>Background</i>	1
3.1.1 Large Whale Entanglement Risk in the Action Area	1
3.1.2 Justification for an Emergency Closure in 2023	5
3.2 <i>Purpose and Need</i>	9
3.2.1 Scope of the Analysis	9
4 SUMMARY OF MANAGEMENT ALTERNATIVES	9
4.1 <i>Alternative 1: No Action (Status Quo)</i>	9
4.2 <i>Alternative 2: Preferred</i>	10
4.3 <i>Alternative 3</i>	11
5 DESCRIPTION OF THE AFFECTED ENVIRONMENT	12
5.1 <i>Protected Species</i>	12
5.1.1 Protected Species: Atlantic Large Whales	14
5.1.2 Species and Critical Habitat Not Likely to be Impacted	21
5.2 <i>Habitat</i>	21
5.2.1 Alteration of Physical Structure	26
5.2.2 Mortality of Benthic Organisms	26
5.2.3 Changes to Benthic Communities and Ecosystems	27
5.2.4 Sediment Suspension	27
5.2.5 Chemical Modifications	27
5.2.6 American Lobster Habitat	28
5.3 <i>Human Community</i>	30
5.3.1 Affected Fisheries	30
5.3.2 Affected Human Communities	32
6 IMPACTS OF THE MANAGEMENT ALTERNATIVES	35
6.1 <i>Impact Designation Descriptions</i>	35
6.1.1 Protected Species	37
6.1.2 Habitat	37
6.1.3 Human Community	38

6.2	<i>Evaluating the Protected Species Impacts of the Alternatives</i>	38
6.2.1	Overview of the Decision Support Tool Analysis	38
6.2.2	The Impacts of Alternative 1 on Protected Species	41
6.2.3	The Impacts of Alternative 2 on Protected Species	45
6.2.4	The Impacts of Alternative 3 on Protected Species	51
6.2.5	Comparison and Summary of the Alternatives	53
6.3	<i>Habitat Impacts of the Alternatives</i>	53
6.3.1	Alternative 1: No Action	53
6.3.2	Alternative 2: Preferred Action	53
6.3.3	Alternative 3	54
6.3.4	Comparison and Summary of the Alternatives	55
6.4	<i>Human Community Impacts of the Alternatives</i>	56
6.4.1	Economic Impacts of the Alternatives	56
6.4.2	Social Impacts of the Alternatives	61
6.4.3	Comparison and Summary of Impacts to Human Communities	62
6.5	<i>Cumulative Impacts of the Alternatives</i>	62
6.5.1	Introduction	62
6.5.2	Summary of Direct and Indirect Impacts of the Alternatives	64
6.5.3	Status Quo Conditions	65
6.5.4	Past, Present, and Reasonably Foreseeable Future Actions	65
6.5.5	Cumulative Effects Analysis	70
6.6	<i>Summary</i>	71
7	APPLICABLE LAWS AND REGULATIONS	71
7.1	<i>Endangered Species Act</i>	71
7.2	<i>Marine Mammal Protection Act</i>	71
7.3	<i>Paperwork Reduction Act</i>	72
7.4	<i>Magnuson-Stevens Fishery Conservation and Management Act including Essential Fish Habitat</i>	72
7.5	<i>Information Quality Act (Public Law 106-554)</i>	72
7.6	<i>Administrative Procedure Act</i>	74
7.7	<i>Coastal Zone Management Act</i>	74
7.8	<i>Executive Order (E.O.) 13132 Federalism</i>	75
7.9	<i>Regulatory Flexibility Act</i>	75
7.10	<i>E.O. 12866 Regulatory Planning and Review</i>	75
7.11	<i>Consolidated Appropriations Act, 2023</i>	75
7.12	<i>National Environmental Policy Act</i>	76
7.12.1	Environmental Assessment	76
7.12.2	Point of Contact	77
7.12.3	Agencies Consulted	77
7.12.4	List of Preparers	77
8	References	79

2.1 List of Tables

Table 1: Species protected under the ESA and/or MMPA that may occur in the Affected Environment.....	13
Table 2: Estimated abundance, potential biological removal level, and average annual observed mortality for Atlantic Large Whales	17
Table 3: List of Essential Fish Habitat for different species and life history stages within Affected Environment.....	22
Table 4: American lobster landing pounds in Massachusetts counties by year from 2017-2021	31
Table 5: Jonah crab landing pounds in Massachusetts counties by year from 2017-2021	32
Table 6: Social-economic indicators for coastal communities	34
Table 7: Impact determinations for Valued Ecosystem Components.....	36
Table 8: Estimated total number of Atlantic large whales in Action Area	42
Table 9: Comparison of North Atlantic right whale entanglement risk reduction by month	46
Table 10: Number of affected vessels and landing values 2017-2021 under Alternative 2 (Preferred)	58
Table 11: Number of affected vessels and landing values 2017-2021 under Alternative 3	58
Table 12: Cost savings for vessels that stop fishing during closure months.....	59
Table 13: Economic impacts of Alternative 2 (Preferred) by month.....	60
Table 14: Economic impacts of Alternative 3 by month	61
Table 15: Socio-economic profile of affected communities - Harvest Parameters	62
Table 16: Direct and indirect impacts of the alternatives on Valued Ecosystem Components ...	64
Table 17: Summary of the current status and trends of the Valued Ecosystem Components	65
Table 18: Summary table of the final cumulative impacts analysis of the Preferred Alternative (Alternative 2) on all three VECs	70

2.2 List of Figures

Figure 1: Massachusetts Restricted Area and emergency closure with Alternative 2, MRA Wedge	3
Figure 2: North Atlantic Right Whale sightings and fishing gear observed in 2021 and 2022 within Proposed Emergency Closure, MRA Wedge	4
Figure 3: North Atlantic right whale visual sightings February, March, and April 2018-2022	7
Figure 4: Acoustic detections of North Atlantic right whale within Massachusetts Bay February, March, and April 2018-2022.....	8
Figure 5: Alternative 2 (Preferred) Proposed Emergency Closure, MRA Wedge	10
Figure 6: Alternative 3 Proposed Emergency Closure, MRA Wedge North.....	11
Figure 7: Habitat Areas of Particular Concern and Essential Fish Habitat in Action Area.....	22
Figure 8: Depiction of the relative direction and magnitude of impacts on valued ecosystem components	37
Figure 9: North Atlantic Right Whale sightings and fishing gear observed in 2021 and 2022 within Proposed Emergency Closure, MRA Wedge	43
Figure 10: Fishing gear observed within Massachusetts Bay in spring 2021 and 2022 within Proposed Emergency Closure, MRA Wedge.....	44
Figure 11: Acoustic detections of fin, sei, and right whales in Massachusetts Bay from March 17, 2021 through April 30, 2021	47

Figure 12: Visual sightings of Atlantic large whales in Massachusetts Bay during February, March, and April 2018-2022.....	49
Figure 13: Acoustic detections of humpback whales in Massachusetts Bay from March 17, 2021 through April 30, 2021.....	50
Figure 14: Cumulative effects analysis steps.....	63

2.3 List of Acronyms

ALWTRP	Atlantic Large Whale Take Reduction Plan
ALWTRT	Atlantic Large Whale Take Reduction Team
APA	Administrative Procedure Act
ASMFC	Atlantic States Marine Fisheries Commission
CCS	Center for Coastal Studies
CEA	Cumulative Effects Analysis
CEQ	Council on Environmental Quality
CZMA	Coastal Zone Management Act of 1972
DST	Large Whale Decision Support Tool
EA	Environmental Assessment
EEZ	Exclusive Economic Zone
EFH	Essential Fish Habitat
EFP	Exempted Fishing Permit
EO	Executive Order
ESA	Endangered Species Act of 1973
FEIS	Final Environmental Impact Statement
FMP	Fishery Management Plan
GARFO	Greater Atlantic Regional Fisheries Office
IQA	Information Quality Act
LMA 1	Lobster Management Area 1
MA LMA 1	Massachusetts' portion of the Lobster Management Area 1
MMPA	Marine Mammal Protection Act of 1972
MRA	Massachusetts Restricted Area
MSA	Magnuson-Stevens Fishery Conservation and Management Act
NEFMC	New England Fishery Management Council
NEFSC	Northeast Fisheries Science Center
NEPA	National Environmental Policy Act
NMFS	National Marine Fisheries Service
NOAA	National Oceanic and Atmospheric Administration
PBR	Potential Biological Removal
PRA	Paperwork Reduction Act
RFA	Regulatory Flexibility Act
RIR	Regulatory Impact Review
SAR	Stock Assessment Report
SSB	Social Science Branch
UME	Unusual Mortality Event
VEC	Valued Ecosystem Component
VTR	Vessel Trip Report

3 INTRODUCTION AND PURPOSE

In accordance with the National Environmental Policy Act (NEPA), the regulations published by the Council on Environmental Quality, and NOAA's Companion Manual for NAO 216-6A (2017), this Environmental Assessment evaluates potential environmental impacts of an emergency rule proposed for implementation by NOAA's National Marine Fisheries Service (NMFS) under Section 118 of the Marine Mammal Protection Act (MMPA) to modify the regulations implementing the Atlantic Large Whale Take Reduction Plan (ALWTRP or Plan). Significant impacts are not anticipated as a result of this action because the emergency closure area is small relative to the entire Massachusetts Restricted Area (MRA), minimal economic impacts are expected, and any indirect effects are likely beneficial for the environment.

3.1 Background

3.1.1 Large Whale Entanglement Risk in the Action Area

The North Atlantic right whale (*Eubalaena glacialis*, hereafter referred to as right whale) population has been in decline since 2010, with the most recent published estimate of right whale population size in 2020 at 338 whales (95 percent confidence interval: 325-350) with a strong male bias (Hayes et al. 2023, Pace et al. 2017, Pace 2021). The steep population decline is a result of high levels of human-caused mortality caused by entanglement in fishing gear and vessel strikes in both the U.S. and Canada. An Unusual Mortality Event (UME) was declared for the population in 2017, as a result of high rates of documented vessel strikes and entanglement in fishing gear. As of January 11, 2023, the UME includes 35 detected mortalities (17 in 2017, 3 in 2018, 10 in 2019, 2 in 2020, 2 in 2021, 0 in 2022, and, 2 in 2023). In addition, 22 serious injuries were documented (2 in 2017, 5 in 2018, 1 in 2019, 4 in 2020, 5 in 2021, 4 in 2022, and, tentatively, 1 in 2023). Lastly, 37 morbidity (or sublethal injury or illness) cases were documented (12 in 2017, 10 in 2018, 7 in 2019, 5 in 2020, 1 in 2021, and 2 in 2022) (<https://www.fisheries.noaa.gov/national/marine-life-distress/2017-2023-north-atlantic-right-whale-unusual-mortality-event>). Population models estimate that 64 percent of all mortalities are not seen and not accounted for in the right whale observed incident data (Pace 2021, Pace et al. 2017).

The right whale is listed as an endangered species under the Endangered Species Act of 1973 (ESA) and considered a strategic stock under the MMPA. NMFS is required by the MMPA to reduce mortality and serious injury incidental to commercial fishing to below a stock's potential biological removal (PBR) level. This is defined as the maximum number of animals that can be removed annually, while allowing a marine mammal stock to reach or maintain its optimal sustainable population level. PBR for the right whale population is 0.7 whales per year in the most recently published stock assessment report (Hayes et al. 2023). Between 2010 and 2021, there has not been one year where observed mortality and serious injury of right whales fell below a PBR of 0.7. With the total estimated mortality well above this number, additional measures are urgently needed to reduce the impact of U.S. Atlantic fisheries on right whales.

The Plan was originally developed pursuant to Section 118 of the MMPA (16 U.S.C. 1387) to reduce mortality and serious injury of three stocks of large whales (fin, humpback, and North


Atlantic right) incidental to certain Category I and II fisheries. Under the MMPA, a strategic stock of marine mammals is defined as a stock: (1) for which the level of direct human-caused mortality exceeds the Potential Biological Removal (PBR) level; (2) which, based on the best available scientific information, is declining and is likely to be listed as a threatened species under the ESA (1973) within the foreseeable future; or (3) which is listed as a threatened or endangered species under the ESA or is designated as depleted under the MMPA (16 U.S.C. 1362(19)). When incidental mortality or serious injury of marine mammals from commercial fishing exceeds a stock's PBR level, the MMPA directs NMFS to convene a take reduction team of stakeholders that includes the following: representatives of Federal agencies, each coastal state that has fisheries interacting with the species or stock, appropriate Regional Fishery Management Councils, interstate fisheries commissions, academic and scientific organizations, environmental groups, all commercial and recreational fisheries groups using gear types that incidentally take the species or stock, and if relevant, Alaska Native organizations or Indian tribal organizations.

The Atlantic Large Whale Take Reduction Team (ALWTRT) was established in 1996 and has 60 members, including approximately 22 trap/pot and gillnet fishermen or fishery representatives. The background for the take reduction planning process and initial development of the Plan is provided in the preambles to the proposed (62 FR 16519, April 7, 1997), interim final (62 FR 39157, July 22, 1997), and final (64 FR 7529, February 16, 1999) rules implementing the initial plan. The ALWTRT met and recommended modifications to the Plan, implemented by NMFS through rulemaking, several times since 1997 in an iterative effort to meet the MMPA take reduction goals.

Mortalities and serious injuries of right whales confirmed in U.S. fishing gear or first seen in U.S. waters with an entanglement continue at levels exceeding the right whale's PBR. NMFS informed the ALWTRT in late 2017 that it was necessary to reconvene to develop recommendations to reduce the impacts of U.S. commercial fisheries on large whales, with a focus on reducing risk to the declining right whale population. During an ALWTRT meeting in April 2019, the ALWTRT recommended a framework of measures to modify lobster and Jonah crab trap/pot trawls within the Northeast Region Trap/Pot Management Area (Northeast Region). In 2021, the recommended measures were estimated to reduce the risk of mortality and serious injury to right whales incidentally entangled in buoy lines in those fisheries by at least 60 percent. At that time, this was the best estimate of the minimum amount of risk necessary to get annual mortality and serious injury rates below PBR based on observed entanglements. NMFS published a Final Environmental Impact Statement (FEIS) on July 2, 2021 (86 FR 35288), with a 30-day comment period. The Record of Decision was signed on August 30, 2021, and the final rule was published on September 17, 2021 (86 FR 51970). The new rule was estimated to meet the minimum 60-percent reduction in risk recommended by the ALWTRT in 2019. Further detail on right whale population estimates, the stock's decline, changes in distribution and reproductive rates, as well as entanglement-related mortalities and serious injuries that were documented in the years leading up to the final rule can be found in Chapters 2 and 4 of the FEIS (NMFS 2021) and the preamble to the 2021 final rule (86 FR 51970).

Figure 1: Massachusetts Restricted Area and emergency closure with Alternative 2, MRA Wedge

The Massachusetts Restricted Area (green shaded area), Massachusetts state waters (crosshatched), and MRA Wedge (simple hatched gray shaded area) are represented. Federal and state waters within the Massachusetts Restricted Area are seasonally closed to the use of vertical buoy lines from February 1 through April 30. Massachusetts State regulations prohibit trap/pot fishing in any waters under the jurisdiction of the Commonwealth from February 1 through May 15, but can be extended the seasonal closure past May 15 in the continued presence of North Atlantic Right whales or rescinded after April 30 in their absence (322 CMR 12.04(2)). The MRA Wedge would remain closed to trap/pot fishing with buoy lines from February 1, 2023 through April 30, 2023, under Alternative 2 (Preferred Alternative). MRA




The 2021 final rule (86 FR 51970, September 17, 2021) left a critical gap in protection of right whales within the MRA, as suggested by sighting data that indicate a high risk of overlap between right whales and buoy lines. The 2021 expansion of the geographic extent of the MRA, to include Massachusetts state waters north to the New Hampshire border (Figure 1) mirrored the Massachusetts 2021 modification of the state water closure (322 CMR 12.04(2)). The implementation of the MRA Expansion, left approximately 200 square miles (518 square kilometers) of federal waters open to trap/pot fishing between state and federal closures. This created the “MRA Wedge” (Figure 1). Center for Coastal Studies (CCS) aerial survey data from 2021 and 2022 indicated that trap/pot gear was concentrated in the MRA Wedge during the MRA closure period (Figure 2). Additionally, CCS and the Northeast Fisheries Science Center (NEFSC) observed right whales within this wedge alongside the presence of aggregated fixed fishing gear during aerial surveys in April 2021 and March and April of 2022 (Figure 2). In early 2022, NMFS received letters and emails from Massachusetts Division of Marine Fisheries,

Stellwagen Bank National Marine Sanctuary, and non-governmental organizations expressing concerns about this gap in restricted waters and the heightened risk of entanglement for right whales. After reviewing available information and due to the high risk of entanglement in this relatively small area, NMFS issued an emergency rule prohibiting trap/pot fishery buoy lines between federal and state waters within the MRA for the month of April in 2022 (87 FR 11590; March 2, 2022).

Figure 2: North Atlantic Right Whale sightings and fishing gear observed in 2021 and 2022 within Proposed Emergency Closure, MRA Wedge

North Atlantic right whale sightings collected February, March, and April 2018-2022 (red and blue crosses) and gear observed (black stars) on four different days are overlaid. Blue crosses show right whale sightings collected through dedicated aerial surveys conducted by the Center for Coastal Studies (CCS) and Northeast Fisheries Science Center (NEFSC) and dedicated shipboard surveys conducted by CCS and Stellwagen Bay National Marine Sanctuary. Red crosses are opportunistic sightings collected by NEFSC. Black stars represent fishing gear observed by CCS on April 19 and 28, 2021 and February 6 and March 11, 2022, four days that were selected as representative snapshots of fishing gear present in survey areas. Surveys concentrate on Cape Cod Bay; surveyors rarely fly north of mid Cape Ann, off Rockport, MA. Massachusetts Restricted Area is shaded in dark gray, and MRA Wedge (Alternative 2) is represented by the light gray area. These maps are used for qualitative not quantitative comparison, and differ from Large Whale Decision Support Tool data.



3.1.2 Justification for an Emergency Closure in 2023

At the time of the 2022 emergency action, NMFS had already begun the rulemaking process for a second round of modifications to the ALWTRP, because new population information indicated a need for further risk reduction to reduce mortality and serious injury of right whales below PBR in U.S. commercial fisheries. Concurrently, NMFS faced litigation on the 2021 Batched Fisheries Biological Opinion issued under the ESA and the 2021 amendment to the ALWTRP issued under the MMPA (86 FR 51970; September 17, 2021). On July 8, 2022 the District Court for the District of Columbia held that the 2021 final ALWTRP rule violated the MMPA for failing to include measures expected to reduce mortality and serious injury to below the PBR level within six months of implementation. (Center for Biological Diversity, et al., v. Raimondo, et al., (Civ. No. 18-112 (D.D.C.))). As a result, on September 9, 2022, NMFS announced it was scoping in advance of additional rulemaking (87 FR 55405) to meet its MMPA mandate as described by the Court's decision. Then on November 17, 2022, the Court ordered NMFS to promulgate a new MMPA compliant ALWTRP rule by December 9, 2024. (Center for Biological Diversity, et al., v. Raimondo, et al., (Civ. No. 18-112 (D.D.C.))).

When the 2022 emergency rulemaking was published, NMFS anticipated that the upcoming modifications to the ALWTRP would address the risk associated with the lack of seasonal restrictions in the MRA Wedge. However, in light of the Court's decisions, an ALWTRP rule addressing the MRA Wedge area was not feasible by February 2023, given that the Court instructed NMFS to promulgate the ALWTRP amendment with measures necessary to meet the PBR level within six months of implementation and the ALWTRT had not completed deliberations on recommended measures until December 2, 2022. Accordingly, the entanglement risk associated with a lack of seasonal restrictions in the MRA Wedge could not be feasibly addressed by an ALWTRP amendment in time to mitigate an immediate and significant adverse impact to right whales in the MRA Wedge during the MRA closure in 2023.

On December 29, 2022, President Biden signed H.R. 2617, the Consolidated Appropriations Act, 2023 (Consolidated Appropriations Act) into law. The Consolidated Appropriations Act establishes that from December 29, 2022, through December 31, 2028, the NMFS' September 17, 2021 rule amending the ALWTRP, Taking of Marine Mammals Incidental to Commercial Fishing Operations; Atlantic Large Whale Take Reduction Plan Regulations, 86 Fed. Reg. 51,970 (Sept. 17, 2021), "shall be deemed sufficient to ensure that the continued Federal and State authorizations of the American lobster and Jonah crab fisheries are in full compliance" with the MMPA and the ESA. H.R. 2617-1631–H.R. 2617-1632 (Division JJ–North Atlantic Right Whales, Title I–North Atlantic Right Whales and Regulations, § 101(a)). The Consolidated Appropriations Act disrupts the Court's 2024 deadline and requires that NMFS promulgate new regulations for the American lobster and Jonah crab fisheries, consistent with the MMPA and ESA, to take effect by December 31, 2028. *Id.* Additionally, the Consolidated Appropriations Act contains an exception at § 101(b), stating that § 101(a) does not apply to "an existing emergency rule, or any action taken to extend or make final an emergency rule" which is a specific reference to this emergency rulemaking. Therefore, this emergency rule is justified, in part, as a consequence of litigation outcomes and changing circumstances following the 2022 emergency rule.

Right whales are known to aggregate in Cape Cod Bay (Bay) in winter and spring to forage on copepods. As this food source declines, right whale distribution accordingly shifts, increasing the presence of right whales in the MRA Wedge as they leave the Bay. Federal waters reopen to trap/pot fishing in May, increasing the area available to fishermen and reducing the likelihood of high gear density from fishermen “storing” their gear in the MRA Wedge. Aerial surveys from 2021 and 2022 capturing gear sightings on specific days when surrounding waters of the MRA are closed to buoy lines, demonstrate the high risk of entanglement that right whales face while in or traversing the waters of the MRA Wedge (Figure 2). Additionally, sightings and acoustic detections of right whales throughout the spring in the MRA Wedge and surrounding waters continue to demonstrate that right whales are in the MRA Wedge or likely traveling through this gap in the MRA closure to feed in waters in and around Massachusetts Bay (Figure 3, Figure 4). Without restrictions in place in the MRA Wedge, gear may increase in this area as fishermen pushed out of surrounding waters move gear into this small open area and continue to actively fish. Gear may also increase if fishermen start bringing gear into the MRA Wedge anticipating the May 1 opening of federal waters. The staging of gear in anticipation of federal waters opening may be especially likely in April when right whale sightings are still high. Given the high likelihood that endangered right whales are present throughout this area and in adjoining waters February through April, the MRA Wedge poses a particularly high risk of mortality or serious injury from entanglement in fishing gear. Accordingly, it is critical that this buoy line closure be implemented simultaneously with similar protections in adjacent federal waters during 2023 to prevent the likelihood of an immediate and significant adverse impact on right whales in the MRA Wedge.

Figure 3: North Atlantic right whale visual sightings February, March, and April 2018-2022

North Atlantic Right Whale sightings spanning February-April 2018-2022 in the closed area (pale gray) are represented below. The black crosses are sightings observed during dedicated Northeast Fisheries Science center (NEFSC) and Center for Coastal Studies (CCS) aerial surveys and dedicated NEFSC, CCS, and Stellwagen Bay National Marine Sanctuary shipboard surveys. The open crosses represent sightings opportunistically observed from various platforms. Surveys concentrate on Cape Cod Bay; surveyors rarely fly north of mid Cape Ann, off Rockport, MA. Massachusetts Restricted Area is shaded in dark gray, and MRA Wedge (Alternative 2) is represented by the light gray area). These maps are used for qualitative not quantitative comparison, and differ from Large Whale Decision Support Tool data.





Figure 4: Acoustic detections of North Atlantic right whale within Massachusetts Bay February, March, and April 2018-2022

Definite (red squares) and possible (green squares) acoustic detections of North Atlantic right whales in Massachusetts Bay from February 1 through April 30, 2018-2022. The acoustic detection data comes from Slocum Glider surveys deployed near the Stellwagen Bank National Marine Sanctuary and Gulf of Maine (Baumgartner 2021). Survey track lines from the Slocum Glider surveys are depicted by the gray lines. Data from Passive Acoustic Cetacean Map Website (PACM 2023, <https://apps-nefsc.fisheries.noaa.gov/pacm/#/narw>). These maps are used for qualitative not quantitative comparison, and differ from Large Whale Decision Support Tool data.



3.2 Purpose and Need

The purpose of the emergency rule is to reduce the acute risk of right whale entanglement in trap/pot fisheries in waters adjacent to the existing MRA from February 1, 2023 through April 30, 2023. Recent survey data demonstrates the likelihood of high overlap between right whales and buoy lines in this area. There is an urgent need to prevent the take of right whales in U.S. commercial fisheries because even one take that causes mortality or serious injury exceeds the Potential Biological Removal (PBR) level for this population. Implementing an emergency closure to fishing with buoy lines in this area will address a critical gap in restrictions. This emergency closure in 2023 will reduce entanglement risk where there is a particularly high chance of entanglement that was not addressed in the Plan's 2021 modifications of the MRA (86 FR 51970).

Need	Purpose
To prevent any right whale mortality and serious injury in U.S. commercial fisheries	Reduce the acute risk of entanglement in trap/pot fisheries in a high risk area adjacent to the MRA

3.2.1 Scope of the Analysis

The scope of this analysis is limited to one No Action Alternative (Alternative 1) that would maintain status quo and two alternatives modifying the Plan to create an emergency restricted area to supplement the existing MRA from February 1, 2023 through April 30, 2023. This analysis only affects a small portion of federal waters within Lobster Management Area 1 (LMA 1) in the Northeast portion of the waters covered under the Plan (see the remaining waters within LMA 1 outside of the MRA seasonal closure shown in Figure 1).

4 SUMMARY OF MANAGEMENT ALTERNATIVES

The alternatives were selected based on the results of surveys conducted by the Center for Coastal Studies (CCS) and the Northeast Fisheries Science Center (NEFSC) that observed North Atlantic right whales (right whales) and/or fixed fishing gear adjacent to the Massachusetts Restricted Area (MRA) seasonal closure from February through April of 2018-2022 and modeling conducted using the Large Whale Decision Support Tool (described further in Subsection 6.2).

4.1 Alternative 1: No Action (Status Quo)

Alternative 1, No Action, leaves the current Plan intact with no regulatory changes proposed. This includes the restricted areas implemented by the Final Rule on September 17, 2021 (86 FR 51970) that went into effect October 18, 2021 and requirements for minimum traps per trawl and weak inserts throughout the buoy line that went into effect May 1, 2022.


4.2 Alternative 2: Preferred

Alternative 2, the Preferred Alternative, implements an emergency closure that restricts the use of persistent trap/pot gear buoy lines from February 1, 2023 through April 30, 2023. The closure area is approximately 200 square miles (518 square kilometers) and begins in federal waters east of Cape Ann, is bounded landward by the Massachusetts state waters and south along the 70°30' W longitude line until it intersects with the MRA at the 42°12'N latitude line, and runs west along that line until it intersects the state water boundary (MRA Wedge; Figure 5).

Authorizations for fishing without buoy lines using ropeless gear in the MRA Wedge during this time must be obtained through an Exempted Fishing Permit.

Figure 5: *Alternative 2 (Preferred) Proposed Emergency Closure, MRA Wedge*

The area closed by this emergency rule in the Preferred Alternative, MRA Wedge, is the shaded green area. The existing Massachusetts Restricted Area is the hatched area. The boundary of MRA Wedge would remain closed to trap/pot fishing with buoy lines from February 1, 2023 through April 30, 2023.




4.3 Alternative 3

Alternative 3 also implements an emergency closure that restricts the use of persistent trap/pot gear buoy lines from February 1, 2023 through April 30, 2023. The closed area is approximately 1,297 square miles (3,359 square kilometers) and extends the northern MRA boundaries up to the New Hampshire border at 42°52.58' N (MRA Wedge North; Figure 6). Authorizations for fishing without buoy lines using ropeless gear in the MRA Wedge North during this time must be obtained through an Exempted Fishing Permit.

Figure 6: Alternative 3 Proposed Emergency Closure, MRA Wedge North

The area closed by this emergency rule in Alternative 3, MRA Wedge North to New Hampshire (MRA Wedge North), is the shaded purple area. The existing MRA is the hatched area. The boundary of MRA Wedge North would remain closed to trap/pot fishing with buoy lines from February 1, 2023 through April 30, 2023.



5 DESCRIPTION OF THE AFFECTED ENVIRONMENT

This section describes the valued ecosystem components (VECs) that may be affected by the three alternatives within the Massachusetts portion of Lobster Management Area 1 (MA LMA 1; action area). The proposed action is not expected to have significant impacts on the biological aspects of the fisheries and therefore fish/lobster biology is not included in this analysis.

The three major VECs potentially affected by the proposed action are as follows:

- **Protected Species:** Subsection 5.1 provides information on species listed under the Endangered Species Act of 1973 and/or protected by the Marine Mammal Protection Act of 1972 that may be affected by elements of the proposed action.
- **Habitat:** Subsection 5.2 provides information on marine habitats, with a focus on Essential Fish Habitat. This includes the physical environment and benthic organisms that provide important ecological functions.
- **Human Community:** Subsection 5.3 describes the fisheries as well as the social and economic environment most likely to be impacted by the alternatives under consideration.

5.1 Protected Species

The following discussion examines the potential impacts of management actions on protected species. Table 1 shows the protected species that were considered and identifies which of those may be impacted by the proposed action. NMFS identified five species of Atlantic large whales that are likely to be directly impacted by the implementation of an emergency restricted area (Subsection 5.1.1). Subsection 5.1.1 is further organized by species for information on stock status, distribution, and current threats for North Atlantic right whales (*Eubalaena glacialis*), Gulf of Maine humpback whales (*Megaptera novaeangliae*), fin whales (*Balaenoptera physalus*), sei whales (*Balaenoptera borealis*), and minke whales (*Balaenoptera acutorostrata*). Subsection 5.1.2 provides information on the protected species not likely to be impacted by the proposed action. For more in-depth details on biology, distribution, and documented mortality or serious injury incidents for protected species in the Atlantic Ocean, including Canadian serious mortality or serious injury incidents, please refer to Subsection 4.1 of the 2021 FEIS Volume 1 (NMFS 2021).

Information regarding marine mammal distribution, abundance, potential biological removal (PBR) levels, and sources of mortality and serious injury can be found in the most recent marine mammal Stock Assessment Report (SAR). NOAA prepares marine mammal SARs annually, as directed by the Marine Mammal Protection Act (MMPA). The Draft 2022 SAR was published on January 24, 2023 (88 FR 4162) and open for public comment until April 24, 2023. The information is considered preliminary; however the Draft 2022 SAR has been reviewed by the Atlantic Scientific Review Group and is not expected to change significantly in the final publication. The 2021 SAR was published on August 3, 2022 (87 FR 47385, Hayes et al. 2022).

Information provided in this Environmental Assessment is from either the Draft 2022 SAR or 2021 SAR unless otherwise indicated.¹

Table 1: Species protected under the ESA and/or MMPA that may occur in the Affected Environment

The species and critical habitat that were considered, their current status, and which ones are likely to be impacted by the proposed regulations. Status refers to species status under the Endangered Species Act (ESA), and Protected indicates the species that are protected under the Marine Mammal Protection Act (MMPA). Critical habitat for the North Atlantic right whale is protected under the ESA.

Potential Effect	Category	Species	Status
Potentially Impacted	Marine Mammals	North Atlantic Right Whale	Endangered
		Humpback Whale	Protected (MMPA)
		Fin Whale	Endangered
		Sei Whale	Endangered
		Minke Whale	Protected (MMPA)
	Fish	Giant Manta Ray	Threatened
		Oceanic Whitetip Shark	Threatened
		Atlantic Salmon	Endangered
		Shortnose Sturgeon	Endangered
		Atlantic Sturgeon	New York Bight, Chesapeake Bay, Carolina, and South Atlantic DPSs - Endangered, Gulf of Maine DPS as Threatened
	Marine Mammals	Sperm Whale	Endangered
		Bryde's Whale	Protected (MMPA)
		Harbor Porpoise	Protected (MMPA)
		Blue Whale	Endangered
		WNA Coastal Bottlenose Dolphin	Protected (MMPA)
		Atlantic White-Sided Dolphin	Protected (MMPA)
		Risso's Dolphin	Protected (MMPA)
		Spotted Dolphin	Protected (MMPA)
		Striped Dolphin	Protected (MMPA)

¹ NMFS determined that the Gulf of Maine stock of humpback whales was not strategic for the 2019 Stock Assessment Report (SAR; Hayes et al. 2020), but was strategic for the 2020 SAR because human-caused mortality exceeds PBR. The humpback whale chapter has not been updated since 2019, thus values on population abundance, stock status, and PBR are from the 2019 SAR.

	Pilot Whale	Protected (MMPA)
	Offshore Bottlenose Dolphin	Protected (MMPA)
	Common Dolphin	Protected (MMPA)
	Harbor Seal	Protected (MMPA)
	Gray Seal	Protected (MMPA)
	Harp Seal	Protected (MMPA)
Sea Turtles	Loggerhead Sea Turtle (Northwest Atlantic Ocean DPS)	Threatened
	Leatherback Sea Turtle	Endangered
	Kemp's Ridley Sea Turtle	Endangered
	Green Sea Turtle (North Atlantic DPS)	Threatened
	Hawksbill Sea Turtle	Endangered
	Olive Ridley Sea Turtle	Threatened
Critical Habitat	North Atlantic Right Whale	ESA

5.1.1 Protected Species: Atlantic Large Whales

The primary management objective of the MMPA is to maintain the health and stability of the marine ecosystem, with a goal of obtaining an optimum sustainable population of marine mammals within the carrying capacity of the habitat. Section 118 of the MMPA specifies that NMFS develop and implement Take Reduction Plans to assist in the recovery or prevent the depletion of strategic marine mammal stocks² that interact with Category I and Category II fisheries, which are fisheries with frequent (Category I) or occasional (Category II) mortalities and serious injuries of marine mammals. All marine mammals are protected by the MMPA.

Five species of Atlantic large whales may be present in the affected environment throughout the spring and have the potential to be impacted by the emergency rule: North Atlantic right whales (right whales), Gulf of Maine humpback whales (humpback whales), fin whales, sei whales, and minke whales. These large whales are also known to interact with Category I and II fisheries in the western North Atlantic Ocean and are susceptible to entanglement in trap/pot fishing gear. Fin, sei, and right whales are also listed as endangered under the Endangered Species Act (ESA)

² A strategic stock is defined under the MMPA as a marine mammal stock for which the level of direct human-caused mortality exceeds the potential biological removal level; which, based on the best available scientific information, is declining and is likely to be listed as a threatened species under the ESA within the foreseeable future; or that is listed as a threatened or endangered species under the ESA, or is designated as depleted under the MMPA.

and considered strategic stocks under the MMPA. Although not currently identified as strategic stocks, humpback and minke whales are protected under the MMPA.

North Atlantic Right Whale

The North Atlantic right whale (*Eubalaena glacialis*) is a baleen whale found in temperate and subpolar latitudes in the North Atlantic Ocean. Historic right whale populations were severely depleted by commercial whaling, and despite protections from commercial harvest, the population remains low. Today, they are mainly found in the western North Atlantic (Kraus and Rolland 2007, Monsarrat et al. 2016). For information on right whale distribution outside of the U.S. Exclusive Economic Zone (EEZ), please refer to Subsection 4.1.1.1 of Chapter 4 in the 2021 FEIS (NMFS 2021). Although some individuals are occasionally sighted in the Gulf of Mexico, the current geographic range of right whales within the U.S. EEZ is primarily along the East Coast of North America, from Florida, Georgia, and South Carolina in the south, where calving occurs, through the mid-Atlantic to the coastal waters of Massachusetts to the Gulf of Maine (Morano et al. 2012, NMFS 2013, Wikgren et al. 2014, Oedekoven et al. 2015, Davis et al. 2017, Krzystan et al. 2018). Other than right whales that aggregate in small numbers on the calving grounds in the winter, aggregations are most frequently observed in New England, particularly in southern New England, Cape Cod Bay, and the Gulf of Maine (Wikgren et al. 2014, Davis et al. 2017, Mayo et al. 2018, Quintano-Rizzo et al. 2021) as well as in Canadian waters, such as the Bay of Fundy, Scotian Shelf, and Gulf of Saint Lawrence (Davies et al. 2019, Plourde et al. 2019) where there are sufficient zooplankton patches to support aggregations.

Right whales feed on zooplankton, primarily on copepods, particularly *Calanus finmarchicus*, where they occur in high abundance (Watkins and Schevill 1976, Wishner et al. 1988, Mayo and Marx 1990, Wishner et al. 1995, Woodley and Gaskin 1996, Kenney 2001, Baumgartner et al. 2003, Baumgartner and Mate 2003). In the spring, right whale foraging commonly occurs in Cape Cod Bay (Mayo and Marx 1990), where high densities of *C. finmarchicus* occur. Right whale critical habitat of approximately 29,763 square nautical miles (55,121 kilometers) was designated in 2016 (January 27, 2016, 81 FR 4837; 50 CFR 226).

Shifting *C. finmarchicus* distribution and abundance coincides with changes in spatial distribution and calving rates in right whales (Sorocean et al. 2019). Right whales need to consume large quantities of prey to meet their basic energy requirements and to support population reproduction, migrations, and lactation (Klanjscek et al. 2007, Williams et al. 2013, Meyer-Gutbrod et al. 2015, Irvine et al. 2017). Climate change has already shifted *C. finmarchicus* abundance and phenology in the Gulf of Maine (Record et al. 2019a, 2019b) and model projections suggest resource limitation will likely worsen in the future (Grieve et al. 2017). As prey density and quality shift (namely, reductions in copepod size and nutritional density, while expanding into the northern end of their range), whales need to spend more time foraging and finding areas that have higher quality aggregations of prey. Shifting seasonal patterns and distribution of *C. finmarchicus* throughout the Gulf of Maine (Record et al. 2019a, 2019b), make it challenging to predict locations and timing of aggregations of both right whales and their prey. High abundance of prey species farther north suggests longer travel between calving grounds and feeding grounds, and could contribute further to nutritional stress. Low prey availability also leads to longer interval periods between births (Meyer-Gutbrod and Greene

2018). Lactating females, in particular, appear to be experiencing energy deficits, which could contribute to low reproductive output (Fortune et al. 2013). For more information on distribution of prey and right whale feeding behavior, refer to Subsection 4.1.1.1 of Chapter 4 and Subsection 8.3.3.7 of Chapter 8 in the 2021 FEIS (NMFS 2021).

The right whale is listed as endangered under the ESA. NMFS believes that the right whale is well below the optimum sustainable population level (Hayes et al. 2023). The Draft 2022 Stock Assessment Report (SAR) published by NMFS (Hayes et al. 2023) estimates a minimum population size of 332, and a best estimate of population size to be 338 individuals (95 percent confidence interval: 325-350; Table 2). Potential Biological Removal (PBR) for the North Atlantic right whale population is 0.7 whales per year in the most recently published Draft 2022 SAR (Hayes et al. 2023), down from 0.8 in 2020 (Hayes et al. 2021). Preliminary 2021 and 2022 data suggest the decline has continued and that fewer than 350 individuals remain (Pettis et al. 2022). The estimated total annual estimated mortality for right whales in the U.S. and Canada between 2015 and 2019 is 31.2 (Hayes et al. 2023, Pace 2021). The annual average of observed range-wide total human-caused mortality and serious injury from 2016 to 2020 is 8.1, including 5.7 observed incidental mortalities and serious injuries attributed to fishery interactions and 2.4 observed vessel collisions (Hayes et al. 2023; Table 2), well above PBR. The observed incidental fishery interaction count does not include fishery related serious injuries that were prevented by disentanglement, which is an annual average of 1.2 from 2016 to 2020 (Hayes et al. 2023).

The right whale population is experiencing an Unusual Mortality Event (UME) that began in 2017 due to vessel strikes and entanglement in fishing gear (Daoust et al. 2018, Bourque et al. 2020). Approximately 50 percent of all mortalities and serious injuries since the beginning of the UME have been attributed to entanglements, which make up 68 percent of cases with known cause of death (Hayes et al. 2023). Anthropogenic mortality has limited the recovery of the right whale (Corkeron et al. 2018). While vessel strikes declined after vessel speed regulations were implemented (78 FR 73726; Conn and Silber 2013), both entanglement in fishing gear and vessel strikes remain a significant threat (Kraus et al. 2016, Sharp et al. 2019).

Human-caused mortality heavily influences population demographics (Corkeron et al. 2018). Findings based on the use of a state-space model to estimate abundance of right whales (Pace et al. 2017, Pace 2021) show a strong male survival bias (Pace et al. 2017, Pace 2021). Female right whales may be predisposed to human-caused mortalities because of the increased time spent at the surface in calving grounds and deeper maximum dive depths, which increase their risk for vessel strikes and entanglement in fishing gear, respectively (Dombroski et al. 2021).

Based on the best available information, the greatest entanglement risks to large whales are posed by trap/pot and gillnet fisheries (Angliss and Demaster 1998, Cassoff et al. 2011, Knowlton and Kraus 2001, Hartley et al. 2003, Johnson et al. 2005, Whittingham et al. 2005, Knowlton et al. 2012, Hamilton and Kraus 2019, Sharp et al. 2019, Pace 2021). Specifically, while foraging or transiting, large whales are at risk of becoming entangled in buoy lines and groundlines of trap/pot and gillnet, as well as the net panels of gillnet gear that rise into the water column (Baumgartner et al. 2017, Cassoff et al. 2011, Hamilton and Kraus 2019, Johnson et al. 2005, Knowlton and Kraus 2001, Knowlton et al. 2012, Hayes et al. 2023). Large whale interactions (entanglements) with these features of trap/pot and/or sink gillnet gear often result in

the mortality of or serious injury to the whale (Angliss and Demaster 1998, Cassoff et al. 2011, Henry et al. 2016, Henry et al. 2022, Knowlton and Kraus 2001, Knowlton et al. 2012, Moore and van der Hoop et al. 2013, Sharp et al. 2019, van der Hoop et al. 2016, van der Hoop et al. 2017). Many entanglements, including mortality or serious injury events, go unobserved, and the gear type, fishery, and/or country of origin for reported entanglement events are often not traceable (Henry et al. 2016, Henry et al. 2022). The rates of large whale entanglement, and thus, rates of mortality and serious injury due to entanglement, are likely underestimated (Hamilton et al. 2019, Henry et al. 2016, Henry et al. 2022, Knowlton et al. 2012, Pace et al. 2017, Robbins 2009). Population models estimate that up to 64 percent of right whale mortalities and serious injuries are unobserved (Pace 2021). Additionally, there are mortalities where, despite some evidence of human causes, no cause of death was determined and it is likely that a proportion of these cases also resulted from an entanglement. For a more detailed description of the status and threats to the right whale population in U.S. and Canadian waters see Chapters 2 and 4 in the 2021 FEIS (NMFS 2021).

Table 2: *Estimated abundance, potential biological removal level, and average annual observed mortality for Atlantic Large Whales*

The estimated abundance, potential biological removal level, and average annual observed mortality for Atlantic large whale species likely to be impacted by the proposed action. Refer to the 2022 Draft U.S. Atlantic and Gulf of Mexico Marine Mammal Stock Assessment Report (SAR) for a full summary of abundance and mortality levels (Hayes et al. 2023). The humpback whale chapter has not been updated since 2019, thus values on population abundance, stock status, and PBR are from the 2019 SAR (Hayes et al. 2020) and average annual observed mortality is from Henry et al. (2022). Because observed mortalities and serious injuries only represent a fraction of observed cases, some of these species may be experiencing human-caused mortalities at higher rates once unobserved mortalities are taken into account.

Species	Estimated Abundance	Potential Biological Removal Level	Average Annual Observed Mortality
Right Whale	338	0.7	8.1
Humpback Whale	1,396	22	16.25
Fin Whale	6,802	11	1.8
Sei Whale	6,292	6.2	0.8
Minke Whales	21,968	170	10.55

Gulf of Maine Humpback Whale

The Gulf of Maine humpback whale (formerly Western North Atlantic) was previously listed as endangered under the ESA. In 2016, several distinct population segments were removed from listing, including the West Indies distinct population segment. The Gulf of Maine stock is largely composed of whales that reproduce in the West Indies (81 FR 62259, September 2016). The Gulf of Maine stock is still protected under the MMPA.

Since the early 1990s, humpbacks, particularly juveniles, have been observed stranded dead with increasing frequency in the mid-Atlantic (Swingle et al. 1993, Wiley et al. 1995) and have been sighted in wintertime surveys in the Southeast and mid-Atlantic (Hayes et al. 2020). In the Gulf

of Maine, sightings are most frequent from mid-March through November, with a peak in May and August, from the Great South Channel east of Cape Cod northward to Stellwagen Bank and Jeffreys Ledge (CETAP 1982). Acoustic detections of humpbacks indicate year-round presence in New England waters, including the waters of Stellwagen Bank (Davis et al. 2020).

Distribution in these waters appears to be correlated with prey species, including herring (*Clupea harengus*), sand lance (*Ammodytes spp.*), and other small fishes, as well as euphausiids (Paquet et al. 1997). Changes in humpback distribution in the Gulf of Maine have been found to be associated with changes in herring, mackerel, and sand lance abundance associated with local fishing pressures (Payne et al. 1986).

Current data suggest that the Gulf of Maine humpback whale stock is increasing (Hayes et al. 2020). The most recent population estimate calculated an abundance of 1,396 animals (Table 2) in this stock and the minimum population estimate is 1,380 (Hayes et al. 2020). The maximum productivity rate is 0.065 and the “recovery” factor is assumed to be 0.50, the default for stocks of unknown status, because the listing for the distinct population segment was removed in 2016. Thus, the PBR for the Gulf of Maine humpback whale stock is 22 whales per year (Hayes et al. 2020).

As with right whales, the primary known sources of anthropogenic mortality and serious injury of humpback whales are commercial fishing gear entanglements and ship strikes. Robbins et al. (2009) found that humpback whales experience new scarring at an annual rate of 12.1 percent. From 2010 to 2019, 38.8 percent of all observed mortalities and serious injuries were attributed to entanglements from interactions with trap/pot, monofilament line, netting, and unidentified gear (see Chapter 2 of 2021 FEIS, NMFS 2021). From 2015 through 2019, observed human-caused mortalities averaged 16.25 animals per year, with 9.35 incidental fishery interactions and 6.9 vessel collisions (Henry et al. 2022). These results include only observed mortality and serious injury. Unobserved anthropogenic impacts on humpback whales is likely, but has not been calculated to date. A UME was declared in 2017 after a spike in humpback whale strandings along the East Coast of the U.S. and 40 percent of the cases where cause of death was examined had evidence of ship strike or entanglement (<https://www.fisheries.noaa.gov/national/marine-life-distress/2016-2023-humpback-whale-unusual-mortality-event-along-atlantic-coast>).

Fin Whale

The fin whale is found in all major oceans and was considered to be composed of three subspecies until recently: *Balaenoptera physalus* in the Northern Hemisphere, and *B. p. quoyi* and *B. p. patachonica* (a pygmy form) in the Southern Hemisphere. New genetic data suggest that fin whales in the North Atlantic and North Pacific oceans represent two different subspecies (Archer et al. 2019). The International Whaling Commission defines a single stock of the North Atlantic fin whale off the eastern coast of the U.S., north to Nova Scotia, and east to the southeastern coast of Newfoundland (Donovan 1991). Fin whales are common in the waters of the U.S. EEZ, principally from Cape Hatteras northward (Hayes et al. 2022). In a globally scaled review of sightings data, Edwards et al. (2015) found evidence to confirm the presence of fin whales in every season throughout much of the U.S. EEZ north of 35° N; however, densities vary seasonally. Acoustic detections of fin whale singers in Massachusetts Bay, New York Bight, and

deep-ocean areas confirm whale presence September through June throughout the western North Atlantic (Watkins et al. 1987, Clark and Gagnon 2002, Morano et al. 2012). Davis et al. 2020 detected year-round acoustic presence of fin whales within the EEZ, particularly in areas north of Cape Hatteras, North Carolina.

Of the three to seven stocks thought to occur in the North Atlantic Ocean, one occurs in U.S. waters, where NMFS best estimate of abundance is 6,802 individuals (Hayes et al. 2023; Table 2). The maximum productivity rate is 0.04, the default value for cetaceans. The “recovery” factor is assumed to be 0.1 because the fin whale is listed as endangered under the ESA. Thus, PBR for the western North Atlantic fin whale is 11 (Hayes et al. 2023). The species’ overall population size may provide some resilience to current threats, but trends remain largely unknown (Hayes et al. 2022).

Documented sources of anthropogenic mortality of fin whales include entanglement in commercial fishing gear and ship strikes. Additional threats include reduced prey availability and anthropogenic sound. Experts believe that fin whales are struck by large vessels more frequently than any other cetaceans (Laist et al. 2001). Approximately 22.7 percent of all observed mortality and serious injury were attributed to entanglements between 2010 and 2019, with most interactions occurring with trap/pot and unidentified gear (see Chapter 2 of 2021 FEIS, NMFS 2021). The minimum annual rate of anthropogenic mortality and serious injury to fin whales, between 2015 and 2019, was 1.85 per year, 1.45 of those from fishing entanglement, and 0.4 per year from ship strikes (Hayes et al. 2022, Henry et al. 2022).

Sei Whale

Sei whales are listed as endangered throughout their range under the ESA. The western North Atlantic sei whale population belongs to the Northern Hemisphere subspecies (*B. b. borealis*) and consists of two stocks, a Nova Scotian Shelf stock and a Labrador Sea stock (Baker and Clapham 2004, Mitchell and Chapman 1977). The Nova Scotian Shelf stock is the only sei whale stock within ALWTRP boundaries and ranges from the U.S. East Coast to Cape Breton, Nova Scotia and east to 42°00’W longitude (Hayes et al. 2022). The Nova Scotia stock in the North Atlantic is estimated to be 6,292 individuals with a minimum population size of 3,098 individuals (Hayes et al. 2023; Table 2). Population growth rates for sei whales are not available at this time as there are little to no systematic survey efforts to study sei whales.

Sei whales are often found in the deeper waters that characterize the edge of the continental shelf (Hain et al. 1985), but NMFS aerial surveys also found substantial numbers of sei whales south of Nantucket in spring and summer (Stone et al. 2017) and on Georges Bank in the spring and summer (CETAP 1982). Sei whales have also been documented inshore, near the Great South Channel (in 1987 and 1989) and Stellwagen Bank (in 1986; Payne et al. 1990). Davis et al. 2020 detected sei whale acoustic presence along the U.S. and Canadian East Coast year round, with the highest detections north of Cape Hatteras, North Carolina during the spring through fall. Sei whales (like right whales) are largely planktivorous, primarily feeding on euphausiids and copepods, which has resulted in reports of sei whales in more inshore locations. Sei whales are also opportunistically piscivorous, consuming species of small schooling fish and squid (Wiles 2017, Prieto et al. 2012).

Current threats include vessel strikes, fisheries interactions (including entanglement), climate change, habitat loss, reduced prey availability, and anthropogenic sound. Between 2010 and 2019, 18 serious injuries and mortalities were observed: 8 with unknown causes, 5 vessel strikes (all confirmed U.S.), 2 entanglements, and 3 non-human caused mortality (see Chapter 2 of 2021 FEIS, NMFS 2021). Between 2015 and 2019, the average annual rate of confirmed human-caused mortality and serious injury to sei whales is 0.8 incidents per year (Hayes et al. 2022; Table 2). This value includes incidental fishery interaction records (0.4), records of vessel collisions (0.2), and other human-induced mortalities (0.2). Possible causes of natural mortality, particularly for compromised individuals, are shark attacks, killer whale attacks, and endoparasitic helminthes (Perry et al. 1999).

Minke Whale

The minke whale is not listed as endangered or threatened under the ESA but is protected under the MMPA. Minke whales off the East Coast of the U.S. are considered to be part of the Canadian East Coast population, which inhabits the area from the eastern half of Davis Strait south to the Gulf of Mexico. They are common and widely distributed within the U.S. Atlantic EEZ (CETAP 1982). Minke whales are most frequently observed in New England waters from spring to fall, and acoustic surveys have commonly detected their presence on the shelf (Hayes et al. 2022, Risch et al. 2013). Acoustic detections in Stellwagen Bank peaked in fall and winter (September to December), while detections off New York appear to peak in spring (Risch et al. 2014). Where recording effort spanned the shelf, acoustic detections were highest near the shelf break and deep waters.

Data are insufficient for determining a population trend for this species. The best estimate of population size is 21,968 (CV=0.31) minke whales, with the minimum population size of 17,002 (Hayes et al. 2023; Table 2). The observed annual estimated average human-caused mortality and serious injury for the Canadian East Coast stock of minke whales is 10.55, including 9.55 mortalities due to incidental fishery interactions, 0.2 from observed fishery interaction, and 0.8 caused by vessel collisions (Hayes et al. 2022).

As with other large whales, documented sources of anthropogenic mortality of minke whales include entanglement in commercial fishing gear and vessel strikes. Minke whales have been entangled in a variety of fishing gear, including unspecified fishing nets, unspecified cables or lines, fish traps, weirs, seines, gillnets, and lobster gear. Between 2010 and 2019, nearly 30 percent of all observed mortalities and serious injuries were attributed to entanglements, most of which resulted from interactions with trap/pot, netting, and unidentified gear (see Chapter 2 for the 2021 FEIS, NMFS 2021). A UME was declared in 2017 following an uptick in strandings along the East Coast of the U.S. Though the specific cause of the high mortality has not been determined, several stranded whales have shown evidence of human interaction (<https://www.fisheries.noaa.gov/national/marine-life-distress/2017-2022-minke-whale-unusual-mortality-event-along-atlantic-coast>).

5.1.2 Species and Critical Habitat Not Likely to be Impacted

Based on the best available information, Table 1 provides a list of species not likely to be impacted by the proposed action. This determination has been made because either the occurrence of the species has either limited or no overlap with the trap/pot fisheries operating in the proposed action area and/or interactions have never been documented or are extremely rare between the species and trap/pot gear (see Marine Mammal Stocks Assessment Reports at <https://www.fisheries.noaa.gov/national/marine-mammal-protection/marine-mammal-stock-assessment-reports-region>; Hayes et al. 2022; Sea Turtle Disentanglement Network, unpublished data; NMFS Observer Program, unpublished data; see OBIS-SMAP at <https://seamap.env.duke.edu/>). The proposed actions will not affect the essential physical and biological features of critical habitat designated for North Atlantic right whales, the Northwest Atlantic Ocean DPS of loggerhead sea turtle, or the Gulf of Maine DPS of Atlantic salmon. Therefore, the proposed action will not result in the destruction or adverse modification of any designated critical habitat (NMFS 2014, NMFS 2015a, 2015b).

5.2 Habitat

Modification of the Plan may affect Essential Fish Habitat (EFH), which is defined by the Magnuson-Stevens Act as “those waters and substrate necessary to fish for spawning, breeding, feeding or growth to maturity” (16 U.S.C. 1802(10)). Regulations developed by NMFS encourage Regional Fishery Management Councils to describe and identify EFH, and, to the extent practicable, to minimize adverse effects caused by fishing activities. Atlantic trap/pot fisheries are geographically widespread on the Atlantic coast and target a diverse array of fish and shellfish species. In the context of this Environmental Assessment, EFH includes the habitat for all non-target species during relevant life history stages that take place within the proposed area (Figure 7; Table 3). Because this action is not expected to affect pelagic habitats, the species and life stages listed in Table 3 are all benthic. For detailed discussion of Essential Fish Habitat and Habitat Areas of Particular Concern regulatory requirements, key components of lobster habitat in detail, and how the Atlantic Large Whale Take Reduction Plan can influence habitat, reference Subsection 4.2 of the 2021 FEIS (NMFS 2021).

Figure 7: Habitat Areas of Particular Concern and Essential Fish Habitat in Action Area

The Habitat Areas of Particular Concern (HAPC) and Essential Fish Habitat (EFH) currently protected from fishing within the proposed area, including those overseen by the New England Fishery Management Council (NEFMC).




Table 3: List of Essential Fish Habitat for different species and life history stages within Affected Environment

Species	Life Stage	Depth (meters)	Habitat Type and Description
Acadian redfish	Juveniles	50-200 in Gulf of Maine, to 600 on slope	Sub-tidal coastal and offshore rocky reef substrates with associated structure-forming epifauna (e.g., sponges, corals), and soft sediments with cerianthid anemones
American plaice	Juveniles	40-180	Subtidal benthic habitats on mud and sand, also found on gravel and sandy substrates bordering bedrock
American plaice	Adults	40-300	Subtidal benthic habitats on mud and sand, also gravel and sandy substrates bordering bedrock
Atlantic cod	Juveniles	Mean high water-120	Structurally-complex intertidal and subtidal habitats, including eelgrass, mixed sand and gravel, and rocky habitats (gravel pavements, cobble, and boulder) with and without attached macroalgae and emergent epifauna

Atlantic cod	Adults	30-160	Structurally complex sub-tidal hard bottom habitats with gravel, cobble, and boulder substrates with and without emergent epifauna and macroalgae, also sandy substrates and along deeper slopes of ledges
Atlantic halibut	Juveniles & Adults	60-140 and 400-700 on slope	Benthic habitats on sand, gravel, or clay substrates
Atlantic herring	Eggs	5-90	Subtidal benthic habitats on coarse sand, pebbles, cobbles, and boulders and/or macroalgae
Atlantic sea scallop	Eggs	18-110	Inshore and offshore benthic habitats (see adults)
Atlantic sea scallop	Larvae	No information	Inshore and offshore pelagic and benthic habitats: pelagic larvae (“spat”), settle on variety of hard surfaces, including shells, pebbles, and gravel and to macroalgae and other benthic organisms such as hydroids
Atlantic sea scallop	Juveniles	18-110	Benthic habitats initially attached to shells, gravel, and small rocks (pebble, cobble), later free-swimming juveniles found in same habitats as adults
Atlantic sea scallop	Adults	18-110	Benthic habitats with sand and gravel substrates
Atlantic surfclams	Juveniles and adults	Surf zone to about 61, abundance low >38	In substrate to depth of 3 ft
Atlantic wolffish	Eggs	<100	Subtidal benthic habitats under rocks and boulders in nests
Atlantic wolffish	Juveniles	70-184	Subtidal benthic habitats
Atlantic wolffish	Adults	<173	A wide variety of sub-tidal sand and gravel substrates once they leave rocky spawning habitats, but not on muddy bottom
Black sea bass	Juveniles and adults	Inshore in summer and spring	Benthic habitats with rough bottom, shellfish and eelgrass beds, man-made structures in sandy-shelly areas, also offshore clam beds and shell patches in winter
Haddock	Juveniles	40-140 and as shallow as 20 in coastal Gulf of Maine	Subtidal benthic habitats on hard sand (particularly smooth patches between rocks), mixed sand and shell, gravelly sand, and gravel
Haddock	Adults	50-160	Subtidal benthic habitats on hard sand (particularly smooth patches between rocks), mixed sand and shell, gravelly sand, and gravel and adjacent to boulders and cobbles along the margins of rocky reefs
Little skate	Juveniles	Mean high water-80	Intertidal and subtidal benthic habitats on sand and gravel, also found on mud
Little skate	Adults	Mean high water-100	Intertidal and subtidal benthic habitats on sand and gravel, also found on mud
Monkfish	Juveniles	50-400 in the Mid-Atlantic, 20-400 in the Gulf of Maine, and to 1000 on the slope	Subtidal benthic habitats on a variety of habitats, including hard sand, pebbles, gravel, broken shells, and soft mud, also seek shelter among rocks with attached algae
Monkfish	Adults	50-400 in the Mid-Atlantic, 20-400 in the Gulf of Maine, and to 1000 on the slope	Subtidal benthic habitats on hard sand, pebbles, gravel, broken shells, and soft mud, but seem to prefer soft sediments, and, like juveniles, utilize the edges of rocky areas for feeding
Ocean pout	Eggs	<100	Sub-tidal hard bottom habitats in sheltered nests, holes, or rocky crevices

Ocean pout	Juveniles	Mean high water-120	Intertidal and subtidal benthic habitats on a wide variety of substrates, including shells, rocks, algae, soft sediments, sand, and gravel
Ocean pout	Adults	20-140	Subtidal benthic habitats on mud and sand, particularly in association with structure forming habitat types; i.e. shells, gravel, or boulders
Ocean quahogs	Juveniles and adults	9-244	In substrate to depth of 3 ft
Offshore hake	Juveniles	160-750	Pelagic and benthic habitats
Pollock	Juveniles	Mean high water-180 in Gulf of Maine, Long Island Sound, and Narragansett Bay; 40-180 on Georges Bank	Intertidal and subtidal pelagic and benthic rocky bottom habitats with attached macroalgae, small juveniles in eelgrass beds, older juveniles move into deeper water habitats also occupied by adults
Pollock	Adults	80-300 in Gulf of Maine and on Georges Bank; <80 in Long Island Sound, Cape Cod Bay, and Narragansett Bay	Pelagic and benthic habitats on the tops and edges of offshore banks and shoals with mixed rocky substrates, often with attached macro algae
Red hake	Juveniles	Mean high water-80	Intertidal and subtidal soft bottom habitats, esp those that provide shelter, such as depressions in muddy substrates, eelgrass, macroalgae, shells, anemone and polychaete tubes, on artificial reefs, and in live bivalves (e.g., scallops)
Red hake	Adults	50-750 on shelf and slope, as shallow as 20 inshore	Subtidal benthic habitats in shell beds, on soft sediments (usually in depressions), also found on gravel and hard bottom and artificial reefs
Scup	Juveniles	No information	Benthic habitats, in association with inshore sand and mud substrates, mussel and eelgrass beds
Scup	Adults	No information, generally overwinter offshore	Benthic habitats
Silver hake	Juveniles	40-400 in Gulf of Maine, >10 in Mid-Atlantic	Pelagic and sandy subtidal benthic habitats in association with sand-waves, flat sand with amphipod tubes, shells, and in biogenic depressions
Silver hake	Adults	>35 in Gulf of Maine, 70-400 on Georges Bank and in the Mid-Atlantic	Pelagic and sandy subtidal benthic habitats, often in bottom depressions or in association with sand waves and shell fragments, also in mud habitats bordering deep boulder reefs, on over deep boulder reefs in the southwest Gulf of Maine
Smooth skate	Juveniles	100-400 offshore Gulf of Maine, <100 inshore Gulf of Maine, to 900 on slope	Benthic habitats, mostly on soft mud in deeper areas, but also on sand, broken shells, gravel, and pebbles on offshore banks in the Gulf of Maine
Smooth skate	Adults	100-400 offshore Gulf of Maine, to 900 on slope	Benthic habitats, mostly on soft mud in deeper areas, but also on sand, broken shells, gravel, and pebbles on offshore banks in the Gulf of Maine
Summer flounder	Juveniles	To maximum 152	Benthic habitats, including inshore estuaries, salt marsh creeks, seagrass beds, mudflats, and open bay areas
Summer flounder	Adults	To maximum 152 in colder months	Benthic habitats
Thorny skate	Juveniles	35-400 offshore Gulf of Maine, <35 inshore Gulf of Maine, to 900 on slope	Benthic habitats on a wide variety of bottom types, including sand, gravel, broken shells, pebbles, and soft mud

Thorny skate	Adults	35-400 offshore Gulf of Maine, <35 inshore Gulf of Maine, to 900 on slope	Benthic habitats on a wide variety of bottom types, including sand, gravel, broken shells, pebbles, and soft mud
White hake	Juveniles	Mean high water-300	Intertidal and subtidal estuarine and marine habitats on fine-grained, sandy substrates in eelgrass, macroalgae, and un-vegetated habitats
White hake	Adults	100-400 offshore Gulf of Maine, >25 inshore Gulf of Maine, to 900 on slope	Subtidal benthic habitats on fine-grained, muddy substrates and in mixed soft and rocky habitats
Windowpane flounder	Juveniles	Mean high water-60	Intertidal and subtidal benthic habitats on mud and sand substrates
Windowpane flounder	Adults	Mean high water-70	Intertidal and subtidal benthic habitats on mud and sand substrates
Winter flounder	Eggs	0-5 south of Cape Cod, 0-70 Gulf of Maine and Georges Bank	Sub-tidal estuarine and coastal benthic habitats on mud, muddy sand, sand, gravel, submerged aquatic vegetation, and macroalgae
Winter flounder	Juveniles	Mean high water-60	Intertidal and subtidal benthic habitats on a variety of bottom types, such as mud, sand, rocky substrates with attached macro algae, tidal wetlands, and eelgrass; young-of-the-year juveniles on muddy and sandy sediments in and adjacent to eelgrass and macroalgae, in bottom debris, and in marsh creeks
Winter flounder	Adults	Mean high water-70	Intertidal and subtidal benthic habitats on muddy and sandy substrates, and on hard bottom on offshore banks; for spawning adults, also see eggs
Winter skate	Juveniles	0-90	Subtidal benthic habitats on sand and gravel substrates, are also found on mud
Winter skate	Adults	0-80	Subtidal benthic habitats on sand and gravel substrates, are also found on mud
Witch flounder	Juveniles	50-400 and to 1500 on slope	Subtidal benthic habitats with mud and muddy sand substrates
Witch flounder	Adults	35-400 and to 1500 on slope	Subtidal benthic habitats with mud and muddy sand substrates
Yellowtail flounder	Juveniles	20-80	Subtidal benthic habitats on sand and muddy sand
Yellowtail flounder	Adults	25-90	Subtidal benthic habitats on sand and sand with mud, shell hash, gravel, and rocks

A reduction in fishing effort is likely to decrease the time that fishing gear is in the water, thereby reducing the potential for interactions between fishing gear and habitat. However, most habitat areas where lobsters are fished have been heavily fished by multiple fishing fleets over many decades and are unlikely to see a measurable improvement in their condition in response to a short-term decrease in effort of an individual fishery.

Experts believe that fixed fishing gear (e.g. traps/pots) has a more direct impact on benthic habitat than on non-benthic (water column) habitat because it generally comes in contact with the seafloor. Therefore, the sections below review how fixed-gear fishing can affect habitat, with a primary focus on benthic habitat. The potential effects examined include:

- Alteration of physical structure;
- Mortality of benthic organisms;
- Changes to the benthic community and ecosystem;

- Sediment suspension; and,
- Chemical modifications.

5.2.1 Alteration of Physical Structure

Any type of fishing gear that is towed, dragged, or dropped on the seabed will disturb the sediment and the resident community to varying degrees. The intensity of disturbance is dependent on the type of gear, how long the gear is in contact with the bottom, sediment type, sensitivity of habitat features in contact with the gear, and frequency of disturbance. Physical effects of fishing gear, such as plowing, smoothing of sand ripples, removal of stones, and turning of boulders, can act to reduce the heterogeneity of the sediment surface. For example, boulder piles, crevices, and sand ripples can provide fish and invertebrates hiding areas and a respite from currents and tides. Removal of taxa, such as tube worms, corals, and gorgonians that provide relief, and the removal or shredding of submerged vegetation, can also occur, thereby reducing the number of structures available to biota as habitat.

Most studies on habitat damage due to fishing gear focus on the effects of bottom trawls and dredges. It has been noted by Rogers et al. (1998) that the reason there are few accounts of static gear (e.g. traps/pots) having measurable effects on benthic biota may be because the area of seabed affected by such gear is almost insignificant when compared to the widespread effects of mobile gear. It is possible that benthic structures (both living and non-living) could be affected as traps/pots are dropped or dragged along the bottom. Most studies investigating small numbers of traps or pots per buoy line (1-3) have found minimal, short-term impacts on physical structures (Eno et al. 2001, Chuenpagdee et al. 2003, Stephenson et al. 2017). Similarly, a panel of experts that evaluated the habitat impacts of commercial fishing gears used in the Northeast of the U.S. (Maine to North Carolina) found bottom-tending static gear (e.g. traps/pots) to have a minimal effect on benthic habitats when compared to the physical and biological impacts caused by bottom trawls and dredges (NMFS 2002). The vulnerability of benthic EFH for all managed species in the region to the impacts of pots/traps and bottom gillnets is considered to be low (NMFS 2004). However, less is known about longer trap/pot trawls and there is limited information that trawls with 20 or more pots may have impacts more similar to mobile gear, though at a smaller spatial scale (Schweitzer et al. 2018).

5.2.2 Mortality of Benthic Organisms

In addition to effects on physical habitat, fishing gear can cause direct mortality to emergent epifauna. In particular, erect, foliose fauna, or fauna that build reef-like structures have the potential to be destroyed by towed gear, longlines, or traps/pots (Hall 1999). Physical structure of the biota sometimes determines their ability to withstand and recover from the physical impacts of fishing gear. For example, thinner shelled bivalves and sea stars often suffer higher damage than solid shelled bivalves (Rumohr and Krost 1991). Animals that can retract below the penetration depth of the fishing gear and those that are more elastic and can bend upon contact with the gear also fare much better than those that are hard and inflexible (Eno et al. 2001). Longer trap/pot trawls likely pose a greater threat to benthic organisms than individual trap/pots or short trap/pot trawls (Schweitzer et al. 2018).

5.2.3 Changes to Benthic Communities and Ecosystems

The mortality of benthic organisms as a result of interaction with fishing gear can alter the structure of the benthic community, potentially causing a shift in the community from low productive long-lived species to highly-productive, short-lived, rapidly-colonizing species. For example, motile species that exhibit high fecundity and rapid generation times will recover more quickly from fishery-induced disturbances than non-mobile, slow-growing organisms, which may lead to a community shift in chronically fished areas (Levin 1984).

Increased fishing pressure in a certain area may also lead to changes in species distribution. Changes (e.g., localized depletion) could be evident in benthic, demersal, and even pelagic species. Scientists have also speculated that mobile fishing may lead to increased populations of opportunistic feeders in chronically fished areas.

5.2.4 Sediment Suspension

Resuspension of sediment can occur as fishing gear is pulled or dragged along or immediately above the seafloor (NMFS 2002). Although resuspension of sediment is typically associated with mobile fishing gear, it also can occur with gear such as traps/pots.

Chronic suspension of sediments and resulting turbidity can affect aquatic habitat by reducing available light for photosynthesis, burying benthic biota, smothering spawning areas, and causing negative effects on feeding and metabolic rates. If it occurs over large areas, resuspension can redistribute sediments, which has implications for nutrient budgets (Mayer et al. 1991, Messieh et al. 1991, Black and Parry 1994, Pilskaln et al. 1998).

Species' reaction to turbidity depends on the particular life history characteristics of the organism. Effects are likely to be more significant in waters that are normally clear as compared to areas that typically experience high natural turbidity (Kaiser 2000). Mobile organisms can move out of the affected area and quickly return once the turbidity dissipates (Coen 1995). Even if species experience high mortality within the affected area, those with high levels of recruitment or high mobility can re-populate the affected area rapidly. However, sessile or slow-moving species would likely be buried and could experience high mortality. Furthermore, if effects are protracted and occur over a large area, recovery through recruitment or immigration will be hampered. Additionally, chronic resuspension of sediments may lead to shifts in species composition by favoring those species that are better suited to recover or those that can take advantage of the additional nutrient supply as the nutrients are released from the seafloor to the euphotic zone (Churchill 1989).

5.2.5 Chemical Modifications

Disturbances associated with fishing gear also can cause changes in the chemical composition of the water column overlying affected sediments. In shallow water, the impacts may not be noticeable relative to the mixing effects caused by tidal surges, storm surges, and wave action. However, in deeper, calmer areas with more stable waters, the changes in chemistry may be more evident (NMFS 2002). Increases in ammonia content, decreases in oxygen, and pulses of

phosphate have been observed in North Sea waters, although it is not clear how these changes affect fish populations. Increased incidence of phytoplankton blooms could occur during seasons when nutrients are typically low. The increase in primary productivity could have a positive effect on zooplankton communities and on organisms up the food chain.

Eutrophication, often considered a negative effect, could also occur. However, it is important to note that these releases of nutrients to the water recycle existing nutrients and make them available to benthic organisms (ICES 1992). This recycling is thought to be less influential in the eutrophication process than the input of new nutrients from rivers and land runoff.

5.2.6 American Lobster Habitat

Bottom dwelling American lobster (*Homarus americanus*) is distributed throughout the Northwest Atlantic Ocean from Newfoundland to Cape Hatteras, North Carolina. Juvenile and adult American lobsters occupy a wide variety of benthic habitats from the intertidal zone to depths of 2300 feet (700 meters). They are most abundant in relatively shallow coastal waters. Temperature and salinity, as well as substrate and diet, are critical habitat components (ASMFC 2015). Lobsters feed on a variety of plants and animals according to seasonal availability, and bait in lobster traps is believed to be an important food source in areas of intense fishing pressure ((Lawton and Lavalli 1995, Grabowski et al. 2010) cited in ASMFC 2015).

The affected area includes the Massachusetts portion of Lobster Management Area 1 (MA LMA 1), including Massachusetts Bay (Alternative 2 and Alternative 3), Ipswich Bay, and other waters offshore of northern Massachusetts (Alternative 3). Water depth ranges from 3.3 feet (1 meter) to 656.2 feet (200 meters) (CZM 1999). Within this area, the affected habitat can be further categorized into inshore and offshore lobster habitat. A full description of lobster habitat that includes estuarine inshore and offshore canyon in addition to rock inshore and other offshore lobster habitats can be found in Chapter 4 of the 2021 FEIS (NMFS 2021).

Inshore estuarine and rock areas make up two key components of inshore lobster habitat. For the purpose of this proposed action, only the inshore rock areas are included within this discussion because inshore estuarine areas are outside the scope of the action area. Inshore rock habitat areas for lobster include the following:

- **Mud Base with Burrows:** These habitats occur primarily in harbors and quiet estuaries with low currents. Lobster shelters are formed from excavations in soft substrate. This is an important habitat for juveniles, and densities can be very high, reaching 20 animals per square meter (per square 3.3 feet).
- **Rock, Cobble, and Gravel:** Juveniles and adolescents have been reported on shallow bottom with gravel and gravelly sand substrates in the Great Bay Estuary, New Hampshire; on gravel/cobble substrates in outer Penobscot Bay, Maine (Steneck and Wilson 1998); and in rocky habitats in Narragansett Bay, Rhode Island (Lawton and Lavalli 1985). Densities in Penobscot Bay exceeded 0.5 juveniles and 0.75 adolescents per square meter (per square 3.3 feet). According to unpublished information cited by Lincoln (1998), juvenile lobsters in Great Bay prefer shallow bottoms with gravelly sand substrates.

- **Rock/Shell:** Adult lobsters in the Great Bay Estuary utilize sand and gravel habitats in the channels, but appear to prefer a rock/shell habitat more characteristic of the high temperature, low salinity regimes of the central bay.
- **Sand Base with Rock:** This is the most common inshore rock type in depths greater than 40 meters. It consists of sandy substrate overlain by flattened rocks, cobbles, and boulders. Lobsters are associated with abundant sponges, Jonah crabs, and rock crabs. Lobsters excavate sand under a rock to form U-shaped, shallow tunnels for shelter. Densities of sub-adult lobsters are fairly high in these areas.
- **Boulders Overlaying Sand:** This habitat type is relatively rare in inshore New England waters. Compared to other inshore rocky habitats, lobster densities are low.
- **Cobbles:** Lobsters occupy shelters of varying size in the spaces between rocks, pebbles, and boulders. Densities as high as 16 lobsters per square meter (per square 3.3 feet) have been observed, making this the most densely populated inshore rock habitat for lobsters in New England.
- **Bedrock Base with Rock and Boulder Overlay:** This rock type is relatively common inshore, from low tide to depths of 49 to 148 feet (15 to 45 meters). Shelters are formed by rock overhangs or crevices. Encrusting coralline algae and attached organisms such as anemones, sponges, and mollusks cover exposed surfaces. Green sea urchins and starfish are common. Cunner, tautog, sculpin, sea raven, and redfish are the most abundant fish. Lobster densities generally are low.
- **Mud-Shell/Rock Substrate:** This habitat type is usually found where sediment discharge is low and shells make up the majority of the bottom. Lobster densities in this habit type are generally low.

Other lobster habitat types are significant. For example, kelp beds represent another form of lobster habitat. Kelp beds in New England consist primarily of *Laminaria longicruris* and *L. saccharina*. Lobsters were attracted to transplanted kelp beds at a nearshore study site in the midcoast region of Maine, reaching densities almost ten times higher than in nearby control areas (Bologna and Steneck 1993). Lobsters did not burrow into the sediment, but sought shelter beneath the kelp. Only large kelp (greater than 1.6 feet (50 cm) in length) was observed sheltering lobsters and was used in the transplant experiments.

Lobster shelters are formed from excavations cut into peat. Reefs form from blocks of salt marsh peat that break and fall into adjacent marsh creeks and channels. The reefs appear to provide moderate protection for small lobsters from predators (Barshaw and Lavalli 1988). Densities are high—up to 61.4 square feet (5.7 square meters) in these areas.

Offshore lobster habitats can be subdivided into canyons and other offshore habitats. The canyon offshore lobster habitats are beyond the scope of the proposed emergency closures. Other offshore habitat includes the following:

- **Sand Base with Rocks:** Although common inshore, this habitat is rather restricted in the offshore region except along the north flank of Georges Bank.
- **Clay Base with Burrows and Depressions:** This habitat is common on the outer continental shelf and slope. Lobsters excavate burrows up to 4.9 feet (1.5 meters) long. There are also large, bowl-like depressions that range in size from one to five meters in

diameter and may shelter several lobsters at a time. Minimum densities of 0.001 lobsters per square meter (per square 3.3 feet) have been observed in summer.

- **Mud-Clay Base with Anemones:** This is a common habitat for lobsters on the outer shelf or upper slope. Forests of mud anemones (*Cerianthus borealis*) may reach densities of three or four per square meter. Depressions serve as shelter for relatively small lobsters at minimum densities of 0.001 per square meter (per square 3.3 feet).
- **Mud Base with Burrows:** This habitat occurs offshore mainly in the deep basins, in depths up to 821 feet (250 meters). This environment is extremely common offshore. Lobsters occupy this habitat, but no density estimates are available.

5.3 Human Community

5.3.1 Affected Fisheries

American Lobster

The American lobster (*Homarus americanus*) is a bottom-dwelling, marine crustacean characterized by a large shrimp-like body and ten legs, two of which are enlarged to serve as crushing and gripping appendages. The American lobster range extends from Newfoundland south to the Mid-Atlantic region. In U.S. waters, the species is most abundant from the inshore waters of Maine to Cape Cod, Massachusetts, and abundance declines from north to south (ASMFC 2015). In Massachusetts, the trap/pot fishery has consistently landed about 17 million pounds of live lobsters per season in the past few years. Based on federal Vessel Trip Report (VTR) data from 2017 to 2021, most of the vessels affected by the action alternatives fished from the ports of Gloucester, Rockport, and Beverly in Essex County, Cohasset in Norfolk County, and Scituate in Plymouth County.³ However, human communities can extend beyond the boundaries of a particular port or city, so our analysis focuses on the county level. Essex and Plymouth county land the most lobsters, Barnstable and Bristol County also land a significant amount of lobsters, while Suffolk and Norfolk land a small fraction of the total amount. Table 4 displays the lobster landing pounds by county in Massachusetts from 2017 to 2021.

³ Vessels that only have lobster permits are not required to submit VTR; therefore, there is some uncertainty in quantifying the number of affected vessels. Vessels that do not have VTR requirements may be underestimated in this analysis.

Table 4: American lobster landing pounds in Massachusetts counties by year from 2017-2021

	2017	2018	2019	2020	2021
Barnstable	3,352,363	2,968,793	2,671,530	2,435,627	2,149,544
Bristol	1,993,809	2,105,482	2,263,234	2,425,182	2,218,481
Dukes	90,776.50	80,010.70	75,597.90	80,167.30	93,559.80
Essex	6,659,528	8,165,759	7,640,883	7,055,319	8,742,081
Middlesex			340.9		
Nantucket	2,887.25	5,573.50	2,583.90	2,856.90	9,937
Norfolk	426,431	425,454	398,522	320,552	265,730
Plymouth	3,331,984	3,271,027	3,346,593	2,890,407	2,961,195
Suffolk	635,346	675,144	630,176	501,741	386,176
Total	16,493,125	17,697,243	17,029,462	15,711,853	16,826,704

Data source: ACCSP dealer report 2017-2021

Jonah Crab

Jonah crab (*Cancer borealis*) is distributed in the waters of the Northwest Atlantic Ocean primarily from Newfoundland, Canada to Florida. The life cycle of Jonah crab is poorly described and what is known is largely compiled from a patchwork of studies. Female crabs are believed to move nearshore during the late spring and summer and then return offshore in the fall and winter.

Jonah crab is managed under the Interstate Fishery Management Plan (FMP) for Jonah Crab (ASMFC 2015) and its three addenda. The FMP lays out specific management measures in the commercial fishery, including a 4.75 inch (12.07 cm) minimum size with zero tolerance, a prohibition on the retention of egg-bearing females, and requiring harvesters to have a lobster permit. Addendum I (May 2016) establishes a bycatch limit of 1,000 crabs per trip for non-trap gear (e.g., otter trawls, gillnets) and non-lobster trap gear (e.g., fish, crab, and whelk pots). Addendum II (February 2017) establishes a coast-wide standard for claw harvest to respond to concerns regarding the equity of the claw provision established in the FMP. Specifically, the Addendum allows Jonah crab fishermen to detach and harvest claws at sea, with a required minimum claw length of 2.75 inches (6.99 cm) if the volume of claws landed is greater than five gallons. Addendum III (February 2018) addresses concerns regarding deficits in existing lobster and Jonah crab reporting requirements by expanding the mandatory harvester reporting data elements, improving the spatial resolution of harvester data, establishing a 5-year timeline for implementation of 100 percent harvester reporting, and prioritizing the development of electronic harvester reporting.

Jonah crabs are primarily caught in pots and traps and have long been taken as incidental catch in the lobster fishery, or more recently as a secondary target, in the lobster fishery. In Massachusetts, most Jonah crabs are landed in Barnstable and Bristol counties. Table 5 displays the yearly landing pounds of Jonah crab by county from 2017 to 2021.

Table 5: *Jonah crab landing pounds in Massachusetts counties by year from 2017-2021*

	2017	2018	2019	2020	2021
Barnstable	2,084,672	1,642,723	1,606,744	1,645,983	1,506,681
Bristol	9,416,030	11,370,068	7,507,008	6,600,901	3,821,477
Dukes	3,757.10	668.47	333.3	390.6	308.66
Essex	171,207	223,045	513,091	304,979	293,024
Nantucket	1,410	151			
Norfolk	721	6,009.28	2,067.71	4,168	
Plymouth	20,545.40	7,841.35	44,358.90	20,352.60	78,382
Suffolk			505	18	
Total	11,698,342	13,250,506	9,674,107	8,576,792	5,699,873

Data source: ACCSP dealer report 2017-2021.

5.3.2 Affected Human Communities

When considering the effect of this emergency rule on human communities, one approach is to focus the analysis on the affected vessels' individual ports or municipalities. However, human communities can extend beyond the boundaries of a particular port or city. Fish can be landed in one town and processed in a neighboring town. Likewise, a fisherman can land catch in one town, live in a neighboring town, and register his vessel in yet another location. In recognition of these factors, this analysis focuses on the county level.⁴ While a county's political boundaries do not limit the network of social interactions and economic resource flows, the use of counties as an analytic focus offers two advantages. First, the geographic range of the county includes individual towns/ports as well as the areas in between with which they likely interact. In addition, many of the data used to characterize communities (e.g., unemployment rate, population) are readily available at the county level. This analysis focuses on four counties in Massachusetts adjacent to the proposed restricted areas: Essex County, Norfolk County, Suffolk County, and Plymouth County.

In both fishing and non-fishing communities, the ability to adapt to change varies with social, political, and economic considerations. The vulnerability of fishing communities, however, is influenced by additional factors, including the importance of familial relationships, the vulnerability of infrastructure, and the commitment to fishing as a culture and way of life (Clay

⁴ This discussion uses the terms "counties" and "communities" interchangeably.

and Olson 2008). From an analytic perspective, vulnerability includes the characteristics of “exposure, sensitivity, and capacity of response to change or perturbation” ((Gallopín 2006) cited in Colburn and Jepson 2012). Consistent with Gallopín’s definition, this social impact assessment considers each county’s vulnerability to be a function of the extent to which its fishing industry is affected by the regulations (i.e., exposure), the significance of the fishing industry within the county (i.e., sensitivity), and baseline factors that may affect each community's ability to absorb the economic costs imposed by the regulations (i.e., capacity to respond to change). The discussion that follows briefly describes the parameters used to evaluate each aspect of vulnerability.

Exposure - The analysis first considers the extent to which the local fishing industry is exposed to the new regulations. Exposure is defined in two ways:

- **Value/proportion of harvest associated with affected gear** – The counties most likely to experience adverse social impacts are those close to the restricted area, and in which the lobster and Jonah crab trap/pot fishery is an important source of commercial fishing revenue, either on an absolute or a relative basis.
- **Number of entities affected** – Similarly, the most vulnerable counties are likely to be those that are home to the greatest number of vessels that fish with lobster trap/pots in the closed area.

Sensitivity - Those communities that are more heavily dependent (both economically and socially) on the fishing industry are more likely to experience adverse social impacts due to fishing regulations. This analysis relies upon a measure of fishing dependence designed to take additional factors into account. This measure, the Occupational Alternative Ratio Summary, emphasizes the importance of fishing as an occupation to participants in the labor force as a whole, and the dependence of the local economy on the fishing industry. In general, a higher score indicates a greater dependence on fishing as an occupation, and a lower likelihood that displaced fishermen can easily enter into alternate occupations.⁵

Capacity to Respond to Change - A number of economic and demographic factors will influence a community’s ability to absorb economic stress, tempering or exacerbating vulnerability to social impacts stemming from regulations:

- **Unemployment Rate, Poverty Rate, Median Income** – Fundamental economic indicators such as the unemployment rate, poverty rate, and median income can indicate the local economy’s resilience to regulatory impacts. Communities that are already economically depressed may find it more difficult to absorb the economic effects of regulatory changes and may be subject to greater social impacts.

⁵ Measures of fishing dependence and gentrification (see below) are based on Hall-Arber et al. (2001). At the time the analysis was developed, these data represented the most recent published attempt to address these issues systematically, allowing for a direct comparison between counties. Colburn and Jepsen (2012) have developed additional indices allowing for evaluation of fishing dependence and gentrification; however, they have yet to be broadly applied. For a qualitative discussion of these issues, see the Community Profiles for Northeast U.S. Marine Fisheries developed by the NMFS Northeast Fisheries Science Center (2010). These profiles are available online at: <http://www.nefsc.noaa.gov/read/socialsci/communityProfiles.html>

- **Gentrification** – Gentrification can be a key source of coastal community vulnerability ((Jacob et al. 2010, Clay and Olson 2008) cited in Colburn and Jepson 2012). According to Hall-Arber et al. (2001), as former working waterfronts succumb to the pressures of gentrification, community character and culture are lost, diversity diminishes, and the fishing community is less able to adapt to changes in the environment. Additional fishing regulations can make it even more difficult for individuals to maintain a “fishing way of life.” Communities that are already experiencing gentrification will likely be more susceptible to social impacts as new regulations are implemented. Hall-Arber et al. (2001) integrate various measures of gentrification into a score that can be used to characterize community vulnerability.

The major ports in the affected area that land lobsters and Jonah crabs include Rockport, Gloucester, Boston, Cohasset, Scituate, and Plymouth. Complete community profiles for these ports can be found in Appendix 3.1. As described in the community profiles, except for Boston, which lands mostly groundfish, and Gloucester, which lands a significant amount of both groundfish and lobsters, all other ports land lobsters as their primary seafood harvest. Table 6 shows the social-economic indicators of each affected community. Essex and Plymouth County have more traditional fishing ports, and their commercial reliance scores are higher than Suffolk and Norfolk County. Norfolk County has the highest income level and lowest unemployment rate. Its low commercial engagement rate indicates that fishermen might have more alternative occupations when fishing is not available. The only major port in Suffolk County is Boston Harbor. It lands a small amount of lobsters and Jonah crabs from a very limited number of vessels.

Table 6: Social-economic indicators for coastal communities

State	County	Key Ports	Population (2018)	Median Household Income (2014-2018)	Persons below Poverty Level (2014-2018)	Unemployment Rate (2018)	Population Composition	Personal Disruption	Housing Disruption	Urban Sprawl	Commercial Engagement	Commercial Reliance
MA	Essex	Gloucester, Rockport, Marblehead	790,638	75,878	10.7%	3.6%	1.24	1.21	1.55	2.79	1.42	1.06
MA	Suffolk	Boston Harbor	807,252	64,582	17.5%	4.5%	3.33	2.33	2.67	4	2	1
MA	Norfolk	Cohasset	705,388	99,511	6.5%	3.0%	1.16	1.08	1.68	2.84	1.04	1
MA	Plymouth	Plymouth, Scituate, Hingham	518,132	85,654	6.2%	3.2%	1.11	1.11	2.25	2.46	1.5	1.04

Source: NMFS social indicator data from 2016.

U.S. Census Bureau 2018: ACS 1-year estimates data profiles; FRED

<https://fred.stlouisfed.org/series/MADUKE7URN>

Notes: Social indicator data are categorical, ranging from 0 to 4. Higher numbers indicate communities that are more vulnerable.

6 IMPACTS OF THE MANAGEMENT ALTERNATIVES

6.1 Impact Designation Descriptions

Using the criteria outlined below and summarized in Table 7, this Environmental Assessment (EA) analyzes the expected impacts of the proposed alternatives for the valued ecosystem components (VECs): protected species, habitat, and human communities as defined in Section 5. For each alternative, impacts to each VEC will be evaluated against the current condition of the VEC (i.e. resource described in the affected environment), as well as relative to the other alternatives proposed. Impacts are described both in terms of their direction (negative, positive, or no impact) and their magnitude (slight, moderate, or high) based on the guidelines shown in Table 7 and Figure 8.

Table 7: Impact determinations for Valued Ecosystem Components

A key of the direction and magnitude of the actions being assessed in the effects analysis. ESA = Endangered Species Act. MMPA = Marine Mammal Protection Act. PBR = potential biological removal level. The Zero Mortality Rate Goal is the requirement for commercial fisheries to reduce incidental mortality and serious injury of marine mammals to insignificant levels approaching a zero mortality and serious injury rate.

General Definitions

VEC	Resource Condition	Direction of Impact		
		Positive (+)	Negative (-)	No Impact (0)
<i>Protected Species</i>	For ESA listed species: populations at risk of extinction (endangered) or endangerment (threatened). For MMPA protected species: stock health may vary but populations remain impacted	For ESA listed species: alternatives that contain specific measures to ensure no interactions with protected species (i.e., no take). For MMPA protected species: alternatives that will maintain takes below PBR and approaching the Zero Mortality Rate Goal	For ESA listed species: alternatives that result in interactions/take of listed resources, including actions that reduce interactions. For MMPA protected species: alternatives that result in interactions with/take of marine mammals that could result in takes above PBR	For ESA listed species: alternatives that do not impact ESA listed species, For MMPA protected species: alternatives that do not impact marine mammals
<i>Habitat</i>	Many habitats degraded from historical effort	Alternatives that improve the quality or quantity of habitat	Alternatives that degrade the quality, quantity or increase disturbance of habitat	Alternatives that do not impact habitat quality
<i>Human Community (Socio-economic)</i>	Highly variable but generally stable in recent years	Alternatives that increase revenue and social well-being of fishermen and/or communities	Alternatives that decrease revenue and social well-being of fishermen and/or communities	Alternatives that do not impact revenue and social well-being of fishermen and/or communities
Magnitude of Impact				
<i>A range of impact qualifiers is used to indicate any existing uncertainty</i>	Negligible	To such a small degree to be indistinguishable from no impact		
	Slight	To a lesser degree / minor	e.g. Slight Negative or Slight Positive	
	Moderate	To an average degree (i.e., more than “slight,” but not “high”)	e.g. Moderate Negative or Moderate Positive	
	High	To a substantial degree (not significant unless stated)	e.g. High Negative or High Positive	
	Significant	Affecting the resource condition to a great degree, see 40 CFR 1508.27.		
	Likely	Some degree of uncertainty associated with the impact		




Figure 8: Depiction of the relative direction and magnitude of impacts on valued ecosystem components

6.1.1 Protected Species

The impacts of the alternatives on protected species take into account impacts to ESA-listed species, as well as impacts to non-ESA listed MMPA protected species in good condition (i.e., marine mammal stocks whose potential biological removal (PBR) level have not been exceeded) or poor condition (i.e., marine mammal stocks that have exceeded or are near exceeding their PBR). These impact descriptors apply to the Protected Species VEC.

6.1.1.1 ESA-Listed Species

For ESA-listed species, any action that results in an interaction or take is expected to have negative impacts, including actions that reduce but do not prevent interactions. Actions expected to result in positive impacts on ESA-listed species include only those that contain specific measures to ensure no interactions (i.e., no take). None of the alternatives considered in this document would ensure no interactions with ESA-listed species. By definition, all ESA-listed species are in poor condition and any take can negatively impact their recovery.

6.1.1.2 MMPA Protected Species

The stock conditions for marine mammals not listed under the ESA varies by species; however, all are legally protected under the MMPA. For non-ESA listed marine mammal stocks, negative impacts would be expected from alternatives that result in the potential for interactions between fisheries and those stocks. For species with PBR that have not been exceeded, alternatives not expected to increase fishing behavior or effort may positively benefit the species by maintaining takes below the PBR and approaching the Zero Mortality Rate Goal, which is the MMPA requirement for commercial fisheries to reduce incidental mortality and serious injury of marine mammals to insignificant levels approaching a zero mortality and serious injury rate. However, none of the alternatives considered in this document ensure no interactions with MMPA protected species, and therefore would be expected to have negative impacts.

6.1.2 Habitat

Alternatives that improve the quality or quantity of habitat are expected to have positive impacts on habitat. Alternatives that degrade the quality or quantity, increase disturbance of habitat, or allow for continued fishing effort are expected to have negative impacts. A reduction in fishing effort is likely to decrease the time that fishing gear is in the water, thus reducing the potential for interactions between fishing gear and habitat.

6.1.3 Human Community

Socioeconomic impacts are considered in relation to potential changes in landings, prices, revenues, and fishing opportunities. Alternatives which could lead to increased availability of target species and/or an increase in catch per unit effort could lead to increased landings. Increased landings are generally considered to have positive socioeconomic impacts because they could result in increased revenues; however, if an increase in landings leads to a decrease in price or a decrease in future availability for any of the landed species, then negative socioeconomic impacts could also occur. Conservation measures that drastically reduce catch and revenue may have negative impacts in the short term, but could ensure access to the fishery in the future, potentially with fewer restrictions.

On the other hand, similar conservation measures could have different impacts on communities depending on their vulnerability and resilience. Communities with lower income and higher fishery dependency, like ports in Essex County, would be more sensitive to stricter restrictions. These communities have less business diversity, and so the communities are less resilient than those with more diverse options, like the Boston-area ports.

6.2 Evaluating the Protected Species Impacts of the Alternatives

6.2.1 Overview of the Decision Support Tool Analysis

The Large Whale Decision Support Tool (DST) is a peer reviewed model that was developed by the Northeast Fisheries Science Center to help managers, decision makers, and stakeholders understand both the spatiotemporal overlap between fishing gear and North Atlantic right whale (right whale) distributions along the East Coast and how risk of entanglement may change after implementing new management measures. This model calculates right whale entanglement risk based on three components: the density of buoy lines in the water, the distribution of right whales, and the threat that gear poses to serious injury as a function of rope strength. The following analyses use DST Version 4.1.0, an updated version of the model used in the 2021 FEIS (NMFS 2021).

The first layer of the model describes the line density per unit area independent of the characteristics of individual lines (e.g. rope strength). The line density component of the DST for the geographic area relevant to this action is based on fishery inputs developed from state and federal trip reports and vessel permits. Permits were used to build state and federal fishery inputs separately, dividing these further by species subgroups. Depending on the level of detail available in trip reports, fishery inputs were built by allocating gear to reported spatial areas, across depth gradients within spatial areas, or distributing gear around spatial coordinates. For each reporting vessel, gear configurations associated with each trip report were used to determine the summed total number of endlines by month and area. This layer was constructed using data collected from 2015 through 2018 for lobster, 2010 through 2020 for other federal trap/pot fisheries, and 2012 through 2019 for other trap/pot fisheries in state waters, all prior to the

implementation of the 2021 Final Rule (86 FR 51970) and the Massachusetts state waters' commercial trap/pot gear closure (322 CMR 12.04(2)).

A second layer in the DST model assesses the risk associated with different gear configurations, accounting for the various breaking strengths of individual lines. Gear with higher breaking strength is expected to be more risky to whales because it is harder to break out of and therefore more likely to result in serious injury or mortality. An empirical gear threat model was built using information on the strength of ropes involved in serious whale entanglements and how the strength of the ropes observed in entanglements compares to the strength of ropes that whales would be expected to encounter. The model estimates uncertainty within the gear threat model and can provide an upper and lower bound within the model output. Models for the upper and lower confidence bounds were calculated by bootstrapping the observed line strength data to generate a ratio of observed to expected line strengths and fitting the data to a binomial generalized linear model. See Appendix 3.1 of the 2021 FEIS (NMFS 2021) for more detail about the development of the gear threat model and associated uncertainty.

The final layer is a right whale habitat density model. The DST employs a right whale habitat density model built by researchers at Duke University's Marine Geospatial Ecology Laboratory in the Nicholas School of the Environment (hereafter referred to as the Duke habitat model or whale habitat density layer) that predicts the spatiotemporal distribution and density of right whales throughout the proposed area based on observations of whales from standardized surveys from January 2010 through September 2020 and co-located oceanographic and habitat variables (Version 12; Roberts et al. 2016; Roberts et al. 2020). The right whale habitat density model then uses oceanographic and habitat variables across the region to create a map of likely whale presence. The DST also includes a humpback whale habitat density model (2009 through 2019; Roberts et al. 2016) and a fin whale habitat density model (1998 through 2020; Roberts et al. 2017) that were used to determine predicted density of these two additional species Alternative 2 and Alternative 3 closure areas. The whale habitat model does not include more recent whale distribution data, such as aerial and shipboard survey data collected during 2021, 2022, or 2023 nor does it include any acoustic detection and opportunistic sightings. However, the Duke habitat model does undergo a rigorous, peer-reviewed validation process in which empirical data, such as passive acoustic monitoring data, are compared to the final predicted whale density layers (for more information on the whale density model data use and validation process see Roberts et al. 2016; Roberts et al. 2020, Roberts 2022).

The DST analyzes information on a common spatial grid with consistent positioning and resolution (i.e., cell size). It employs two spatial resolutions for analysis: a low resolution (10 square nautical miles/16 square kilometers) and a high resolution (1 square nautical mile/1.6 square kilometers) option. The analysis in this EA was conducted using the high resolution setting on a one nautical mile (1.6 square kilometer) grid. For detailed information on the model, including its limitations and uncertainties, refer to Volume I, Chapter 3 of the 2021 FEIS and the model documentation in Volume II, Appendix 3.1 (NMFS 2021).

Each model run allows for the selection of a variety of spatially explicit management measures for a particular month with a focus on measures that reduce the number or strength of lines in the water column, such as changes in the number of traps per trawl, the proportion of traps fished,

line strength, line number, restricted areas with lines out and/or lines moved to adjacent fishing areas, and number of lines per trawl. The output provides the mean reduction in risk throughout an entire year. Suites of measures can be run in tandem to best estimate overall changes in risk while taking into account how different management measures may interact with one another to alter the risk landscape.

We used relative risk reduction to estimate the risk of a serious entanglement, which takes into account the overlap between whale density, line density, and the strength of the line. Together, these components roughly estimate the approximate risk of an entanglement that will result in mortality or serious injury, where a higher density of lines, higher predicted density of right whales, and/or high line strength increase the eventual estimate of relative risk. This enables a semi-quantitative comparison of how different management scenarios and gear modifications are predicted to change the risk of entanglements that result in mortality or serious injury.

Relative risk was calculated by spatially constraining the DST model in two ways. First, the model was constrained to the Massachusetts portion of Lobster Management Area 1 (MA LMA 1; the action area). MA LMA 1 limits the scope of the analysis to the action area and calculates entanglement risk reduction relative to the total risk landscape within the spatial scope of this emergency action (that is, risk reduction relative to all risk in MA LMA 1). The second round of analyses expanded the spatial constraint to the trap/pot fishery in the Northeast Region Trap/Pot Management Area (Northeast Region), which makes up approximately 94 percent of fixed gear buoy lines in the right whale range within U.S. waters (risk reduction of the action relative to all of the Northeast Management Area evaluated in the 2021 FEIS (NMFS 2021)). Increasing the spatial constraint to the Northeast Region provides a much broader analysis of entanglement risk reduction provided by the proposed alternatives and allows a comparison of the emergency rule with the measures implemented on September 17, 2021 that went into effect May 1, 2022.

It is important to model fishing effort across the region to understand how this emergency action may redistribute effort, and consequently entanglement risk, across the Northeast Region. Within the MA LMA 1 and Northeast Region spatial constraints, restricted areas were analyzed two ways, taking into account gear location and whether gear would be removed or relocated. We know from existing closures that removal of all gear is more likely for nearshore restricted areas, particularly the Massachusetts Restricted Area (MRA), when fishermen would have a long transit to open areas and where those without federal permits are restricted in area choices. However, some fishermen with federal permits would be able to move their lines and could increase risk outside of restricted areas, as may have occurred in 2021. Discussions with Massachusetts fishermen in 2022 suggested that, due to good lobster prices in 2021 and again in the spring of 2022, relocating gear outside of the closure is attractive to fishermen, if possible (Robert Martin, pers. comm. 2022). A similar mixed response is expected in 2023, with fishermen from more northern ports relocating gear to open federal waters, and fishermen from the more southern ports with less access to federal waters removing their gear. This analysis considered both extremes (gear reduction under which all lines were removed and a closure under which all lines were relocated outside of the emergency closure) to estimate the maximum and minimum level of risk reduction anticipated based on fishing behavior. The effects of these two responses differ slightly depending on how they correspond to overlap between right whales and trap/pot gear. When fishing is suspended or ropeless technologies are employed and lines are

removed from the water entirely, there is typically a straightforward decrease in risk of entanglement. For optimal conservation, the emergency restricted area needs to be sufficiently large to provide protection for whales, but not designed such that fishermen would relocate large numbers of lines to other areas of high risk and/or create a fencing effect along the borders of the restricted area.

In this EA, the DST is used in a similar manner as in the 2021 Final Rule of the modifications to the Plan to assess the possible risk reduction of the two action alternatives on large whale species. The analyses evaluated risk reduction that occur on top of the baseline measures implemented with the 2021 Final Rule within the two spatial constraints (MA LMA 1 and the Northeast Region) achieved by the two anticipated fishery behavior options in which fishermen either remove gear from the water (Gear Reduction) or relocate gear outside of the emergency restricted area (Closure). Closure scenarios assume that the number of buoy lines within the water is the same before and after a management intervention. The analysis of the status quo assumes compliance with these current requirements. For a detailed description of model settings and results see Appendix 4.1

6.2.2 The Impacts of Alternative 1 on Protected Species

Under Alternative 1 (No Action), the current Plan management regime consisting of time/area closures, minimum trap per trawl requirements, use of weak buoy line inserts or buoy line, and gear marking requirements remain in place. The closures included in the Preferred Alternative of the 2021 Final Rule are considered part of the status quo for this proposed action (see Subsection 4.1) because they were implemented on September 17, 2021 and gear modifications went into effect on May 1, 2022. Under No Action, high negative impacts are expected because there would be a risk of entanglement due to the present number of buoy lines that would remain in the water when right whales are abundant in the area.

In MA LMA 1, right whale habitat density increases each month as time passes February through April (Table 8). Right whale density peaks in April, reflecting the importance of that month in the action area. The high density aggregation of right whales in Cape Cod Bay motivated the implementation of the large seasonal MRA that was designed to separate right whales from fishing gear. On average, trawls are longer and lines are stronger in federal waters compared to state waters within the action area, so the gear present in the alternative closure areas poses a higher risk compared to state waters.

Table 8: *Estimated total number of Atlantic large whales in Action Area*

Comparison of estimated total number of whales by species within Alternative 2 (Preferred) and Alternative 3 emergency closures areas by month (February, March, and April). The estimated total number of whales are the estimates of whales present during each month at any given time as projected by the whale habitat density models created by Jason Roberts and Duke University (Roberts et al. 2016, Roberts et al. 2017, Roberts et al. 2020).

Area	Right Whale			Humpback Whale			Fin Whale		
	Feb	Mar	Apr	Feb	Mar	Apr	Feb	Mar	Apr
MRA Wedge (Alternative 2: Preferred)	0.04	1.4	3.3	0.2	0.2	0.9	0.4	0.3	0.8
MRA Wedge North (Alternative 3)	6.7	2.4	4.6	2	2.4	17.6	8.3	7.1	7.6

The DST model uses whale survey data from January 2010 through September 2020 to predict right whale distribution, 2009 through 2019 for humpback whale distribution, and 1998 through 2020 for fin whale distribution. Table 8 shows the expected whale densities for right, humpback, and fin whales by month within the closed areas of Alternative 2 and Alternative 3. As described in Subsection 6.2.1, the Duke habitat model does not include more recent whale distribution data, such as the empirical data collected during aerial surveys conducted by the Center for Coastal Studies (CCS), nor does it include aerial and shipboard surveys conducted by Northeast Fisheries Science Center (NEFSC), acoustic detections, or opportunistic sightings, and therefore may be underestimating right whale density in some areas. The CCS observations include right whales within the affected wedge and observations of aggregated fishing gear (Figure 9). The fishing gear is thought to be a mix of actively fished gear and, in April, staged gear that is set in preparation for the opening of the MRA (Figure 10). Federal waters within the MRA are open to the trap/pot fishery on May 1 and Massachusetts State regulations prohibit commercial trap/pot fishing in any waters under the jurisdiction of the Commonwealth February 1 through May 15 (322 CMR 12.04(2); Figure 1).

Without taking emergency action, the aggregation of gear remains an imminent threat to the right whale population as well as to other large whales in the area. No Action continues the status quo of Phase I implementation of seasonal closures and presents a high to moderate negative risk to ESA-listed (right, fin, and sei whales) and MMPA protected species (humpback and minke whales) within the affected area. Relative to Alternative 2, No Action has a slight negative impact because No Action would allow fishing activities with buoy lines that present an entanglement risk to protected species in a relatively small area (200 square miles/518 square kilometers) of federal waters outside of the MRA. Relative to Alternative 3, No Action will have a moderate negative impact because Alternative 3 would close a substantial area (1,297 square miles/3,359 square kilometers) to the lobster and Jonah crab, and other trap/pot fisheries that under No Action would remain open. As has been noted, buoy lines used by the trap/pot fisheries present an entanglement risk. Under No Action, no additional risk reduction to prevent mortality and serious injury would occur, meaning that ESA and MMPA protected species would experience high negative to moderate negative impacts.

Figure 9: North Atlantic Right Whale sightings and fishing gear observed in 2021 and 2022 within Proposed Emergency Closure, MRA Wedge

North Atlantic right whale sightings collected February, March, and April 2018-2022 (blue and red crosses) and gear observed (black stars) on four different days are overlaid. Blue crosses show right whale sightings collected through dedicated aerial surveys conducted by the Center for Coastal Studies (CCS) and Northeast Fisheries Science Center (NEFSC) and dedicated shipboard surveys conducted by CCS and Stellwagen Bay National Marine Sanctuary. Red crosses are opportunistic sightings collected by NEFSC. Black stars represent fishing gear observed by CCS on April 19 and 28, 2021 and February 6 and March 11, 2022, four days that were selected as representative snapshots of fishing gear present in survey areas. Surveys concentrate on Cape Cod Bay; surveyors rarely fly north of mid Cape Ann, off Rockport, MA. Massachusetts Restricted Area is shaded in dark gray, and MRA Wedge (Alternative 2) is represented by the light gray area. These maps are used for qualitative not quantitative comparison, and differ from Large Whale Decision Support Tool data.





Figure 10: Fishing gear observed within Massachusetts Bay in spring 2021 and 2022 within Proposed Emergency Closure, MRA Wedge

Fishing gear observed by the Center for Coastal Studies (CCS) on April 19, 2021 (white circles), April 28, 2021 (encircled “x”), February 6, 2022 (open diamonds), and March 11, 2022 (black diamonds), were selected as representative snapshots of fishing gear present in survey areas. Surveys concentrate on Cape Cod Bay; surveyors rarely fly north of mid Cape Ann, offshore Rockport, MA. Massachusetts Restricted Area is shaded in dark gray, and the MRA Wedge (Alternative 2) is represented in pale gray. These maps are used for qualitative not quantitative comparison, and differ from Large Whale Decision Support Tool data.



6.2.3 *The Impacts of Alternative 2 on Protected Species*

Alternative 2 would implement an emergency closure prohibiting trap/pot fishery buoy lines from February 1 through April 30 within the 200 square miles (518 square kilometers) of federal waters referred to as MRA Wedge. The boundary of MRA Wedge lies between state and federal waters within the MRA and coincides with the existing MRA closure season (Figure 1).

The maximum and minimum relative risk reduction for Alternative 2 was estimated based on two assumptions of what happens to gear during a closure, as discussed in Subsection 4.2.1. The maximum relative risk reduction relies on the assumption that all lines are removed from the water, whereas the minimum risk reduction estimate assumes a closure redistributes the gear to areas outside of the emergency closure. The actual risk reduction will likely fall between the two analyzed extremes. The greater the reduction in overlap between right whales and buoy lines, such as when all lines are removed, the smaller the likelihood of a right whale dying or becoming seriously injured in buoy lines. Removing lines provides greater benefit to right whales present during February, March, and April than if the lines are moved elsewhere within the range of the right whales.

Within MA LMA 1, the MRA Wedge emergency closure February 1, 2023 through April 30, 2023 reduces entanglement risk posed by trap/pot gear by 13.2 to 16.6 percent, depending on whether gear is relocated outside of the emergency closure or removed (Table 9). Relative to the entire Northeast Region, trap/pot gear in the MRA Wedge this is an approximate 1.9 to 2.4 percent reduction of trap/pot entanglement risk (Table 9). These estimates suggest that the difference in risk reduction between gear removal and relocation may be fairly small, but there are limitations in the ability of the model to predict where gear is reset and in what density. Gear that is relocated in particular areas in high numbers could pose more of a risk than the model results reflect. February shows the possibility of a slight increase in risk if all gear was moved outside of the closure but, as indicated in Subsection 6.2.1, we anticipate a mix of responses with a good portion of the gear being removed from the water instead of relocated into other areas in high density. The proposed area is a particular concern for storage of gear in April as the May 1 opening of the federal MRA, which is expected to have the highest whale density. In the absence of an emergency closure, fishermen may relocate gear along the boundaries of these waters, staging for a rapid relocation after May 1, which is similarly not captured in the DST analyses and may be underestimated. Since right whales are expected to disperse during this period, the potential for a gear fence is of some concern, but this risk is outweighed by the high risk during February through April across the MRA and emergency closure area and potential for gear storage to concentrate within the Proposed Area. Whale density outside along the border of MRA Wedge is expected to be lower than within the emergency closure area, particularly relative to MA LMA 1, which is one of the most valuable regions for the species.

Table 9: Comparison of North Atlantic right whale entanglement risk reduction by month

Comparison of risk reductions in North Atlantic right whale entanglement risk by month within proposed emergency closure within Alternative 2 (Preferred) and Alternative 3. Closure scenarios assume that the gear is relocated outside of the emergency restricted area, leaving the number of buoy lines within the water the same before and after a management intervention. Gear Removed scenarios that assume gear is removed from the waters. The MA portion of LMA 1 refers to the Massachusetts portion of Lobster Management Area 1. The Northeast Region refers to the lobster and Jonah crab trap/pot fishery in the Northeast Region Trap/Pot Management Area.

Risk reduction relative to MA portion of LMA 1	Closures				Gear Removed			
Area	Feb	Mar	Apr	Feb 1-Apr 30	Feb	Mar	Apr	Feb 1-Apr 30
MRA Wedge (Alternative 2: Preferred)	-1.9%	3.6%	11.5%	13.2%	0.1%	4.1%	12.4%	16.6%
MRA Wedge North (Alternative 3)	9.3%	2.8%	10.2%	22.3%	14.9%	6.4%	16.1%	37.4%
Risk reduction relative to All Northeast Trap/Pot	Closures				Gear Removed			
MRA Wedge (Alternative 2: Preferred)	-0.3%	0.5%	1.7%	1.9%	0.0%	0.6%	1.8%	2.4%
MRA Wedge North (Alternative 3)	1.4%	0.4%	1.5%	3.3%	2.2%	0.9%	2.4%	5.5%


Relative to the recent empirical data, it is likely that the right whale habitat density model underestimates the number of whales in this area in recent years (see Figure 9 for February, March and April 2018-2022 right whale sightings data provided by CCS and NEFSC). As discussed in Subsection 6.2.1, the DST right whale habitat density model relies on oceanographic and habitat variables to create a map of likely whale presence using whale data from January 2010 through September 2020. Therefore, not all sources of data are included within the model, including more recent survey data or acoustic and opportunistic detections of right whales. Dedicated right whale survey efforts conducted by CCS and NEFSC are centralized in Cape Cod Bay, and surveys northward in Massachusetts Bay are not conducted with the same frequency. Therefore, there are limited data on whale density in this area, and shifting right whale distribution within the last five years makes predicting whale aggregations and distribution challenging. However, the sightings data of right whale distribution suggest a higher concentration of right whales than is estimated by the DST. Multiple aggregations of two to three right whales were observed by NEFSC dedicated aerial and shipboard surveys in the middle and western portion of the MRA Wedge area around 42°20' (Figure 9). The opportunistic sightings collected by NEFSC on whale presence along the southern border of the MRA Wedge area, range from individual whales to groups as large as ten. However, the DST estimates only up to three right whales in the same area at any given point February through April (Table 8). Therefore the risk estimated by the DST likely underestimates the risk reduction of the proposed emergency closure.

Acoustic detections of right whales also indicated that whale presence may be higher or more persistent than what is estimated by the whale habitat model used in the DST. Acoustic data were collected by Slocum Glider surveys deployed near the Stellwagen Bank National Marine

Sanctuary and Gulf of Maine for the purpose of surveying tagged fish and baleen whales, including right, fin, sei, and humpback whales (Baumgartner 2021; Figure 11). Acoustic data do not confirm the number of individual whales present nor do these particular data provide exact locations of the animals; however, the data do confirm sustained large whale presence in the action area during March and April 2021 the proposed seasonal closures of Alternative 2 (Figure 11, Figure 13).

Figure 11: Acoustic detections of fin, sei, and right whales in Massachusetts Bay from March 17, 2021 through April 30, 2021

Definite and possible acoustic detections of fin, sei, and right whales in Massachusetts Bay from March 17, 2021 through April 30, 2021. The acoustic detection data comes from Slocum Glider surveys deployed near the Stellwagen Bank National Marine Sanctuary and Gulf of Maine (Baumgartner 2021). Restricted areas proposed in Alternative 2 and Alternative 3 are represented in the simple hatched areas. Massachusetts Restricted Area is outlined in black. These maps are used for qualitative not quantitative comparison and differ from Large Whale Decision Support Tool data.



Data Credit: Julianne Wilder and Genevieve Davis, acoustic analysts; Mark Baumgartner, glider operator; mission funding from SanctSound and NERACOOS.

Empirical survey evidence indicates that dense aggregations of gear were present within the action area in 2021 and 2022, along with right whales in higher numbers than projected by the whale habitat model used in the DST. Systematic surveys for whales are done differently than surveys for gear, and gear surveys are not typically conducted so data on gear density within the action area are not available for comparison across seasons or years. Changes in fishing effort distribution data would not be reflected by the DST, which models gear abundance and distribution based on fishing data before the MRA Wedge was created. The DST analyses may underestimate risk reduction because the fishing effort data does not reflect the additional risk of potential gear storage in the area. It remains uncertain if the expansion of the MRA in 2021 resulted in an increase in the amount of trap/pot gear in this area in response to the Massachusetts state water closure, in which case, the DST would underestimate the amount of gear in this area and the associated risk. Alternative 2 would reduce overlap of gear and whales

through an emergency closure prohibiting the use of buoy lines by the trap/pot fishery in an area of observed right whale presence.


Introducing ropeless fishing, where buoy lines are stored on the bottom until retrieval, in the action area where right whale aggregations may be high, may pose a slightly higher threat of entanglement in the short-term compared to a full fishery closure. Gangions and sinking groundline would be present in ropeless and on-demand fishing gear, and some on-demand gear includes short term deployment of a buoy line while a vessel is on site. However, there are long-term benefits to the accelerated development of gear that protects right whales. To reduce potential risks in the short term, an Exempted Fishing Permit (EFP) would be required that would include conditions on fishing to reduce the impact of this gear on right whales. No more than 7 vessels fishing 10 trawls each are anticipated to be fishing with ropeless gear in the MRA federal waters under an EFP in 2023, including the buoy line closure area during February, March, and April. These vessels will be fishing without persistent buoy lines and with extensive monitoring and reporting to verify that the effort would add only a negligible risk above a full closure.

Overall, large whale entanglement risk following the implementation of the emergency closure will likely decline substantially during a critical period when the right whales are likely to be exiting Cape Cod Bay. While entanglement risk is not completely eliminated, the action does significantly reduce risk in the action area. With the addition of 200 square miles (518 square kilometers) February, March, and April 2023, there is a decrease in entanglement risk for right whales during those three months. This emergency measure is expected to substantially reduce the potential of a right whale mortality or serious injury as a result of interactions with fishing gear.

Alternative 2 could reduce overlap of minke, humpback, and fin whales with trap/pot buoy lines, but any reduction would be minimal. The MRA Wedge emergency closure is unlikely to impact minke, humpback, or fin whales substantially but any impact is expected to have a positive impact, similar to the conclusion in the 2022 Emergency Rule EA (NMFS 2022). Minke, humpback, and fin whales are present in the Massachusetts portion of LMA 1 and were observed by dedicated surveys and opportunity sightings February, March, and April 2018-2022 (Figure 12). Acoustic detection of fin whales indicates abundance in Massachusetts Bay peaks in September to January, though fin whales may be present throughout the year (Hain et al. 1992, Morano et al. 2012). Passive acoustic monitoring data indicate that humpback whales persistently utilize Massachusetts Bay April through December to feed (Murray et al. 2013, Clapham et al. 1993). February, March, and April 2018-2022 observation data show low numbers of confirmed visual sightings of sei whales within the action area.

Figure 12: Visual sightings of Atlantic large whales in Massachusetts Bay during February, March, and April 2018-2022

Atlantic large whale (minke, fin, humpback, and sei whales) sightings in the Massachusetts portion of LMA 1 reported during Northeast Fisheries Science Center dedicated aerial surveys conducted in February, March, and April 2018-2022. Dedicated surveys are concentrated on Cape Cod Bay, Massachusetts, which may underestimate whale abundance north of Cape Ann. Restricted areas proposed in Alternative 2 and Alternative 3 are represented in the simple hatched areas. Massachusetts Restricted Area is outlined in black. These maps are used for qualitative not quantitative comparison, and differ from Large Whale Decision Support Tool data.




The DST is unable to provide a quantitative estimate of risk reduction for minke or sei whales, both of which have been observed in the action area over spring of 2018 through 2022 in low numbers. Minke whales have been frequently observed in Massachusetts Bay and Cape Cod Bay throughout the year. Feeding behavior was observed most frequently May through October, and the highest abundance of minke whale sightings were documented July through October (Murphy 1995). NEFSC aerial surveys conducted February, March, and April 2018-2022 reported five minke whales within the MRA Wedge and six total minke whales observed in the MRA Wedge North area during that same period (Figure 12). Minke whales were more frequently documented within the MRA and on the continental shelf south of Rhode Island. Sei whale sightings in February, March, and April 2018-2022 were concentrated within federal waters of the MRA and offshore waters beyond LMA 1. Though sei whales have been observed in the Great South Channel and Stellwagen Bank in the spring, they typically remain offshore in the deeper waters (Payne et al. 1990, Schilling et al. 1992, Waring et al. 2009).

Acoustic detections of fin, sei, right, and humpback whales confirm their presence in federal waters east of Massachusetts Bay, Massachusetts (Baumgartner 2021; Figure 11, Figure 13). Humpback and fin whales were detected closer inshore than sei whales, but the acoustic detections do indicate sei whales were also present in the MRA Wedge March and April 2021. Detectability for baleen whales may vary depending on abundance, distance from the glider, whale calling behavior, and environmental conditions affecting interfering noise (Baumgartner et al. 2020). However, the acoustic data does confirm continued presence of these four species of Atlantic large whales in the action area during the time period of the proposed closures of Alternative 2.

Figure 13: *Acoustic detections of humpback whales in Massachusetts Bay from March 17, 2021 through April 30, 2021*

Definite and possible acoustic detections data of humpback whales in Massachusetts Bay from March 17, 2021 through April 30, 2021. The acoustic detection data comes from Slocum Glider surveys deployed near the Stellwagen Bank National Marine Sanctuary and Gulf of Maine (Baumgartner 2021). Restricted areas proposed in Alternative 2 and Alternative 3 are represented in the simple hatched areas. Massachusetts Restricted Area is outlined in black. These maps are used for qualitative not quantitative comparison and acoustic detections data differ from Large Whale Decision Support Tool data.



Data Credit: Julianne Wilder and Genevieve Davis, acoustic analysts; Mark Baumgartner, glider operator; mission funding from SanctSound and NERACOOS.

The evidence available for large whales does not suggest that new entanglement risk hotspots would be created by relocated gear outside of the emergency closure area. Therefore, the implementation of Alternative 2 is unlikely to further negatively impact protected species. Relative to No Action, Alternative 2 would have a negligible to slight positive impact on ESA-listed (right, fin, and sei whales) and MMPA protected species (humpback and minke whales) because large whale entanglement risk in trap/pot gear is reduced. Relative to Alternative 3, Alternative 2 has a negligible to slight negative impact because the proposed emergency closure area of Alternative 2, MRA Wedge, provides less risk reduction than the proposed emergency closure of Alternative 3, MRA Wedge North. Considered alone, ESA-listed and MMPA protected species would be moderately negative to slightly negatively impacted by Alternative 2

because this proposed emergency action does not eliminate the potential for all interaction risk between fishing gear and marine mammals that could result in takes above PBR.

6.2.4 The Impacts of Alternative 3 on Protected Species

Alternative 3 would also implement an emergency closure, similar to Alternative 2, but would close an area more than six times larger than Alternative 2 (1,297 square miles/3,359 square kilometers). The proposed emergency restricted area in Alternative 3, MRA Wedge North, would remain closed to the use of buoy lines by the trap/pot fishery from February 1, 2023 to April 30, 2023. Between February and April, the total number of right whales within the MRA Wedge North at any given time is estimated to be between 3 to 7 (Table 8). This may be an underestimate of whale density south of Cape Ann, Massachusetts according to survey data as discussed in Section 6.2.3. Surveys are less frequent north of Cape Ann; therefore, similar 2020, 2021, 2022 sightings data are not available and it is unknown how the right whale habitat density model aligns with more recent distribution of whales. All age classes have been observed near Jeffreys Ledge located in offshore waters of Massachusetts, and sightings data indicate whales may be traveling from coastal waters of Massachusetts north toward this region (Weinrich et al. 2000) suggesting that right whales may be present north of Cape Ann more often than is being reported by dedicated surveys. Further information is needed to confirm abundance in this area.

Implementing the MRA Wedge North emergency closure reduces mean risk in the MA LMA 1 by 22.3 to 37.4 percent depending on if gear is relocated outside the emergency closure or removed, respectively (Table 9). As discussed in Section 6.2.2, the DST is likely underestimating the risk reduction benefits of an emergency closure to right whales because empirical survey sightings indicate a larger presence of whales than the DST average. The value of this closure is also likely higher because extending the closure farther northeast is more likely to remove lines rather than move them given the distance from Massachusetts ports. Additionally, the potential for dense concentrations of gear along the closed area border is diluted given the larger and offshore border. Removal of gear reduces entanglement risk while also preventing the formation of new hotspots where newly relocated gear and right whales overlap. This is a significant risk reduction relative to this portion of the Northeast trap/pot fishery.

In the entire Northeast Region, mean risk reduction ranges from 3.3 to 5.5 percent depending on whether gear is relocated or removed from this area (Table 9). For reasons discussed in Section 6.2.3, risk reduction may be underestimated but results demonstrate a precautionary benefit to large whales through the removal of buoy lines within the affected area. It is unlikely that gear would be relocated outside of this area given the distance from MA home ports. Only vessels with dual LMA permits for LMAs 1 and 3 would be able to relocate into LMA 3 during the closure. Therefore, we assume lines would be removed from the water.

Similar to Alternative 2, introducing ropeless fishing may slightly increase the entanglement risk in the groundline in the short-term, compared to a full closure, but the amount of gear expected to be fished in this area (no more than 70 trawls) during the emergency closures is very small relative to current effort and sinking ground line requirements will prevent most of the gear from being in the water column and available for entanglement. The presence of buoy lines in areas

where whales are aggregating and transiting will decline during the seasonal closure period. Therefore, this emergency measure would be expected to substantially reduce the potential of a right whale mortality or serious injury as a result of interactions and entanglement with fishing gear.

Removing buoy lines from MRA Wedge North from MA LMA 1 would also decrease the risk of mortality or serious injury for minke, humpback, and fin whales. The DST projects 2 to 18 humpback whales and 7 or 8 fin whales from February through April within the boundaries of the MRA North (Table 8). As discussed in Section 6.2.3, minke, humpback, and fin whales were present in the MA LMA 1 in the spring 2018-2022, and visual observations and acoustic detection data indicate they were also present within the boundaries of MRA Wedge North during the closure period (Figure 11, Figure 12, Figure 13). Minke whales were not detected acoustically, and were observed infrequently through visual surveys. Humpback whales were observed visually in Massachusetts Bay and scattered within MRA Wedge North. However, acoustic detection data strongly indicates that they are present throughout federal waters north of MRA to the New Hampshire border. The fin whale observations suggest they do feed or travel within the boundaries of the emergency closure, and the sightings and acoustic data confirm the importance of multiple survey efforts to monitor stock status, presence, and abundance for Atlantic large whales. More information is needed to fully quantify minke, humpback, and fin whale abundance and habitat use in this area.

The DST is unable to provide a quantitative estimate of risk reduction for minke or sei whales, both of which have been observed in low numbers within the action area (Figure 11, Figure 12). NEFSC aerial surveys in 2021 reported a single minke whale offshore Cape Ann within the MRA Wedge North and the remaining minke whale observations during that time period were documented within Massachusetts Bay. Sei whales were sighted less frequently by NEFSC aerial Surveys conducted February through April 2018-2022, and no sei whales were sighted within the MRA Wedge North. However, the Slocum Glider detected sei whales acoustically in Massachusetts Bay and federal waters (Baumgartner 2021; Figure 11). As discussed in Section 6.2.3, dedicated aerial surveys conducted by NEFSC focus on Cape Cod Bay. The lack of sightings appear to be attributed to insufficient survey effort in the region, as they were detected acoustically in March and April of 2021. The sightings and acoustic data indicate that humpback, fin, minke, and sei whales would not be negatively impacted by the MRA Wedge North closure. However, according to the DST, extending the boundary of MRA to the New Hampshire state water boundary.

Reducing lines through a closure may lower entanglement risk for these species during February 1, 2023 through April 30, 2023. Compared to No Action, Alternative 3 would have a moderate positive impact on protected species because the emergency closure proposed in Alternative 3, MRA Wedge North, would close a large area to the trap/pot fishery which reduces the risk of entanglement in buoys lines for ESA-listed (right, fin, and sei whales) and MMPA protected species (humpback and minke whales). Relative to Alternative 2, Alternative 3 has a slight positive impact on ESA-listed and MMPA protected species. When considered alone, Alternative 3 would have negligible to slight negative impacts on large whales due to the continued operation of the fishery, and potential risk of interaction between the fishery and take of ESA-listed and MMPA-protected species remains.

6.2.5 Comparison and Summary of the Alternatives

The primary difference in biological impacts between the alternatives is the removal of buoy lines within the water column that are directly related to the reduction in right whale entanglement risk. Alternative 2 would remove or relocate fewer lines than Alternative 3, but the Alternative 2 emergency closure achieves more risk reduction per square mile than Alternative 3. Removing the vertical lines from February 1, 2023 to April 30, 2023 would reduce entanglement risk for these large whales when they are more likely to occur in high densities.

The impact on ESA-listed (right, fin, and sei whales) and MMPA protected species (humpback and minke whales) is expected to be slightly positive under Alternative 2 and moderately positive under Alternative 3, when compared to No Action. The MRA Wedge North (Alternative 3) is substantially bigger in area (by a magnitude of six) than MRA Wedge (Alternative 2). Alternative 2 provides less overall risk reduction (1.9 to 2.4 percent, out of all Northeast trap/pot fisheries) compared to Alternative 3 (3.3 to 5.5 percent, out of all Northeast trap/pot fisheries). The difference in impact between the two is even greater when considering local risk in MA LMA 1, an area with particularly high entanglement risk during spring (13.2 to 22.3 percent under Alternative 2 compared to 14.9 to 37.4 percent under Alternative 3).

6.3 Habitat Impacts of the Alternatives

6.3.1 Alternative 1: No Action

The No Action alternative (Alternative 1) would maintain current regulations seasonally closing the MRA to trap/pot gear, while continuing to allow access to trap/pot fisheries in the portion of LMA 1 primarily used by Massachusetts vessels. Although the footprint of each trap on the bottom is minimal, as the gear will be weighted to sit on the ocean floor, some level of disturbance to the habitat is likely, particularly when placed in long trawls. The Plan currently requires a minimum of 10 to 25 traps per trawl depending on distance from shore throughout the action area but average trawl length is closer to 25 traps. Baseline conditions may already contribute to some disturbance on the seafloor when the gear is hauled and set.

No Action represents the status quo and will likely continue to have a moderately negative to slightly negative impact on the habitat. Trap/pot fishing will continue at current levels and disturbance of the habitat will not change. Relative to Alternative 2 and Alternative 3, No Action is expected to result in negligible to slight negative impacts because fishing gear will be concentrated within the MRA Wedge during the emergency closure period.

6.3.2 Alternative 2: Preferred Action

Under Alternative 2, fishing and setting trap/pot gear with persistent buoy lines would be restricted in the MRA Wedge from February 1, 2023 through April 30, 2023. As noted in Subsection 4.2, authorizations for fishing without buoy lines (ropeless fishing) in the MRA Wedge must be obtained through an EFP. At this time, there are no significant efforts within the fishery to obtain EFPs that could be issued during this time period.

Alternative 2 would close the MRA Wedge to trap/pot fishing February 1, 2023 through April 30, 2023. Outside of this, fishing operations will continue to occur. The removal of traps may decrease benthic community disturbance, protect local community structure, and may increase local lobster and/or Jonah crab abundance (Uhrin 2016). However, trap/pot gear may be relocated outside of the emergency closure, and there is uncertainty around what percentage of gear will be removed from the water and what percentage of gear will be relocated elsewhere. It is difficult to predict and quantify impacts to habitat if gear is displaced because there is uncertainty around where the gear will be relocated for active fishing operations or “wet storage” of gear being hauled only once per 30 days (consistent with regulations) set near or along closed area borders for a quick move once seasonal restrictions are lifted on May 1, 2023. There is a potential for trap/pot gear to be redistributed in an area that has not historically been disturbed by fishing. We anticipate that it is difficult for vessels in the southern portion of the restricted area to redistribute their traps outside the northern or eastern boundaries, given the cost of operation and expected landings in April, so we assume some fishing from these ports remove their gear while those farther north will move their gear into other areas.⁶ However, operational trap/pot gear is not considered to cause long-term benthic impacts and lobster fishing is believed to negligibly impact EFH (Uhrin 2016, Goode et al. 2021). At this point, it is unlikely that ropeless fishing could expand widely during the emergency closure, but if it did, the habitat could experience similar levels of disturbance as described under No Action.

Relative to No Action, impacts of Alternative 2 on the habitat are expected to be negligible to slight positive because while fishery operations for the trap/pot fisheries will be prohibited in this area from February 1, 2023 through April 30, 2023, other fisheries will continue to operate in this area. Relative to Alternative 3, impacts of Alternative 2 on the habitat are expected to be negligible. The emergency closure area of Alternative 3 is 6 times larger than the emergency closure area of Alternative 2, meaning that a much smaller area will be temporarily closed to disturbances created by trap/pot fishing operations in Alternative 2. However, given the temporary nature of the proposed emergency closure, these habitat impacts are not considered to be significantly different. The closure that would be implemented under either alternative will only be in place for 3 months, which is not sufficient time to allow the habitat to recover after a previously occurring disturbance. Relocated gear may disturb benthic habitat not previously utilized by the fishery. The overall impacts to biological communities would be the same since most affected organisms would require more than a few months to recover from disturbance. Considered alone, Alternative 2 has a negligible to slight negative impact on the habitat because fishing activity outside of the trap/pot fisheries would continue to operate within the emergency closure that may disturb the benthic habitat.

6.3.3 Alternative 3

Alternative 3 would restrict the fishing and setting of trap/pot gear with buoys lines within MRA Wedge North February 1, 2023 through April 30, 2023. As discussed in Subsection 6.2.4, it is unlikely that gear would be relocated outside of this area because the remaining waters in LMA 1 require a Maine Zone permit. Only vessels with dual LMA permits for LMAs 1 and 3 would be

⁶ Per personal communication with Massachusetts Division of Marine Fisheries on Jan 12, 2023.

able to relocate into LMA 3 during the closure. Therefore, we assume lines would be removed from the water, reducing the likelihood that fishing gear will be relocated to habitat areas previously undisturbed by trap/pots. Removing the gear may decrease short-term disturbances, maintain local community structure, and may increase local lobster and/or Jonah crab abundance (Uhrin 2016). Similar to Alternative 2, if ropeless fishing expands widely under Alternative 3, the habitat could experience similar levels of disturbance as described under No Action, but it is unlikely that ropeless fishing levels will reach existing effort during the restricted area time period.

Relative to No Action, impacts of Alternative 3 on the habitat are expected to be negligible to slight positive because the emergency closure, MRA Wedge North, would only temporarily suspend fishing activity by the lobster and Jonah crab fishery and other fisheries would continue to operate in this area. Relative to Alternative 2, impacts of Alternative 3 on the habitat are expected to be negligible because while Alternative 3 restricts fishing activities in a larger area than the emergency area proposed in Alternative 2, the benefits of suspended fishing activities are limited. Given the temporary nature of the proposed emergency closure, habitat impacts are not considered to be significantly different. The closure that would be implemented under either Alternative will only be in place for 3 months, which is not sufficient time to allow the habitat to recover after a previously occurring disturbance. Relocated gear may disturb benthic habitat not previously utilized by the fishery. Alone, Alternative 3 has a negligible to slight negative impact on the habitat because other fisheries will continue to operate in the emergency closure area that may disturb the benthic habitat. The overall impacts to biological communities would be the same since most affected organisms would require more than a few months to recover from disturbance.

6.3.4 Comparison and Summary of the Alternatives

No quantitative criteria are available to formally compare the biological effect of the alternatives on habitat. No Action will maintain baseline levels of biological impacts on benthic habitats, negligible to slight negative impacts to habitat due to disturbance to benthic habitat.

Given the information above, in comparison to No Action, Alternative 2 and Alternative 3 are expected to have negligible to slight positive impacts on the MA LMA 1 habitat. If ropeless fishing is implemented in closed areas, it is not expected that effort under Alternative 2 or Alternative 3 will significantly increase the amount of gear that comes into contact with the seafloor and is unlikely to occur at levels similar to those prior to the closure. Similarly, Alternative 3 would likely have a negligible impact compared to Alternative 2. Compared to Alternative 2 and Alternative 3, No Action is expected to have negligible impacts on affected fish habitats. Considered on their own, Alternative 2 and 3 will likely have a negligible to slight negative impact on the environment due to continued disturbance from long trawls outside of the closure period.

6.4 Human Community Impacts of the Alternatives

6.4.1 *Economic Impacts of the Alternatives*

Alternative 1 (No Action) leaves the provisions of the Plan unchanged, and thus there would be no change in economic impacts relative to current regulatory requirements. Alternative 2, the Preferred Alternative, proposes to temporarily expand the Massachusetts Restricted Area (MRA) to include the MRA Wedge northward to Cape Ann, Massachusetts from February 1, 2023 through April 30, 2023. It would add 200 square miles (518 square kilometers) to the current MRA and bring slight short-term negative economic impacts to a number of lobster vessels in Southern Essex County, Suffolk County, Norfolk County and Northern Plymouth County. Alternative 3 expands the boundary of the MRA from February 1, 2023 through April 30, 2023 to include a larger area of approximately 1,297 square miles (3,359 square kilometers) that is bounded landward by the Massachusetts state waters, north at 42°52.58', seaward at 69°45', and south along the northern border of the MRA. Alternative 3 would impact more vessels, particularly those fishing from Northern Essex County, compared to Alternative 2.

The following subsection gives an overview of the analytic approach and results of economic impacts.

6.4.1.1 Analytic Approach

Vessels that fished within the proposed restricted area have two options to comply with the Emergency Rule: relocate their traps to the northern waters adjacent to the closure and keep fishing, or bring their traps back to dock and suspend fishing activity. Vessel Trip Report (VTR) gear distribution data from the past few seasons show that at least half of the traps were placed at the southern portion of the restricted areas proposed in Alternative 2 and Alternative 3. Because these traps are about 20 to 30 miles from the northern boundary of the proposed restricted area, they would be difficult to relocate, as the distance is beyond the normal range of fishing vessels.

For this analysis, we evaluated the two scenarios for the economic impacts on lobster vessels: we assume half of the vessels would relocate their traps, and the other half would remove their gear from the ocean and stop fishing. For relocated vessels, the cost differences come from reduced revenue in a different fishing area than their familiar and preferred, and extra operating costs to move gear. For vessels that stop fishing, the cost differences include lost revenue, gear removal costs, and saved operating costs from not fishing. The lower and higher end of cost estimates include the range of lost revenue of the relocated vessels, and a range of gear moving costs for all vessels (see details in the following section).

To estimate catch impacts of the proposed alternatives, we first used the VTR data for 2017-2021 to identify the vessels impacted by each alternative by using their self-reported fishing coordinates. Although the VTR coordinates only represent the general location of the vessels, it is the best available data for spatial analysis. We then determined the number of vessels and their landings weight for both lobster and Jonah crab. Finally, we calculated the landings value by multiplying the weight and price. The monthly average prices were calculated from NMFS dealer data for 2017-2021. All final values are adjusted to 2021 U.S. dollars by using GDP

deflator from U.S. Bureau of Economic Analysis (BEA 2022).

It should be noted that federal permitted fishing vessels that only carry lobster permits are not required to submit VTRs. In order to determine the total number of vessels fishing in this area, we divided the VTR landing value by the percent of VTR vessel coverage. NMFS federal permit data show that from 2017 to 2021, about 41 percent of Massachusetts federal lobster vessels in LMA 1 do not have VTR requirement, which means the landing value from VTR data need to be divided by 59 percent.

Another factor that needs to be considered is the operating cost savings from vessels that stop fishing. Vessel operating costs usually include fuel, bait, ice, fresh water, food and other incidentals. Labor costs are not included because many nearshore vessels are owner-operated, and mates are often paid based on landings rather than by the hour. These costs only occur when the vessel goes on a fishing trip. If a vessel does not fish, then these costs should be considered as savings. We use VTR data to determine the total number of fishing days, and then we apply an average daily operating cost to estimate the total savings.

For the operating costs of transporting gear back to the dock, or to resume fishing outside the restricted area, we assume that fishermen need three to six days to move all their traps around, and multiply that by the daily operating costs based on the average annual operating costs and fishing days for lobster vessels. The detailed results are presented in the next section.

6.4.1.2 Analysis Results

Vessel Lost Revenue

The emergency restricted area would be in place from February to April. During these months, few vessels were actively fishing and the landings were relatively low compared to summer/fall season. In Table 10, we list all lobster and Jonah crab vessels and landing values during February, March, and April from 2017 to 2021 for Alternative 2 and Alternative 3. We also provide the adjusted value by dividing the average value by 59 percent, as not all vessels were reporting their trips. We estimated that about 30 vessels would be affected by Alternative 2, with a total lobster and Jonah crab landing value of \$272,500 (Table 10). Alternative 3 would impact over 50 vessels with a total landing value of \$698,800 (Table 11).

Table 10: Number of affected vessels and landing values 2017-2021 under Alternative 2 (Preferred)

The number of affected vessels and landing values in Alternative 2 (Preferred) from 2017-2021 (in 2021 \$).

February			March		April	
Year	Number of vessels	Landing Value	Number of vessels	Landing Value	Number of vessels	Landing Value
2017	18	\$44,672	18	\$37,343	24	\$99,552
2018	25	\$130,445	18	\$64,155	19	\$144,306
2019	16	\$46,591	14	\$35,915	20	\$80,831
2020	19	\$47,206	12	\$22,222	14	\$33,499
2021	13	\$61,224	15	\$43,883	12	\$47,748
Average	18	\$66,028	15	\$40,704	18	\$81,187
Adjusted Average	31	\$112,004	26	\$69,046	30	\$137,719

Notes: 1. Landing values include both lobster and Jonah crab.

2. Both vessel number and landings are from federal VTR data. Based on federal vessel permit data, only 59 percent of Massachusetts federal lobster vessels are required to submit VTR, so the final number are adjusted proportionally to reflect the whole lobster fleet.

Table 11: Number of affected vessels and landing values 2017-2021 under Alternative 3

The number of affected vessels and landing values in Alternative 3 from 2017-2021 (in 2021 \$).

February			March		April	
	Number of vessels	Landing Value	Number of vessels	Landing Value	Number of vessels	Landing Value
2017	32	\$144,973	31	\$83,673	39	\$163,309
2018	48	\$488,671	35	\$264,741	39	\$391,033
2019	37	\$194,738	31	\$155,475	35	\$179,161
2020	42	\$250,343	32	\$99,482	31	\$102,029
2021	35	\$266,417	26	\$175,252	26	\$143,211
Average	39	\$269,028	31	\$155,725	34	\$195,749
Adjusted Average	66	\$456,358	53	\$264,159	58	\$332,052

Notes: 1. Landing values include both lobster and Jonah crab.

2. Both vessel number and landings are from federal VTR data. Based on federal vessel permit data, only 59 percent of Massachusetts federal lobster vessels are required to submit VTR, so the final number is adjusted proportionally to reflect the whole lobster fleet.

Vessel Operating Cost Savings

Vessels that decide to stop fishing during closure months could save some operating costs. We estimated the vessel operating costs based on the cost surveys conducted by the Social Science Branch (SSB) of the Northeast Fisheries Science Center for fishing years 2011, 2012, and 2015. Survey data show that the average annual operating costs for lobster vessels in the Northeast Region is about \$50,365 (in 2021 dollars). Table 12 displays the potential cost savings. We calculate the percentage of trips in each month, and then assign the operating cost to each month based on the trip percentage. At the end, we multiply the cost per vessel and the affected vessel number to get the total cost saving for each month.

Table 12: *Cost savings for vessels that stop fishing during closure months*
Cost savings for vessels that stop fishing during closure months (in 2021 \$).

	Month	Affected vessel number	Annual cost per vessel	Closure month trip %	Monthly cost per vessel	Total cost
Alternative 2	Feb	15	\$50,365	4.77%	\$2,403	\$37,092
	Mar	13	\$50,365	3.31%	\$1,669	\$21,806
	Apr	15	\$50,365	4.10%	\$2,067	\$31,210
Alternative 3	Feb	33	\$50,365	4.77%	\$2,403	\$79,075
	Mar	26	\$50,365	3.31%	\$1,669	\$43,894
	Apr	29	\$50,365	4.10%	\$2,067	\$59,614

Notes: 1. We assume that half of the vessels would stop fishing.

2. Annual cost per vessel is based on SSB survey results.

3. Closure month trip percentage is from VTR data.

6.4.1.3 Final Results

We estimated that 26 to 31 vessels would be affected by Alternative 2 and over 50 vessels affected by Alternative 3. For Alternative 2, the total costs including gear transportation costs and lost revenue range from \$339,000 to \$608,000 for February to April. For vessels moving their gear to new fishing grounds, the total costs are around \$139,000 to \$278,000, about \$3,200 to \$6,400 per vessel; for vessels that stop fishing, the total costs are around \$200,000 to \$331,000, about \$4,600 to \$7,600 per vessel (Table 13). For Alternative 3, the total cost ranges from \$898,000 to \$1,453,000. Total costs for vessels moving their gear to new fishing grounds range from \$290,000 to \$581,000, about \$3,300 to \$6,600 per vessel. Total costs for vessels that stop fishing are from \$608,000 to \$872,000, about \$6,900 to \$9,900 per vessel (Table 14).

Table 13: Economic impacts of Alternative 2 (Preferred) by month
Economic impacts of Alternative 2 (Preferred) by month (in 2021\$).

	Feb		March		April		Total	
	Lower	Higher	Lower	Higher	Lower	Higher	Lower	Higher
Relocating costs (half vessels)								
Lost revenue	\$2,800	\$5,600	\$1,726	\$3,452	\$3,443	\$6,886	\$7,969	\$15,938
Gear moving	\$46,310	\$92,619	\$39,185	\$78,370	\$45,292	\$90,583	\$130,786	\$261,572
Sum	\$49,110	\$98,219	\$40,911	\$81,822	\$48,735	\$97,469	\$138,755	\$277,511
Stop fishing costs (half vessels)								
Lost revenue	\$56,002	\$56,002	\$34,523	\$34,523	\$68,860	\$68,860	\$159,385	\$159,385
Gear moving	\$46,310	\$92,619	\$39,185	\$78,370	\$45,292	\$90,583	\$130,786	\$261,572
(Cost savings)	\$37,092	\$37,092	\$21,806	\$21,806	\$31,210	\$31,210	\$90,107	\$90,107
Sum	\$65,219	\$111,529	\$51,903	\$91,088	\$82,942	\$128,233	\$200,064	\$330,850
Total cost	\$114,329	\$209,748	\$92,814	\$172,910	\$131,676	\$225,703	\$338,819	\$608,361

Notes:

- 1 We estimate lost revenue of the relocating vessels to be between 5 and 10 percent of the total landing value.
2. We estimate gear moving costs to take between 3 and 6 days at \$1,000/day.

Table 14: Economic impacts of Alternative 3 by month
Economic impacts of Alternative 3 by month (in 2021 \$).

	Feb		March		April		Total	
	Lower	Higher	Lower	Higher	Lower	Higher	Lower	Higher
Relocating costs (half vessels)								
Lost revenue	\$11,409	\$22,818	\$6,604	\$13,208	\$8,301	\$16,603	\$26,314	\$52,628
Gear moving	\$98,726	\$197,452	\$78,879	\$157,758	\$86,512	\$173,025	\$264,117	\$528,234
Sum	\$110,135	\$220,270	\$85,483	\$170,966	\$94,814	\$189,627	\$290,431	\$580,862
Stop fishing costs (half vessels)								
Lost revenue	\$228,179	\$228,179	\$132,079	\$132,079	\$166,026	\$166,026	\$526,285	\$526,285
Gear moving	\$98,726	\$197,452	\$78,879	\$157,758	\$86,512	\$173,025	\$264,117	\$528,234
(Cost savings)	\$79,075	\$79,075	\$43,894	\$43,894	\$59,614	\$59,614	\$182,584	\$182,584
Sum	\$247,829	\$346,555	\$167,064	\$245,943	\$192,924	\$279,437	\$607,818	\$871,935
Total cost	\$357,964	\$566,825	\$252,547	\$416,908	\$287,738	\$469,064	\$898,249	\$1,452,797

Notes:

- 1 We estimate lost revenue of the relocating vessels to be between 5 and 10 percent of the total landing value.
2. We estimate gear moving costs to take between 3 and 6 days at \$1,000/day.

6.4.2 Social Impacts of the Alternatives

Table 15 presents socio-economic data for each county identified as potentially vulnerable to social impacts due to Alternative 2 or Alternative 3. Essex and Plymouth counties have the most potentially affected vessels and land a large amount of seafood that are from the regulated gear. They also have a higher commercial reliance score than Suffolk and Norfolk counties. Norfolk County has a small number of vessels, but almost all of its seafood landings are from the lobster and Jonah crab trap/pot fishery. The fishermen in these two counties are the most impacted by this Emergency Rule at the individual level, but not the community level. Norfolk County has the highest income level and lowest unemployment rate. Its low commercial engagement rate indicates that fishermen might have more alternative occupations when fishing is not available. The only major port in Suffolk County is Boston Harbor. It lands a small amount of lobsters and Jonah crabs from a very limited number of vessels. Both Suffolk and Norfolk counties have a much lower commercial reliance score than Essex and Plymouth counties.

Considering all factors, trap/pot vessels in Plymouth County could be the community most vulnerable to the implementation of this Emergency Rule. Essex County could also be heavily

impacted, but its fisheries are more diversified so individual fishermen may be more flexible. The Norfolk County fishery would be totally shut down during the restricted time period, which came with only one month's notice, so although the community has more access to alternative jobs than some other counties, it may be difficult to make up lost income given the short notice of the closure. Suffolk County might be the least vulnerable to the Emergency Rule.

Table 15: Socio-economic profile of affected communities - Harvest Parameters

State	County	Major Ports	Top Species by Value	2021 lobster/Jonah Crab Harvest (\$)	Lobster/Jonah Crab Value as Percentage of Total Value	Number of lobster trap/pot Vessels
MA	Essex	Gloucester, Rockport, Marblehead	Lobster, cod, pollock	62,781,295	52%	552
MA	Suffolk	Boston Harbor	Cod, lobster, pollock	2,768,326	14%	72
MA	Norfolk	Cohasset	Lobster, softshell clam, bluefin tuna	1,920,738	91%	32
MA	Plymouth	Plymouth, Scituate, Hingham	Lobster, oysters, cod	22,280,221	60%	315

Data source: ACCSP dealer report 2021

6.4.3 Comparison and Summary of Impacts to Human Communities

Alternative 1 (No Action) would maintain the status quo, which has a slight negative to moderate positive impact on fishing communities. Alternative 2, the Preferred Alternative, is expected to have a slight negative impact on the fishing communities impacted by this action. Overall, the economic impacts of the Alternative 2 results in an estimated total cost (including lost revenue) of from \$339,000 to \$608,000 with about 30 affected vessels, compared to No Action.

Alternative 2 would impact lobster and Jonah crab vessels in Southern Essex County, Suffolk County, Norfolk County, and Northern Plymouth County. Vessels in Plymouth County could be the most vulnerable to the Emergency Rule, while Suffolk County might be the least vulnerable. Alternative 3 is expected to have a moderate negative impact on the human community VEC, as defined here. Alternative 3 is estimated to impact 53 to 66 vessels for a total estimated cost (including lost revenue) of \$898,000 to \$1,453,000, compared to No Action. Alternative 3 has similar social impacts to Alternative 2, except it will affect a few more vessels in Essex County that fish in the Northern waters offshore and north of Cape Ann.

6.5 Cumulative Impacts of the Alternatives


6.5.1 Introduction

A cumulative effects analysis (CEA) is required by the Council on Environmental Quality (CEQ; 40 CFR part 1508.7) and NOAA policy and procedures for the National Environmental Policy Act (NEPA), found in NOAA Administrative Order 216-6A (Companion Manual, January 13, 2017). A CEA examines the impact of the actions in conjunction with other factors that affect the

physical, biological, and socioeconomic resource components of the affected environment. The purpose of the CEA is to ensure that federal decisions consider the full range of an action's consequences, incorporating this information into the planning process. The CEQ guidelines recognize that it is not practical to analyze the cumulative effects of an action from every conceivable perspective, but, rather, the intent is to focus on those effects that are truly meaningful. This CEA is based upon a more detailed analysis that was conducted in a 2021 Final Environmental Impact Statement (NMFS 2021) and follows the steps depicted in Figure 14. The CEA analysis relies upon the impact designations defined in Subsection 6.1 to determine the cumulative effects of each valued ecosystem component (VEC).

Figure 14: Cumulative effects analysis steps

Cumulative effects analysis steps, and how they inform the cumulative effects analysis (adapted from Canter 2012).



6.5.1.1 Geographic and Temporal Scope

The geographic scope of this CEA is focused on the southern portion of Lobster Management Area 1 (LMA 1) that includes waters from New Hampshire state waters south to the Massachusetts Restricted Area boundary at 40° 12' N. latitude bounded on the west by Massachusetts state waters, and on the east by the LMA 1/LMA 3 boundary. This is an area currently subject to the requirements of the Plan and includes the seawater and sea bottom of the Atlantic Ocean within U.S. jurisdiction. Though some of the activities included in this analysis do not occur within the small geographic scope of this Environmental Assessment, they are still considered in this analysis for each VEC due to the potential for negative impacts on the right whale population. This CEA focuses on the lobster and Jonah crab fishery given this is the trap/pot fishery that will most likely be impacted by this action.

The temporal scope of the analysis varies by resource. In all instances, the analysis attempts to take into account past (primarily the past two decades), present, and reasonably foreseeable future actions (within five years) that could affect valuable physical, biological, or socioeconomic resources. The discussion here focuses on impacts of management actions as well as the direct impact of potential stressors: interactions with commercial and recreational fisheries, vessel strikes, pollution, noise, climate change, renewable energy development, oil and gas development, harmful algal blooms, and prey availability. Stressors that are not expected to impact a VEC may be noted but will not be analyzed.

6.5.1.2 Analysis of Total Cumulative Effects

A CEA ideally makes effect determinations based on the combination of: 1) impacts from past, present, and reasonably foreseeable future actions; 2) status quo condition of the VECs (the combined effects from past, present, and reasonably foreseeable future actions plus the present condition of the VEC); and 3) impacts of the alternatives under consideration for this action.

6.5.2 Summary of Direct and Indirect Impacts of the Alternatives

The direct and indirect impacts of the alternatives on the VECs were discussed in Subsections 6.1 to 6.4 and summarized in Table 16.

Table 16: Direct and indirect impacts of the alternatives on Valued Ecosystem Components

<i>Alternatives</i>	<i>Protected Species</i>	<i>Habitat</i>	<i>Human Communities</i>
Alternative 1 (No Action)	High Negative to Moderate Negative Mortality and serious injury would continue to occur and impact ESA-listed species' population health. More so for right whales and other large whales, and to a lesser degree for other ESA-listed or MMPA protected species	Negligible to Slight Negative Areas with trawls above 15 traps per trawl may have a short-term impact.	Slight Negative to Moderate Positive Positive in that there are no new impacts or costs to harvesters and markets, but the lack of recovery of whale species has a slight negative impact on public intrinsic value benefits due to whale population declines.
Alternative 2 (Preferred)	Moderate Negative to Slight Negative Would reduce entanglement risk for ESA-listed and MMPA protected species. However, risk of interactions will not be entirely eliminated.	Negligible to Slight Negative Areas with trawls above 15 traps per trawl may have a short-term impact.	Slight Negative Fisheries would experience extra costs and catch reduction in the short term that could ease over the long term.
Alternative 3	Negligible to Slight Negative Would reduce entanglement risk for ESA-listed and MMPA protected species. However, risk of interactions will not be entirely eliminated.	Negligible to Slight Negative Areas with trawls above 15 traps per trawl may have a short-term impact.	Moderate Negative Catch reduction could be significant.

6.5.3 Status Quo Conditions

The status and trends of each VEC is summarized in Table 17. Additional details can be found in the Affected Environment Section (Section 5) of this document and Chapter 4 in the recent 2021 FEIS (NMFS 2021).

Table 17: Summary of the current status and trends of the Valued Ecosystem Components

VEC	Historical Conditions	Current Conditions	Possible Future Conditions	Implications of Conditions Relative to Sustainability
Protected Species	Stocks were depleted by whaling and other anthropogenic impacts.	Right, fin, and sei whales are endangered. Right whale stock is declining, humpbacks are slightly increasing, and the trends of the others are unknown. Trends are unavailable for other whale species.	Under current conditions, right whales are likely to continue declining. Certain protected species may be more resilient to future changes while others may remain small or continue to decline.	Certain stocks that are still depleted are still vulnerable to additional anthropogenic stressors and population decline (right whales and fin whales).
Habitat	The habitat has slowly degraded over time with increasing exposure to anthropogenic stressors.	The habitat is rapidly shifting from historical baselines from the impacts of climate change as well as other anthropogenic stressors.	Shifts in habitat features are expected to continue as the climate shifts and alters the frequency and magnitude of disturbance.	The habitat is vulnerable to additional disturbance.
Human Community	American lobster stocks have been abundant in GOM but depleted in SNE waters; Jonah crab fishery was supplement of lobster fishery.	Total lobster landings peaked in 2015 and started to decrease. GOM represents about 80 percent of all lobster landings; Southern MA and RI landed the most Jonah crabs.	GOM lobster landings are trending down and SNE stock stays depleted; more Jonah crabs will be landed from SNE.	Target species, lobster and Jonah crab, are vulnerable to anthropogenic and environmental stressors, posing a threat to fishing communities that depend on commercial fisheries.

6.5.4 Past, Present, and Reasonably Foreseeable Future Actions

Detailed information on the past, present, and reasonably foreseeable future actions that may impact this action were evaluated as part of the cumulative effects assessment found in the environmental assessment prepared for the last substantial modification to the Plan (NMFS 2021). Much of that information remains applicable, though unlikely to affect the implementation of Alternative 2 because the temporal scope of the action is limited to the months of February, March, and April in 2023. The following provides a brief summary of updates on the pertinent fishing activities, and the proposed rule expanding reporting and vessel tracking in the American lobster and Jonah crab fishery. Past, present, and reasonably foreseeable future fisheries management actions were summarized in more detail recently in an FEIS released on July 2, 2021 (NMFS 2021). A summary of the impact of past, present, and

anticipated actions on each VEC is summarized here, but more detail can be found in Chapter 8 of the FEIS (NMFS 2021).

6.5.4.1 Fisheries Management

Fishery management actions include the creation of a new Fishery Management Plan (FMP) and additional amendments and addenda that modify how the fishery is conducted. These amendments and addenda can include actions such as quotas, trap reductions, administration of taxes, and guidelines on how data is collected and shared with management agencies. These actions can have a variety of impacts on the economic aspects of fisheries as well as the environment.

A list of past, present, and reasonably foreseeable future actions listed in Table 8.3 of the FEIS (NMFS 2021). There are several additional management actions underway that affect the Northeast American lobster and Jonah crab fishery. Currently, the Atlantic States Marine Fisheries Commission is reviewing public comments on the proposed Draft Addendum XXIX to Amendment 3 to the American Lobster FMP and Draft Addendum IV to the Jonah Crab FMP for the purposes of collecting high resolution spatial and temporal data to characterize fishing effort in the federal American lobster and Jonah crab fisheries for management and enforcement needs.

Protected Species

FMPs and their amendments can mitigate the impact of fishing gear on protected large whale species. The amendments and addenda referenced in this analysis were primarily intended to optimize fishing practices, restrict overfishing, manage bycatch, and gather information to better manage the stock. Lobster and Jonah crab management measures that reduce rope in the water column would be an improvement compared to current conditions; improved reporting and monitoring would inform future management and may have an indirect net positive impact; and modifications to maintain or restrict fishing on other species would likely cause negligible impacts. However, any fishing generally has a negative effect on protected species because any gear in the water has some risk of interaction. While fisheries management can mitigate some of this, the overall effect is anticipated to be between slight negative to moderate negative. Future actions that aim to improve monitoring of lobster and Jonah crab trap/pot fisheries are likely to positively impact protected species by improving data collection on fishing effort, which will inform updates to the Large Whale Decision Support Tool analysis and discussions to support fishery management decisions related to protected species, marine spatial planning, and offshore enforcement. Management actions in the past, present, and reasonably foreseeable future are likely to benefit or have negligible impacts on protected resources. Overall, the cumulative effects are likely to be a slight negative to moderate negative impact on the protected species VEC.

Habitat

Trap/pot fisheries that operate longer trap trawls could have a slightly deleterious impact on the habitat. Setting quotas and trap limits that reduce gear on the bottom are likely indirectly better

for the habitat than unmanaged fisheries. Overall, the impact of trap/pot fisheries management on habitat is considered to be negligible to slight negative.

Human Communities

The aims of many of the fishery management actions in the past, present, and reasonably foreseeable future aim to improve the maintenance of the target stock and mitigating bycatch. Both of these goals are likely to have a slight positive impact on the economics of the fishery by allowing the continuation of a healthy fishery as a source of income for fishing communities.

6.5.4.2 Non-Fisheries Management

Several management actions have been implemented to mitigate the impact of stressors on protected species, habitats, and human communities. These include actions to reduce the impact of pollution, climate change, entanglement, and vessel strikes on the environment and protected species. The impact of other past, present, and foreseeable future conservation actions are discussed below.

Protected Species

Conservation mitigation measures aim to reduce the impact of known human or environmental stressors. Mitigating the impact of multiple stressors in the environment by protecting habitats and habitat quality can reduce the overall stress by reducing the energy necessary to adapt to new baselines. Many stressors are known to negatively impact large whales and, therefore, mitigating actions are expected to improve impacts on this VEC. Actions like speed reductions for vessels and on-board observers would also benefit other large whale species. However, the risk of entanglement with buoy lines and vessel strikes remains, albeit less so for entanglement, after these mitigation measures are taken. Therefore, ESA-listed species of large whales (right, fin, and sei whales) are expected to experience moderate negative to slight positive impacts, and MMPA-protected species of large whales (humpback and minke whales) are expected to have slight positive impacts (i.e., PBR not exceeded).

Habitat

Some of the environmental mitigation actions that occurred in the past, present, and reasonably foreseeable future are likely to reduce the number or magnitude of stressors on fish habitat and benthic organisms in the Northeast Region, particularly those related to regulating pollutants. Pollution and climate change can contribute to habitat degradation through mechanical disruption of habitat structure and negative impacts on the health of organisms (see the next section). Measures that directly protect habitats, address the effects of climate change, or protect water and sediment quality via pollution mitigation will prevent additional environmental degradation as a result of these stressors. These measures are expected to have positive impacts on marine habitats. Other regulations likely have a negligible impact on habitat, such as vessel strike regulations that are not expected to impact the physical environment. However, continued fishing effort will continue to impact habitats. The net impact of all actions is likely slightly negative to slightly positive.

Human Communities

Most of the mitigation actions included in this analysis are expected to have negligible impact on the fishing communities that rely on fisheries. Actions that have been implemented to mitigate entanglement, such as reductions in gear, expensive changes in gear configurations, and exclusion from areas they have fished in the past, likely have a negative impact on this VEC, whereas those that have a positive impact on fishery habitat are expected to have a slight positive impact on human communities by supporting healthy fisheries.

6.5.4.3 Non-Management

There are several anthropogenic actions that could potentially impact the VECs included in this analysis, including fishing, aquaculture, manufacturing, agriculture, construction, oil and gas activities, wind farm exploration and operation, military activities, shipping, and climate change. These activities can have an impact individually as well as collectively and should be considered when proposing management actions. The impact of these individual activities on the VECs are discussed in greater detail in Table 8.4 of a 2021 FEIS and summarized below (NMFS 2021).

Protected Species

Human activities have directly or indirectly increased the number and magnitude of stressors protected species are exposed to, which is a concern for vulnerable protected species such as the North Atlantic right whale. Climate change, vessel strikes, entanglement, and Canadian mortalities are all anticipated to have high negative impacts on protected species due to the severity of the impact on the declining right whale population in particular, though these do impact other large whale species in this VEC as well. Aquaculture, offshore wind farm exploration and operation, oil and gas related activities, prey availability, and harmful algal blooms are estimated to have a moderate negative impact on protected species. The use of aquaculture and offshore wind farms specifically are expected to grow in the foreseeable future and there is concern for the increase in risk of entanglement, vessel traffic, and noise on protected species. Changes in prey availability related to climate change are expected to become more pronounced and variable over time. Underwater anthropogenic noise more broadly likely has a slight to moderate negative impact on protected species depending on the source, severity, duration, and species. McCauley et al. (2017) suggest that noise pollution from seismic surveys cause significant mortality to zooplankton populations, an important prey species of large whales. Pollution and water quality likely have the least impact on protected species (slight negative) since baleen whales are typically less at risk of bioaccumulation compared to higher trophic level marine mammals. Together, non-management human activities have a moderate negative impact on this protected species VEC.

Habitat

Climate change is the factor that likely has the greatest impact on the habitat VEC and is anticipated to be a high negative. Offshore wind farms, oil and gas activities, and harmful algal blooms generally have a moderate negative impact on marine habitats due to the level of disturbance and disruption they can cause, though there are currently no new plans for offshore

wind farms or oil and gas activities within the specific geographic scope of this EA. Pollution has a slight negative impact on marine habitats, partly due to past mitigation that was discussed in the previous section. Aquaculture likely has a negligible to slight negative impact on marine habitats in this area, though there are few aquaculture projects currently in the area analyzed in this EA. Overall, non-management human activities likely have a moderate negative to high negative impact on this VEC, given past, present and foreseeable future activities.

Human Communities

Climate change will also have a high negative impact on human communities due to the reliance of these communities on natural resources that are already being affected. Offshore wind development, oil and gas development, and harmful algal blooms likely have a moderate negative impact on fishing communities in general. However, the extent to which the fishing communities in the geographic scope of this are impacted by offshore wind or oil and gas activities may be slightly negative to moderate negative within the scope of this analysis. Entanglements have a slight negative impact on fishing communities. Aquaculture is estimated to have a slight negative to negligible impact on fishing communities. The impact of noise and pollution has a negligible impact on the fishing community VEC. When combined, non-management human activities have a negligible to high negative impact on fishing communities.

6.5.5 Cumulative Effects Analysis

A summary of the cumulative impacts on all VECs for Alternative 2 (Preferred) is summarized in Table 18.

Table 18: Summary table of the final cumulative impacts analysis of the Preferred Alternative (Alternative 2) on all three VECs

<i>VECs</i>	<i>Direct and Indirect Impacts</i>	<i>Existing Conditions</i>	<i>All Management Actions and Stressors</i>	<i>Cumulative Impacts</i>
Protected Species	Slight to Moderate Negative Would reduce entanglement risk for ESA-listed and MMPA protected species. However, interaction risk will not be entirely eliminated.	Negative Several protected species are still listed as endangered or threatened.	Moderate Negative to Slight Positive Fisheries negatively impact large whale species, though some management actions may have mitigated the risk. Non-fishery management actions likely improved ocean quality and reduced gear encounters, which benefitted large whales. Anthropogenic and natural stressors have had negative impacts on the VECs and likely will continue to do so in the future.	Slight Negative to Negligible Continued catch and effort controls, is likely to reduce gear encounters through effort reductions. Additional management actions taken under ESA/MMPA should also help mitigate the risk of gear interactions.
Habitat	Negligible to Slight Negative Areas with trawls above 15 traps per trawl may have a short-term impact.	Negative Habitats have experienced degradation from human activities and are shifting as a result of climate change.	Slight Negative to Slight Positive Fishery management actions likely have negligible to slight negative impacts on habitat. Non-fishery management actions likely improved ocean quality, which benefitted habitats. Anthropogenic and natural stressors have had moderate negative impacts on habitats.	Negligible to Slight Positive Continued management is not expected to measurably change habitat quality and existing cumulative impacts.
Human Community	Slight Negative Fisheries would experience extra costs and catch reduction in the short term.	Negative Commercial fisheries are shifting as a result of climate change.	Slight Negative to Slight Positive Overall, fisheries management positively impacts fishing communities, though certain management actions may have had a short term negative effect. Non-fishery management actions likely improved fisheries. Anthropogenic and natural stressors have had negative impacts.	Slight Negative to Slight Positive Continued fishery management is expected to positively benefit fishing communities but conservation measures will likely negatively impact fishing communities, except for the positive social benefits expected from protecting whale species.

6.6 Summary

We analyzed the impacts of all alternatives on physical habitat, protected species, and human communities. The impacts of Alternative 2 (Preferred) considered on each VEC described in the Affected Environment are in Section 6 and are summarized here.

When considered in conjunction with all other pressures placed on protected species by past, present, and reasonably foreseeable future actions, the Emergency Rule is expected to slightly reduce the impact of human activities (i.e. entanglement risk) on ESA-listed (right, fin, and sei whales) and MMPA protected species (humpback and minke whales). Considered alone, ESA-listed and MMPA protected species would be moderately negative to slightly negatively impacted because this proposed emergency action does not eliminate the potential for all interaction risk between fishing gear and marine mammals that could result in takes above PBR. Considered alone, Alternative 2 has a negligible to slight negative impact on habitat due to continued disturbance from long trawls outside of the closure period. Alternative 2 will have a slight negative impact on fishing communities impacted by this action.

When Alternative 2 is considered in conjunction with all other impacts from past, present, and reasonably foreseeable future actions, it is not expected to result in any significant impacts, positive or negative. No significant cumulative effects on the human environment are associated with the proposed action and, therefore, a “Finding of No Significant Impact” is warranted.

7 APPLICABLE LAWS AND REGULATIONS

7.1 Endangered Species Act

Section 7 of the Endangered Species Act (ESA) requires federal agencies to ensure that their actions do not jeopardize the continued existence of any species listed as threatened or endangered or result in the destruction or adverse modification of the Critical Habitat of listed species. The ESA requires the “action” agency to consult with an “expert” agency to evaluate the effects a proposed agency action may have on a listed species. If the action agency determines through preparation of an environmental assessment or informal consultation that the Preferred Alternative is “not likely to adversely affect” listed species or Critical Habitat, formal consultation is not required so long as the expert agency concurs.

A section 7 consultation on the Atlantic Large Whale Take Reduction Plan (Plan) was completed on May 25, 2021 and determined that the Plan would have wholly beneficial effects to ESA-listed species or their critical habitat. An informal consultation concluded on February 11, 2022 that the 2022 emergency regulations modifying the Plan fall within the scope of the 2021 consultation. Given that the emergency rulemaking falls within the scope of the existing section 7 consultation, reinitiation of the existing consultation is not required.

7.2 Marine Mammal Protection Act

Under the Marine Mammal Protection Act (MMPA), federal responsibility for protecting and conserving marine mammals is vested with the Departments of Commerce (NMFS) and Interior (U.S. FWS) and the MMPA is the authority under which much of the current rulemaking is being undertaken. The MMPA prohibits the “take” of marine mammals, with certain exceptions, in waters under U.S. jurisdiction and by U.S. citizens on the high seas. The primary management objective of the MMPA is to maintain the health and stability of the marine ecosystem, with a goal of obtaining an optimum sustainable population of marine mammals within the carrying capacity of the habitat. Section 118 of the MMPA specifies that NMFS develop and implement Take Reduction Plans to assist in the recovery or prevent the depletion of strategic marine mammal stocks that interact with Category I and Category II fisheries, which are fisheries that cause frequent (Category I) or occasional (Category II) serious injuries and mortalities to marine mammals. The goal is to reduce these takes incidental to fishing activities to levels below the Potential Biological Removal level, defined as the maximum number of animals, not including natural mortalities that may be removed from a marine mammal stock while allowing that stock to reach or maintain its optimum sustainable population.

For species not managed under take reduction plans, the MMPA requires consultation within NMFS if impacts on marine mammals are unavoidable. An analysis of the potential impact of the management actions on all marine mammal species that may be affected by this management action are discussed in Subsection 6.2. NMFS has reviewed the impacts of this action on marine mammals and concluded that the management actions proposed are consistent with the provisions of the MMPA.

7.3 Paperwork Reduction Act

This action contains no information collection requirements under the PRA of 1995.

7.4 Magnuson-Stevens Fishery Conservation and Management Act including Essential Fish Habitat

The Essential Fish Habitat (EFH) provisions of the Magnuson-Stevens Fishery Conservation and Management Act (MSA) require the National Marine Fisheries Service to provide recommendations to federal and state agencies for conserving and enhancing EFH if a determination is made that an action may adversely impact EFH. An EFH consultation, as required under the MSA, concluded on January 12, 2023 that adverse impacts to EFH have been minimized to the extent practicable and no further EFH Conservation Recommendations pursuant to 50 CFR 600.925(a) were provided.

7.5 Information Quality Act (Public Law 106-554)

The Information Quality Act (IQA) directed the Office of Management and Budget to issue government-wide guidelines that “provide policy and procedural guidance to federal agencies for ensuring and maximizing the quality, objectivity, utility, and integrity of information (including statistical information) disseminated by federal agencies.” Under the NOAA guidelines, the Plan is considered a Natural Resource Plan. It is a composite of several types of information,

including scientific, management, and stakeholder input, from a variety of sources. An IQA pre-dissemination review was completed on January 13, 2022. Compliance of this document with NOAA guidelines is evaluated below.

- **Utility:** The information disseminated is intended to describe the current management actions and the impacts of those actions. A diversity of public interests may be affected by this emergency rule, but not limited to the Massachusetts fishing community, scientists, conservation groups, and state and federal resource managers. This document presents information in a manner that is understandable to a wide range of users and thoroughly explains why NMFS is publishing an emergency final rule, requirements of the rulemaking action, policy and science justifying the action, and the potential effects of the action.
- **Integrity:** Information and data, including statistics that may be considered as confidential, were used in the analysis of impacts associated with this document. This information was necessary to assess the biological, social, and economic impacts of the alternatives considered as required under the National Environmental Policy Act for the preparation of an Environmental Assessment statement/regulatory impact review. NMFS complied with all relevant statutory and regulatory requirements as well as NMFS policy regarding confidentiality of data. For example, confidential data were only accessible to authorized federal employees and contractors for the performance of legally required analyses. In addition, confidential data are safeguarded to prevent improper disclosure or unauthorized use. Finally, the information to be made available to the public was done so in aggregate, summary, or other such form that does not disclose the identity or business of any person.
- **Objectivity:** The NOAA Information Quality Guidelines for Natural Resource Plans state that plans must be presented in an accurate, clear, complete, and unbiased manner. Because take reduction plans and their implementing regulations affect such a wide range of interests, NMFS strives to draft and present new management measures in a clear and easily understandable manner with detailed descriptions that explain the decision making process and the implications of management measures on marine resources and the public. Although the alternatives considered in this document rely upon scientific information, analyses, and conclusions, clear distinctions were drawn between policy choices and the supporting science. In addition, the scientific information relied upon in the development, drafting, and publication of this EA was properly cited and a list of references was provided. Finally, this document was reviewed by a variety of biologists, policy analysts, economists, and attorneys from the Greater Atlantic Region, the Northeast Fisheries Science Center, NMFS Headquarters, and the Office of General Council. In general, this team of reviewers has extensive experience with the policies and programs established for the protection of marine mammals, and specifically with the development and implementation of the Plan. Therefore, this Natural Resource Plan was reviewed by technically qualified individuals to ensure that the document was complete, unbiased, objective, and relevant. This review was conducted at a level commensurate with the importance of the interpreted product and the constraints imposed by legally-enforceable deadlines.

7.6 Administrative Procedure Act

The Administrative Procedure Act (APA) establishes procedural requirements applicable to informal rulemaking by federal agencies. The purpose of the APA is to ensure public access to the federal rulemaking process and to give the public notice and an opportunity to comment before the agency promulgates new regulations. Pursuant to 5 U.S.C. 553(b)(B), the Assistant Administrator for Fisheries finds prior notice and public comment is not required because it would be impracticable and contrary to the public interest. North Atlantic right whale (right whale) monthly distribution data identifies risk in unrestricted waters surrounded on three sides by the expanded Massachusetts Restricted Area (MRA) from February 1 through April 30. The 2021 expansion of the geographic extent of the MRA to include Massachusetts state waters north to the New Hampshire border mirrors the Massachusetts 2021 modification of the state water closure (322 CMR 12.04(2)). With the implementation of the MRA Expansion, approximately 200 square miles (518 square kilometers) of federal waters remain open to trap/pot fishing between state and federal closures, creating a wedge where 2021 and 2022 data indicates that trap/pot gear is concentrated during the closure period (MRA Wedge; Figure 9). During aerial surveys in February, March, and April 2018-2021, the Center for Coastal Studies (CCS) and Northeast Fisheries Science Center (NEFSC) observed right whales within the MRA Wedge alongside the presence of aggregated fishing gear (Figure 9, Figure 10). The gear in this area is thought to be a mix of actively fished gear and staged gear that is placed in preparation for the opening of MRA federal waters in May. This, in addition to the gear being actively fished outside of the buoy line closure area, could increase gear density in the area. Given the dense concentration buoy lines in an area with persistent right whale presence, the MRA Wedge area presents an imminent entanglement threat.

In summary, this emergency rule is necessary to protect right whales in an area of elevated risk in Massachusetts Bay from February 1, 2023 through April 30, 2023. Providing prior notice through proposed rulemaking and a public comment period in the normal rulemaking process would be impracticable and contrary to the public interest, as it would delay these necessary and time sensitive emergency measures. Delay would prevent NMFS from meeting its obligations to protect right whales from entanglements. For the reasons outlined above, NMFS finds it impracticable and contrary to the public interest to provide prior notice and public comment on these emergency measures. For the same reasons, NMFS finds good cause under 5 U.S.C. 553(d)(3) to waive the 30-day delay of effectiveness period for this.

7.7 Coastal Zone Management Act

Section 307(c)(1) of the Coastal Zone Management Act of 1972 (CZMA) requires that all federal activities that affect any land or water use or natural resource of the coastal zone be consistent with approved state coastal zone management programs to the maximum extent practicable.

NMFS has not conducted a full consistency review under the requirements of section 307 of the CZMA for this Emergency Rule. NMFS has determined that the provisions of 15 CFR 930.32(b), which allow for deviation due to exigent circumstances that present a federal agency with a substantial obstacle that prevents complete adherence to the approved program, apply. NMFS will consult with the relevant Massachusetts state agency to the extent possible prior to

the effective date of the Emergency Rule and attempt to seek concurrence. Once the exigent circumstances have passed, if NMFS is still carrying out an activity with coastal effects, NMFS will comply with the applicable provisions of the CZMA consistency review requirements to ensure that the rule is consistent to the maximum extent practicable with the enforceable policies of those State's management programs. Once NMFS has addressed the exigent circumstance through issuance of this rule, NMFS will provide the state agencies with a description of its action and its coastal effects.

7.8 Executive Order (E.O.) 13132 Federalism

E.O. 13132, otherwise known as the Federalism E.O., was signed by President Clinton on August 4, 1999, and published in the Federal Register on August 10, 1999 (64 FR 43255). This E.O. is intended to guide federal agencies in the formulation and implementation of "policies that have federal implications." Such policies include regulations, legislative comments or proposed legislation, and other policy statements or actions that have substantial direct effects on the states, on the relationship between the national government and the states, or on the distribution of power and responsibilities among the various levels of government. This rule does not contain policies with federalism implications as that term is defined in E.O. 13132.

7.9 Regulatory Flexibility Act

The purpose of the Regulatory Flexibility Act (RFA) is to reduce the impacts of burdensome regulations and recordkeeping requirements on small businesses. The Emergency Rule is exempt from the procedures of the RFA because the rule will not include prior notice or an opportunity for public comment.

7.10 E.O. 12866 Regulatory Planning and Review

The purpose of E.O. 12866, otherwise known as Regulatory Planning and Review, is to enhance planning and coordination with respect to new and existing regulations. This E.O. requires the Office of Management and Budget to review regulatory programs that are considered to be "significant." This Emergency Rule has been determined to be not significant for the purposes of E.O. 12866. The analysis meeting the requirements of the E.O. are found in the Regulatory Impact Review for the Emergency Rule.

7.11 Consolidated Appropriations Act, 2023

On December 29, 2022, President Biden signed H.R. 2617, the Consolidated Appropriations Act, 2023 ("Consolidated Appropriations Act") into law. The Consolidated Appropriations Act establishes that from December 29, 2022, through December 31, 2028, NMFS' September 17, 2021 rule amending the ALWTRP, Taking of Marine Mammals Incidental to Commercial Fishing Operations; Atlantic Large Whale Take Reduction Plan Regulations, 86 Fed. Reg. 51,970 (Sept. 17, 2021), "shall be deemed sufficient to ensure that the continued Federal and State authorizations of the American lobster and Jonah crab fisheries are in full compliance" with the MMPA and the Endangered Species (ESA) Act. H.R. 2617-1631–H.R. 2617-1632 (Division

JJ–North Atlantic Right Whales, Title I–North Atlantic Right Whales and Regulations).

However, the Consolidated Appropriations Act contains an exception at Section 101(b), stating that the preceding mandate does not apply to “an existing emergency rule, or any action taken to extend or make final an emergency rule” which is a specific reference to the subject of this EA, a rulemaking to extend the 2022 emergency rulemaking. H.R. 2617-1631–H.R. 2617-1632 (Division JJ–North Atlantic Right Whales, Title I–North Atlantic Right Whales and Regulations, Section 101 (b)).

7.12 National Environmental Policy Act

The National Environmental Policy Act (NEPA) provides a mechanism for identifying and evaluating the full spectrum of environmental issues associated with federal actions and for considering a reasonable range of alternatives to avoid or minimize adverse environmental impacts. The Council on Environmental Quality (CEQ) has issued regulations specifying the requirements for NEPA documents (40 CFR 1500–1508), as has NOAA in its policy and procedures for NEPA (NAO 216-6A). This EA is being prepared using the 2020 CEQ NEPA Regulations. The effective date of the 2020 CEQ NEPA Regulations was September 14, 2020, and reviews begun after this date are required to apply the 2020 regulations unless there is a clear and fundamental conflict with an applicable statute. 85 Fed. Reg. at 43372-73 (§§ 1506.13, 1507.3(a)).

7.12.1 *Environmental Assessment*

The required elements of an Environmental Assessment (EA) are specified in 40 CFR 1508.9(b). They are included in this document as follows:

- The need for this action is in Subsection 3.2;
- The alternatives that were considered are in Subsection 4.1, 4.2 and 4.3;
- The environmental impacts of the proposed action are in Subsection 6.2, 6.3, and 6.4;
- The agencies and persons consulted on this action are in Subsection 7.12.3; and
- A determination of significance is in the Finding of No Significance included with this EA.

While not required for the preparation of an EA, this document includes the following additional subsections that are based on requirements for an Environmental Impact Statement (EIS).

- Background and purpose are in Subsection 3.1 and 3.2;
- A description of the affected environment is in Subsection 5.1, 5.2, and 5.3;
- Cumulative effects of the proposed action are in Subsection 7.5;
- A list of preparers is in Subsection 7.12.4.

7.12.2 *Point of Contact*

For inquiries about the or to request a copy of the document, please contact the NMFS Greater Atlantic Region Protected Resources Division at (978) 281-9328.

7.12.3 *Agencies Consulted*

The following agencies, in alphabetical order, were consulted in preparing this document:

- Massachusetts Department of Marine Fisheries

7.12.4 *List of Preparers*

Michael J. Asaro
Economist
NMFS, NEFSC, Social Sciences Branch

Diane Borggaard
Right Whale Recovery Coordinator
NMFS, GARFO, Protected Resources Division

Tim Cardiasmenos
NMFS Greater Atlantic Region, NEPA Office

Colleen Coogan
Marine Mammal & Sea Turtle Branch Chief
NMFS, Greater Atlantic Region, Protected Resources Division

Cheryl Cross
Marine Resources Management Specialist
NMFS, Greater Atlantic Region, Protected Resources Division

Marianne Ferguson
NEPA Policy Analyst
NMFS, Greater Atlantic Region, NEPA Office

Jennifer Goebel
Marine Mammal Policy Analyst
NMFS, Greater Atlantic Region, Protected Resources Division

Allison Henry
Research Fishery Biologist
NMFS, NEFSC, Research Evaluation and Assessment Branch

Jean Higgins
Protected Species Conservation Branch Chief
NMFS, Greater Atlantic Region, Protected Resources Division

Alicia Miller
Research Fish Biologist
NMFS, NEFSC, Research Evaluation and Assessment Branch

Allison Murphy
Fishery Policy Analyst
NMFS, Greater Atlantic Region, Sustainable Fisheries Division

Alicia Schuler
Marine Resources Management Specialist
NMFS, Greater Atlantic Region, Protected Resources Division

Burton Shank
Research Fishery Biologist
NMFS, NEFSC, Research Evaluation and Assessment Branch

Kate Swails
Communications and Internal Affairs
NMFS, Greater Atlantic Region

Marisa Trego
Policy Analyst, ALWTRT Coordinator
NMFS, Greater Atlantic Region, Protected Resources Division

Integrated Statistics, Inc.

Crystal Franco
Environmental Specialist,
Integrated Statistics, Inc.

Chao Zou
Economist,
Integrated Statistics, Inc.

Laura Solinger
Quantitative Modeler,
Integrated Statistics, Inc.

Staff members of NMFS GARFO and NEFSC were also consulted in preparing this Environmental Assessment.

8 References

- Angliss, R.P., and D.P. DeMaster. 1998. Differentiating Serious and Non-Serious Injury of Marine Mammals Taken Incidental to Commercial Fishing Operation: Report of the Serious Injury Workshop. 1-2 April 1997, Silver Spring, Maryland. US Department of Commerce, NOAA Tech. Memo. NMFS-OPR-13, 48pp.
- Archer, F. I., R. L. Brownell, B. L. Hancock-Hanser, P. A. Morin, K. M. Robertson, K. K. Sherman, J. Calambokidis, J. Urbán R, P. E. Rosel, S. A. Mizroch, S. Panigada, and B. L. Taylor. 2019. Revision of fin whale *Balaenoptera physalus* (Linnaeus, 1758) subspecies using genetics. *Journal of Mammalogy* 100:1653-1670.
- ASMFC. 2015. American Lobster Stock Assessment Report for Peer Review. Atlantic States Marine Fisheries Commission, Stock Assessment Report. Atlantic States Marine Fisheries Commission.
- Baker, C. S., and P. J. Clapham. 2004. Modeling the past and future of whales and whaling. *Trends in Ecology and Evolution* 19:365–371.
- Barshaw, D. E. and K. L. Lavalli. 1988. Predation upon postlarval lobsters *Homarus americanus* by cunners *Tautoglabrus adspersus* and mud crabs *Neopanope sayi* on three different substrates: eelgrass, mud, and rock. *Marine Ecology Progress Series* 48:119-123.
- Baumgartner, M. 2021. Robots4Whales. Woods Hole Oceanographic Institution.
- Baumgartner, M. F., J. Bonnell, P. J. Corkeron, S.M. Van Parijs, C. Hotchkin, B. A. Hodges, J. Bort Thornton, B. L. Mensi, and S. M. Bruner. 2020. Slocum Gliders Provide Accurate Near Real-Time Estimates of Baleen Whale Presence From Human-Reviewed Passive Acoustic Detection Information. *Front. Mar. Sci.* 7:100.
- Baumgartner, M., and B. Mate. 2003. Summertime foraging ecology of North Atlantic right whales. *Marine Ecology Progress Series* 264:123-135.
- Baumgartner, M., F. Wenzel, N. Lysiak, and M. Patrician. 2017. North Atlantic right whale foraging ecology and its role in human-caused mortality. *Marine Ecology Progress Series* 581:165-181.
- Baumgartner, M., T. V. N. Cole, P. J. Clapham, and B. R. Mate. 2003. North Atlantic right whale habitat in the lower Bay of Fundy and on the SW Scotian Shelf during 1999-2001. *Marine Ecology Progress Series* 264:137-154.
- BEA [U.S. Bureau of Economic Analysis]. 2022. “Table 1.1.4. Price Indexes for Gross Domestic Product,” <https://apps.bea.gov/iTable/iTable.cfm?reqid=19&step=2#reqid=19&step=2&isuri=1&1921=survey> (accessed Feb 10, 2022).
- Black, K. P., and G. D. Parry. 1994. Sediment transport rates and sediment disturbance due to scallop dredging in Port Phillip Bay. *Mem. Queensl. Mus.* 36:327-341.

- Bologna, P. A., and R. S. Steneck. 1993. Kelp beds as habitat for American lobster *Homarus 78 americanus*. *Marine Ecology Progress Series* 100:127-134.
- Bourque, L., T. Wimmer, S. Lair, M. Jones, and P.-Y. Daoust. 2020. Incident Report: North Atlantic Right Whale Mortality Event in Eastern Canada, 2019 Collaborative Report Produced by: Canadian Wildlife Health Cooperative and Marine Animal Response Society. 210 pp.
- Cassoff, R., K. Moore, W. McLellan, S. Barco, D. Rotsteins, and M. Moore. 2011. Lethal entanglement in baleen whales. *Diseases of Aquatic Organisms*. 96, 175-85.
- CETAP [Cetacean and Turtle Assessment Program]. 1982. A characterization of marine mammals and turtles in the mid- and north Atlantic areas of the USA outer continental shelf. Final Report #AA551-CT8-48 Cetacean and Turtle Assessment Program, University of Rhode Island, Bureau of Land Management, Washington, D.C.
- Chuenpagdee, R., L. E. Morgan, S. M. Maxwell, E. A. Norse, and D. Pauly. 2003. Shifting gears: assessing collateral impacts of fishing methods in US waters. *Frontiers in Ecology and the Environment* 1:517-524.
- Churchill, J. H. 1989. The effect of commercial trawling on sediment resuspension and transport over the Middle Atlantic Bight continental shelf. *Continental Shelf Research* 9:841-864
- Clapham, P. J., L. S. Baraff, C. A. Carlson, M. A. Christian, D. K. Mattila, C. A. Mayo, M. A. Murphy, and S. Pittman. 1993. Seasonal occurrence and annual return of humpback whales, *Megaptera novaengliae*, in the southern Gulf of Maine. *Canadian Journal of Zoology*. 71(2): 440- 443.
- Clark, C. W., and G. C. Gagnon. 2002. Low-frequency vocal behaviors of baleen whales in the North Atlantic: Insights from IUSS detections, locations and tracking from 1992 to 1996. *J. Underwater Acoust. (US Navy)*, 52(3):609-640.
- Clay, P. M., and J. Olson. 2008. Defining ‘Fishing Communities’: Vulnerability and the Magnuson-Stevens Fishery Conservation and Management Act. *Human Ecology Review* 15(2): 143–60.
- Coen, L. D. 1995. A review of the potential impacts of mechanical harvesting on subtidal and intertidal shellfish resources. South Carolina Department of Natural Resources, Marine Resources Research Institute.
- Colburn, L. L., and M. Jepson. 2012. Social Indicators of Gentrification Pressure in Fishing Communities: A Context for Social Impact Assessment. *Coastal Management*. 40(3): 289–300.
- Conn, P. B., and G. K. Silber. 2013. Vessel speed restrictions reduce risk of collision-related mortality for North Atlantic right whales. *Ecosphere* 4(4):43.
- Corkeron, P., P. Hamilton, J. Bannister, P. Best, C. Charlton, K. R. Groch, K. Findlay, V. Rowntree, E. Vermeulen, and R. M. Pace. 2018. The recovery of North Atlantic right

- whales, *Eubalaena glacialis*, has been constrained by human-caused mortality. *R. Soc. Open Sci.* 5:180892.
- CZM [Massachusetts Office of Coastal Zone Management]. 1999. Offshore Bathymetry. 1:250,000. <https://www.mass.gov/info-details/massgis-data-offshore-bathymetry-1250000> (accessed January 3, 2023).
- Daoust, P.-Y., É. L. Couture, T. Wimmer, and L. Bourque. 2018. Incident report: North Atlantic right whale mortality event in the Gulf of St. Lawrence, 2017. Report, Canadian Wildlife Health Cooperative, Marine Animal Response Society, and Fisheries and Oceans Canada, Ottawa, Canada.
- Davies, K., M. Brown, P. Hamilton, A. Knowlton, C. Taggart, and A. Vanderlaan. 2019. Variation in North Atlantic right whale *Eubalaena glacialis* occurrence in the Bay of Fundy, Canada, over three decades. *Endangered Species Research* 39:159-171.
- Davis, G. E., M. F. Baumgartner, J. M. Bonnell, J. Bell, C. Berchok, J. Bort Thornton, S. Brault, G. Buchanan, R. A. Charif, D. Cholewiak, C. W. Clark, P. Corkeron, J. Delarue, K. Dudzinski, L. Hatch, J. Hildebrand, L. Hodge, H. Klinck, S. Kraus, B. Martin, D. K. Mellinger, H. MoorsMurphy, S. Nieukirk, D. P. Nowacek, S. Parks, A. J. Read, A. N. Rice, D. Risch, A. Sirovic, M. Soldevilla, K. Stafford, J. E. Stanistreet, E. Summers, S. Todd, A. Warde, and S. M. Van Parijs. 2017. Long-term passive acoustic recordings track the changing distribution of North Atlantic right whales (*Eubalaena glacialis*) from 2004 to 2014. *Sci Rep* 7:13460.
- Davis, G. E., M. F. Baumgartner, P. J. Corkeron, J. Bell, C. Berchok, J. M. Bonnell, J. B. Thornton, S. Brault, G. A. Buchanan, D. M. Cholewiak, C. W. Clark, J. Delarue, L. T. Hatch, H. Klinck, S. D. Kraus, B. Martin, D. K. Mellinger, H. Moors-Murphy, S. Nieukirk, D. P. Nowacek, S. E. Parks, D. Parry, N. Pegg, A. J. Read, A. N. Rice, D. Risch, A. Scott, M. S. Soldevilla, K. M. Stafford, J. E. Stanistreet, E. Summers, S. Todd, and S. M. van Parijs. 2020. Exploring movement patterns and changing distributions of baleen whales in the western North Atlantic using a decade of passive acoustic data. *Glob Change Biol.* 00:1-29.
- Dombroski, J., S. Parks, and D. Nowacek. 2021. Dive behavior of North Atlantic right whales on the calving ground in the Southeast USA: Implications for conservation. *Endangered Species Research*, 46: 35–48.
- Donovan, G. P. 1991. A review of IWC stock boundaries. *Rep. int. Whal. Common* (special issue).13:39-68.
- Edwards, E. F., C. Hall, T. J. Moore, C. Sheredy, and J. Redfern. 2015. Global distribution of fin whales (*Balaenoptera physalus*) in the post-whaling era (1980 to 2012). *Mamm Rev.* 45:197– 214.
- Eno, N. C., D. S. MacDonald, J. A. M. Kinnear, S. C. Amos, C. J. Chapham, R. A. Clard, F. P. D. Bunker, and C. Munro. 2001. Effects of crustacean traps on benthic fauna. *ICES Journal of Marine Science* 58:11-20.

- Fortune, S., A. Trites, C. Mayo, D. Rosen, and P. Hamilton. 2013. Energetic requirements of North Atlantic right whales and the implications for species recovery. *Marine Ecology Progress Series*. 478:253-272.
- Gallopín, Gilberto C. 2006. Linkages between Vulnerability, Resilience, and Adaptive Capacity. *Global Environmental Change, Resilience, Vulnerability, and Adaptation: A Cross-Cutting Theme of the International Human Dimensions Programme on Global Environmental Change*. 16(3): 293–303.
- Goode, A. G., J. H. Grabowski, and D. C. Brady. 2021. Evaluating benthic impact of the Gulf of Maine lobster fishery using the Swept Area Seabed Impact (SASI) model. *Canadian Journal of Fisheries and Aquatic Sciences* 78(6): 693-703.
- Grabowski, J. H., E. J. Clesceri, J. Gaudette, A. Baukus, M. Weber, and P. O. Yund. 2010. Use of herring bait to farm lobster in the Gulf of Maine. *PLoS One* 5: e10188.
- Grieve, B. D., J. A. Hare, and V. S. Saba. 2017. Projecting the effects of climate change on *Calanus finmarchicus* distribution within the U.S. Northeast Continental Shelf. *Scientific Reports* 7:6264.
- Hain, J. H. W., M. A. M. Hyman, R. D. Kenney, and H. E. Winn. 1985. The role of cetaceans in the shelf edge region of the northeastern United States. *Marine Fisheries Review* 47:13-17.
- Hain, J. H. W., M. Ratnaswamy, R. D. Kenney, and H. E. Winn. 1992. The fin whale, *Balaenoptera physalus*, in waters of the northeastern United States continental shelf. Report of the International Whaling Commission. 42:653-669.
- Hall, S. J. 1999. *The Effects of Fishing on Marine Ecosystems and Communities*. Blackwell Science, Oxford.
- Hall-Arber, M., C. Dyer, J. Poggie, J. McNally, and R. Gagne. 2001. New England's Fishing Communities. MITSG 01-15, MIT Sea Grant College Program, 292 Main Street, E38-300, Cambridge, MA 02139.
- Hamilton, P. K., A. R. Knowlton, M. N. Hagbloom, K. R. Howe, H. M. Pettis, M. K. Marx, M. A. Zani, and S. D. Kraus. 2019. Maintenance of the North Atlantic right whale catalog, whale scarring and visual health databases, anthropogenic injury case studies, and near real-time matching for biopsy efforts, entangled, injured, sick or dead right whales. New England Aquarium, Boston, MA. Contract No. 1305M2-18-P-NFFM-0108.
- Hamilton, P., and S. Kraus. 2019. Frequent encounters with the seafloor increase right whales' risk of entanglement in fishing groundlines. *Endangered Species Research* 39:235-246.
- Hartley, D., A. Whittingham, J. Kenney, T. Cole, and E. Pomfret. 2003. Large Whale Entanglement Report 2001. Report to the National Marine Fisheries Service, updated February 2003, 61 pp.
- Hayes, S. A., E. Josephson, K. Maze-Foley, and P. E. Rosel. 2020. US Atlantic and Gulf of Mexico Marine Mammal Stock Assessments - 2019. Page 479.

- Hayes, S. A., E. Josephson, K. Maze-Foley, P. E. Rosel and J. Wallace. 2022. US Atlantic and Gulf of Mexico Marine Mammal Stock Assessments - 2020. Page 386.
- Hayes, S. A., E. Josephson, K. Maze-Foley, P. E. Rosel, and J. Wallace. 2023. Draft US Atlantic and Gulf of Mexico Marine Mammal Stock Assessments - 2022. Northeast Fisheries Science Center Woods Hole, MA.
- Hayes, S. A., E. Josephson, K. Maze-Foley, P. E. Rosel, and J.T. Turek. 2021. US Atlantic and Gulf of Mexico Marine Mammal Stock Assessments - 2020. Page 403.
- Henry, A. G., T. V. N. Cole, L. Hall, W. Ledwell, D. Morin, and A. Reid. 2016. Serious injury and mortality determinations for baleen whale stocks along the Gulf of Mexico, United States East Coast and Atlantic Canadian Provinces, 2010-2014. Page 51.
- Henry, A.G., M. Garron, D. Morin, Smith, A., A. Reid, W. Ledwell, and T. Cole. 2022. Serious Injury and Mortality Determinations for Baleen Whale Stocks along the Gulf of Mexico, United States East Coast, and Atlantic Canadian Provinces, 2015-2019. Page 65. NOAA Tech Memo. NMFS-NE-280.
- ICES. 1992. Report of the Study Group on Ecosystem Effects of Fishing Activities, Copenhagen, 7-14 April. ICES CM 1992/G:11, International Council for the Exploration of the Sea, Study Group on Ecosystem Effects of Fishing Activities, Copenhagen.
- Irvine, L. G., M. Thums, C. E. Hanson, C. R. McMahon, and M. A. Hindell. 2017. Quantifying the energy stores of capital breeding humpback whales and income breeding sperm whales using historical whaling records. *R. Soc. open sci.* 4:160290.
- Jacob, S. P. Weeks, B. Blount, M. Jepson. 2010. Exploring Fishing Dependence in Gulf Coast communities. *Marine Policy.* 34(6):1307-1314.
- Johnson, A., G. Salvador, J. Kenney, J. Robbins, S. Kraus, S. Landry, and P. Clapham. 2005. Fishing Gear Involved in Entanglements of Right and Humpback Whales. *Marine Mammal Science.* 21: 635-645.
- Kaiser, M. J. 2000. The implications of the effects of fishing on non-target species and habitats. Pages 383-392 in M. J. Kaiser and S. J. de Groot, editors. *The Effects of Fishing on Non-target Species and Habitats.* Blackwell Science.
- Kenney, R. D. 2001. Anomalous 1992 spring and summer right whale (*Eubalaena glacialis*) distributions in the Gulf of Maine. *Journal of Cetacean Research and Management (Special Issue)* 2:209-223.
- Klanjscek, T., R. M. Nisbet, H. Caswell, and M. G. Neubert. 2007. A model for energetics and bioaccumulation in marine mammals with applications to the right whale. *Ecological Applications.* 17:2233-2250.
- Knowlton, A. and S. Kraus. 2001. Mortality and serious injury of northern right whales (*Eubalaena glacialis*) in the western North Atlantic Ocean. *Journal of Cetacean Research and Management (Special Issue).* 2.

- Knowlton, A. R., P. K. Hamilton, M. K. Marx, H. M. Pettis, and S. D. Kraus. 2012. Monitoring North Atlantic right whale *Eubalaena glacialis* entanglement rates: a 30 yr retrospective. *Mar Ecol Prog Ser.* 466:293-302.
- Kraus, S. D., and R. M. Rolland. 2007. Right whales in the urban ocean. Pages 1-38 in S. D. Kraus and R. M. Rolland, editors. *The Urban Whale: North Atlantic Right Whales at the Crossroads.* Harvard University Press, Cambridge.
- Kraus, S. D., R. D. Kenney, C. A. Mayo, W. A. McLellan, M. J. Moore, and D. P. Nowacek. 2016. Recent Scientific Publications Cast Doubt on North Atlantic Right Whale Future. *Frontiers in Marine Science.* 3.
- Krzystan, A., T. Gowan, W. Kendall, J. Martin, J. Ortega-Ortiz, K. Jackson, A. Knowlton, P. Naessig, M. Zani, D. Schulte, and C. Taylor. 2018. Characterizing residence patterns of North Atlantic right whales in the southeastern USA with a multistate open robust design model. *Endangered Species Research.* 36:279-295.
- Laist, D. W., A. R. Knowlton, J. G. Mead, A. S. Collet, and M. Podesta. 2001. Collisions between ships and whales. *Marine Mammal Science.* 17:35-75.
- Lawton, P., and K. L. Lavalli. 1995. Post larval, juvenile, adolescent, and adult ecology. Pages 47-88 in J. R. Factor, editor. *Biology of the Lobster Homarus americanus.* Academic Press, New York, NY.
- Levin, L. A. 1984. Life history and dispersal patterns in a dense infaunal polychaete assemblage: community structure and response to disturbance. *Ecology* 65:1185-1200. Lincoln, D. 1998. Lobsters on the edge-essential lobster habitats in New England. Greenlite Consultants, Newton Highlands, MA.
- Lincoln, D. 1998. Lobsters on the edge-essential lobster habitats in New England. Greenlite Consultants, Newton Highlands, MA.
- Mayer, L. M., D. F. Schick, R. H. Findlay, and D. L. Rice. 1991. Effects of commercial dragging on sedimentary organic matter. *Marine Environmental Research.* 31:249-261.
- Mayo, C. A., and M. K. Marx. 1990. Surface behavior of the North Atlantic right whale, *Eubalaena glacialis*, and associated zooplankton characteristics. *Canadian Journal of Zoology.* 68:2
- Mayo, C. A., L. Ganley, C. A. Hudak, S. Brault, M. K. Marx, E. Burke, and M. W. Brown. 2018. Distribution, demography, and behavior of North Atlantic right whales (*Eubalaena glacialis*) in Cape Cod Bay, Massachusetts, 1998-2013: Right Whales in Cape Cod Bay. *Marine Mammal Science.* 34:979-996.
- McCauley, R., R. Day, K. Swadling, Q. Fitzgibbon, R. Watson, and J. Semmens. 2017. Widely used marine seismic survey air gun operations negatively impact zooplankton. *Nature Ecology and Evolution* 1. 214-2220.

- Messieh, S. N., T.W. Rowell, D. L. Peer, and P. J. Cranford. 1991. The effects of trawling, dredging and ocean dumping on the eastern Canadian continental shelf seabed. *Continental Shelf Research*. 11(8–10): 1237-1263.
- Meyer-Gutbrod, E. L., and C. H. Greene. 2018. Uncertain recovery of the North Atlantic right whale in a changing ocean. *Global Change Biology* 24:455-464.
- Meyer-Gutbrod, E. L., C. H. Greene, P. J. Sullivan, and A. J. Pershing. 2015. Climate-associated changes in prey availability drive reproductive dynamics of the North Atlantic right whale population. *Marine Ecology Progress Series*. 535:243-258.
- Mitchell, E., and D. G. Chapman. 1977. Preliminary assessment of stocks of northwest Atlantic sei whales (*Balaenoptera borealis*). Rep. omt/ Whal. Common. Special Issue 1: 117-120.
- Monsarrat, S., M. G. Pennino, T. D. Smith, R. R. Reeves, C. N. Meynard, D. M. Kaplan, and A. S. Rodrigues. 2016. A spatially explicit estimate of the prewhaling abundance of the endangered North Atlantic right whale. *Conserv Biol*. 30:783-791.
- Moore, M. J., and J. M. van der Hoop. 2012. The Painful Side of Trap and Fixed Net Fisheries: Chronic Entanglement of Large Whales. *Journal of Marine Biology*. 2012:1-4.
- Morano, J. L., A. N. Rice, J. T. Tielens, B. J. Estabrook, A. Murray, B. L. Roberts, and C. W. Clark. 2012. Acoustically Detected Year-Round Presence of Right Whales in an Urbanized Migration Corridor: Right Whales in Massachusetts Bay. *Conservation Biology* 26:698-707.
- Murphy, M. A. 1995. Occurrence and group characteristics of minke whales, *Balaenoptera acutorostrata*, in Massachusetts Bay and Cape Cod Bay. *Oceanographic Literature Review*. 5(43): 506.
- Murray, A., A. Rice, and C. Clark. 2013. Extended seasonal occurrence of humpback whales in Massachusetts Bay. *Journal of the Marine Biological Association of the U.K.* 94(6): 1117-1125.
- NMFS. 2002. Workshop on the effects of fishing gear on marine habitats off the Northeastern United States October 23-25, 2001. Boston, Massachusetts. National Marine Fisheries Service, Northeast Fisheries Science Center, Woods Hole, Massachusetts.
- NMFS. 2004. Characterization of the Fishing Practices and Marine Benthic Ecosystems of the Northeast U.S. Shelf, and an Evaluation of the Potential Effects of Fishing on Essential Fish Habitat.
- NMFS. 2013. North Atlantic Right Whale (*Eubalaena glacialis*) Source Document for the Critical Habitat Designation: A review of information pertaining to the definition of “critical habitat”. December 2012. 166pp.
- NMFS. 2014. NMFS-Greater Atlantic Region (GARFO). Memo to the record: Determination regarding reinitiation of Endangered Species Act section 7 consultation on 12 GARFO fisheries and two Northeast Fisheries Science Center funded fisheries research surveys

- due to critical habitat designation for loggerhead sea turtles. Memo issued September 17, 2014.
- NMFS. 2015a. Endangered Species Act Section 4(b)(2) Report: Critical Habitat for the North Atlantic Right Whale (*Eubalaena glacialis*). Prepared by National Marine Fisheries Service Greater Atlantic Regional Fisheries Office and Southeast Regional Office, December 2015.
http://www.greateratlantic.fisheries.noaa.gov/regs/2016/January/16narwchsection4_b__2_report_012616.pdf
- NMFS. 2015b. North Atlantic Right Whale (*Eubalaena glacialis*). Source Document for the Critical Habitat Designation: A review of information pertaining to the definition of “critical habitat” Prepared by National Marine Fisheries Service Greater Atlantic Regional Fisheries Office and Southeast Regional Office, July 2015.
- NMFS. 2017. 62nd Northeast Regional Stock Assessment Workshop (62nd SAW) Assessment Summary Report. US Department of Commerce, Northeast Fisheries Science Center, Woods Hole, MA.
- NMFS. 2019. Guidance on the Application of NEPA for Emergency Response Actions. Memo to National Environmental Policy Act (NEPA) Practitioners. September 23, 2019.
- NMFS. 2021. Final Environmental Impact Statement (FEIS), Regulatory Impact Review, and Final Regulatory Flexibility Analysis for Amending the Atlantic Large Whale Take Reduction Plan: Risk Reduction Rule Volume I.
https://www.greateratlantic.fisheries.noaa.gov/public/nema/aprd/2021FEIS_Volume%20I.pdf
- NMFS. 2022. Environmental Assessment, Finding of No Significance, and Regulatory Impact Review for the 2022 Emergency Final Rule to Reduce Right Whale Interactions with Lobster and Jonah Crab Trap/Pot Gear. <https://media.fisheries.noaa.gov/2022-03/Environmental%20Assessment%20March%202022.pdf>
- Oedekoven, C., E. Fleishman, P. Hamilton, J. Clark, and R. Schick. 2015. Expert elicitation of seasonal abundance of North Atlantic right whales *Eubalaena glacialis* in the mid-Atlantic. *Endangered Species Research*. 29:51-58.
- Pace, R. M. 2021. Revisions and Further Evaluations of the Right Whale Abundance Model: Improvements for Hypothesis Testing. Page 54. NOAA Tech. Memo. NOAA, Northeast Fisheries Science Center, Woods Hole, MA.
- Pace, R.M., III, P. J. Corkeron, and S. D. Kraus. 2017. State-space mark-recapture estimates reveal a recent decline in abundance of North Atlantic right whales. *Ecol. and Evol.* 7:8730-8741.
- PACM [Passive Acoustic Cetacean Map]. 2023. Woods Hole (MA): NOAA Northeast Fisheries Science Center v1.1.3 [access January 23, 2023]. <https://apps-nefsc.fisheries.noaa.gov/pacm>

- Paquet, D., C. Haycock, and H. Whitehead. 1997. Numbers and seasonal occurrence of humpback whales, *Megaptera novaeangliae*, off Brier Island, Nova Scotia. *Canadian Field Naturalist*. 111:548–552.
- Payne, M. P., D. N. Wiley, S. B. Young, S. Pittman, P. J. Clapham, and J. W. Jossi. 1990. Recent fluctuations in the abundance of baleen whales in the southern Gulf of Maine in relation to changes in selected prey. *Fisheries Bulletin*. 88:687-696.
- Payne, P. M., J. R. Nicolas, L. O'Brien, and K. D. Powers. 1986. The distribution of the humpback whale, *Megaptera novaeangliae*, on Georges Bank and in the Gulf of Maine in relation to densities of the sand eel, *Ammodytes americanus*. *Fishery Bulletin*. 84:271-227.
- Perry, S. L., D. P. DeMaster, and G. K. Silber. 1999. The Great Whales: History and Status of Six Species Listed as Endangered Under the U.S. Endangered Species Act of 1973. *The Marine Fisheries Review*. 61:74.
- Pettis, H. M., Pace, R. M. III, and P. K. Hamilton. 2022. North Atlantic Right Whale Consortium 2021 Annual Report Card.
https://www.narwc.org/uploads/1/1/6/6/116623219/2021report_cardfinal.pdf
- Pilskaln, C. H., J. H. Churchill, and L. M. Mayer. 1998. Resuspension of sediment by bottom trawling in the Gulf of Maine and potential geochemical consequences. *Conservation Biology*. 12:1223-1229.
- Plourde, S., C. Lehoux, C. L. Johnson, G. Perrin, and V. Lesage. 2019. North Atlantic right whale (*Eubalaena glacialis*) and its food: (I) a spatial climatology of *Calanus* biomass and potential foraging habitats in Canadian waters. 00:19.
- Prieto, R., D. Janiger, M. A. Silva, G. T. Waring, and J. M. Gonçalves. 2012. The forgotten whale: A bibliometric analysis and literature review of the North Atlantic sei whale *Balaenoptera borealis*: North Atlantic sei whale review. *Mammal Review*. 42(3): 235–272.
- Quintana-Rizzo E., S. Leiter, T. V. N. Cole, M. N. Hagbloom, A. R. Knowton, P. Nagelkirk, O.O'Brian, C. B. Khan, A. G. Henry, P. A. Duley, L. M. Crowe, C. A. Mayo, and S. D. Kraus. 2021. Residency, demographics, and movement patterns of North Atlantic right whales *Eubalaena glacialis* in an offshore wind energy development area in southern New England, USA. *Endang Species Res*. 45:251-268.
- Record, N. R., J. Runge, D. Pendleton, W. Balch, K. Davies, A. Pershing, C. Johnson, K. Stamieszkin, R. Ji, Z. Feng, S. Kraus, R. Kenney, C. Hudak, C. Mayo, C. Chen, J. Salisbury, and C. Thompson. 2019b. Rapid Climate-Driven Circulation Changes Threaten Conservation of Endangered North Atlantic Right Whales. *Oceanography*. 32.
- Record, N. R., W. M. Balch, and K. Stamieszkin. 2019a. Century-scale changes in phytoplankton phenology in the Gulf of Maine. *PeerJ*. 7:e6735.

- Risch, D., C. W. Clark, P. J. Dugan, M. Popescu, U. Siebert, S. M. Van Parijs. 2013. Minke whale acoustic behavior and multi-year seasonal and diel vocalization patterns in Massachusetts Bay, USA. *Marine Ecology Progress Series*. 489:279-295.
- Risch, D., M. Castellote, C. W. Clark, G. E. Davis, P. J. Dugan, L. E. W. Hodge, A. Kumar, K. Lucke, D. K. Mellinger, S. L. Niekirk, C. M. Popescu, C. Ramp, A. J. Read, A. N. Rice, M. A. Silva, U. Siebert, K. M. Stafford, H. Verdaat, and S.M. Van Parijs. 2014. Seasonal migrations of North Atlantic minke whales: novel insights from large-scale passive acoustic monitoring networks. *Mov Ecol*. 2, 24.
- Robbins, J. J., S. Landry, and D. K. Mattila. 2009. Estimating entanglement mortality from scar based studies. International Whaling Commission Scientific Committee, Madeira, Portugal.
- Roberts J. J., L. Mannocci, P. N. Halpin. 2017. Final Project Report: Marine Species Density Data Gap Assessments and Update for the AFTT Study Area, 2016-2017 (Opt. Year 1). Document version 1.4. Report prepared for Naval Facilities Engineering Command, Atlantic by the Duke University Marine Geospatial Ecology Lab, Durham, NC.
- Roberts J. J., R. S. Schick, P. N. Halpin. 2020. Final Project Report: Marine Species Density Data Gap Assessments and Update for the AFTT Study Area, 2018-2020 (Option Year 3). Document version 1.4. Report prepared for Naval Facilities Engineering Command, Atlantic by the Duke University Marine Geospatial Ecology Lab, Durham, NC.
- Roberts, J. J. 2022. Right Whale Model Webinar and Q&A for Take Reduction Team [Webinar]. NMFS. <https://register.gotowebinar.com/register/3942817237491823632>.
- Roberts, J. J., B. D. Best, L. Mannocci, E. Fujioka, P. N. Halpin, D. L. Palka, L. P. Garrison, K. D. Mullin, T. V. N. Cole, C. B. Khan, W. A. McLellan, D. A. Pabst, and G. G. Lockhart. 2016. Habitat-based cetacean density models for the U.S. Atlantic and Gulf of Mexico. *Scientific Reports* 6:22615.
- Rogers, S. I., M. J. Kaiser, and S. Jennings. 1998. Ecosystem effects of demersal fishing: a European perspective. Pages 68-78 in E. D. Dorsey and J. Pederson, editors. *Effect of Fishing Gear on the Sea Floor of New England*. Conservation Law Foundation, Boston, Massachusetts.
- Rumohr, H., and P. Krost. 1991. Experimental evidence of damage to benthos by bottom trawling with special reference to *Arctica islandica*. *Meeresforschung*. 33:340-345.
- Schilling M. R., I. Seipt, M. T. Weinrich, S. E. Frohock, A. E. Kuhlberg, and P. J. Clapham. 1992. Behavior of individually identified sei whales, *Balaenoptera borealis*, during an episodic influx into the southern Gulf of Maine in 1986. *Fishery Bulletin*. 90: 749–755.
- Schweitzer, C. C., R. N. Lipcius, and G. S. Bradley. 2018. Impacts of a multi-trap line on benthic habitat containing emergent epifauna within the Mid-Atlantic Bight. *ICES Journal of Marine Science*. 75(6): 2202–2212.

- Sharp, S., W. McLellan, D. Rotstein, A. Costidis, S. Barco, K. Durham, T. Pitchford, K. Jackson, P. Daoust, T. Wimmer, E. Couture, L. Bourque, T. Frasier, B. Frasier, D. Fauquier, T. Rowles, P. Hamilton, H. Pettis, and M. Moore. 2019. Gross and histopathologic diagnoses from North Atlantic right whale *Eubalaena glacialis* mortalities between 2003 and 2018. *Diseases of Aquatic Organisms*. 135:1-31.
- Sorochan, K. A., S. Plourde, R. Morse, P. Pepin, J. Runge, C. Thompson and C. L. Johnson. 2019. North Atlantic right whale (*Eubalaena glacialis*) and its food: (II) interannual variations in biomass of *Calanus* spp. on western North Atlantic shelves. *Journal of Plankton Research*. 41(5): 687–708.
- Steneck, R. S., and C. Wilson. 1998. Why are there so many lobsters in Penobscot Bay? Pages 72-75 in D. D. Platt, editor. *Rim of the Gulf – Restoring Estuaries in the Gulf of Maine*. The Island Institute, Rockland, ME.
- Stephenson, F., A. C. Mill, C. L. Scott, N. V. C. Polunin, and C. Fitzsimmons. 2017. Experimental potting impacts on common UK reef habitats in areas of high and low fishing pressure. *ICES Journal of Marine Science*. 74:1648-1659.
- Stone, K. M., S. M. Leiter, R. D. Kenney, B. C. Wikgren, J. L. Thompson, J. K. D. Taylor, and S. D. Kraus. 2017. Distribution and abundance of cetaceans in a wind energy development area offshore of Massachusetts and Rhode Island. *Journal of Coastal Conservation*. 21:527-543.
- Swingle, W. M., S. G. Barco, T. D. Pitchford, W. A. McLellan, and D. A. Pabst. 1993. Appearance of Juvenile Humpback Whales Feeding in the Nearshore Waters of Virginia. *Marine Mammal Science*. 9: 309-315.
- Uhrin, A.V. 2016. Tropical cyclones, derelict traps, and the future of the Florida Keys commercial spiny lobster fishery. *Marine Policy*. 69: 84-91.
- van Der Hoop, J. M., M. J. Moore, S. G. Barco, T. V. N. Cole, P.-Y. Daoust, A. G. Henry, D. F. McAlpine, W. A. McLellan, T. Wimmer, and A. R. Solow. 2013. Assessment of management to mitigate anthropogenic effects on large whales. *Cons. Biol*. 27:121-133.
- van der Hoop, J. M., P. Corkeron, J. Kenney, S. Landry, D. Morin, J. Smith, and M. J. Moore. 2016. Drag from fishing gear entangling North Atlantic right whales. *Marine Mammal Science*. 32:619-642.
- van der Hoop, J.M., P. Corkeron, and M. Moore. 2017. Entanglement is a costly life-history stage in large whales. *Ecol Evol*. 7:92-106.
- Waring, G. T., E. Josephson, K. Maze-Foley, and P.E. Rosel eds. 2009. *U.S. Atlantic and Gulf of Mexico Marine Mammal Stock Assessments – 2009*. NOAA Tech. Memo. NMFS NE 213, Woods Hole, Massachusetts, USA.
- Watkins, W. A., and W. E. Schevill. 1976. Right whale feeding and baleen rattle. *Journal of Mammalogy*. 57:58-66.

- Watkins, W.A., P. Tyack, K. E. Moore and J. E. Bird. 1987. The 20-Hz signals of finback whales (*Balaenoptera physalus*). *J. Acoust. Soc. Am.* 82: 901-1912.
- Weinrich, M. T., R. D. Kenney, and P. K. Hamilton. 2000. Right whales (*Eubalaena glacialis*) on Jeffreys Ledge: A Habitat of Unrecognized Importance?. *Marine Mammal Science*. 16: 326-337.
- Whittingham, A., M. Garron, J. Kenney, and D. Hartley. 2005. Large Whale Entanglement Report 2003. Report to the National Marine Fisheries Service, updated June 2005.
- Wikgren, B., H. Kite-Powell, and S. Kraus. 2014. Modeling the distribution of the North Atlantic right whale *Eubalaena glacialis* off coastal Maine by aerial co-kriging. *Endangered Species Research*. 24:21-31.
- Wiles, G. 2017. Periodic status review for blue, fin, sei, North Pacific right, and sperm whales in Washington.
- Wiley, D., R. A. Asmutis, T. D. Pitchford, and D. P. Gannon. 1995. Stranding and mortality of humpback whales, *Megaptera novaeangliae*, in the mid-Atlantic and southeast United States, 1985-1992. *Fishery Bulletin- National Oceanic and Atmospheric Administration*. 93: 196-205.
- Williams, R., G. A. Vikingsson, A. Gislason, C. Lockyer, L. New, L. Thomas, and P. S. Hammond. 2013. Evidence for density-dependent changes in body condition and pregnancy rate of North Atlantic fin whales over four decades of varying environmental conditions. *ICES Journal of Marine Science*. 70:1273-1280.
- Wishner, K. F., E. Durbin, A. Durbin, M. Macaulay, H. Winn, and R. Kenney. 1988. Copepod patches and right whales in the Great South Channel off New England. *Bulletin of Marine Science*. 43:825-844.
- Wishner, K. F., J. R. Schoenherr, R. Beardsley, and C. Chen. 1995. Abundance, distribution and population structure of the copepod *Calanus finmarchicus* in a springtime right whale feeding area in the southwestern Gulf of Maine. *Continental Shelf Research*. 15:475-507.
- Woodley, T. H., and D. E. Gaskin. 1996. Environmental characteristics of North Atlantic right and fin whale habitat in the lower Bay of Fundy, Canada. *Canadian Journal of Zoology*. 74:75.

**ENVIRONMENTAL ASSESSMENT OF A 2023 EMERGENCY RULE TO REDUCE
RIGHT WHALE INTERACTIONS WITH TRAP/POT GEAR
VOLUME II**

**January 2023
Updated March 31, 2023 (See Errata, Appendix A)**

**US DEPARTMENT OF COMMERCE
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
NATIONAL MARINE FISHERIES SERVICE
GREATER ATLANTIC REGIONAL FISHERIES OFFICE**

Contents

GREATER ATLANTIC REGIONAL FISHERIES OFFICE	i
Figure Correction Appendices.....	1
<i>Appendix A Errata to the Environmental Assessment.....</i>	<i>1</i>
Chapter 1 Appendices	4
<i>Appendix 1.1 Letters of Concern</i>	<i>4</i>
Chapter 3 Appendices	49
<i>Appendix 3.1 Community Profiles</i>	<i>49</i>
Chapter 4 Appendices	86
<i>Appendix 4.1 Decision Support Tool Model Runs.....</i>	<i>86</i>
Appendix 4.1.1 Baseline Information.....	86
Appendix 4.1.2 Alternative 2 Model Runs	104
Appendix 4.1.3 Alternative 3 Model Runs	116

Figure Correction Appendices

Appendix A Errata to the Environmental Assessment

1. In February 2023, NMFS published an Environmental Assessment (EA) for the 2023 Emergency Rule to Reduce Right Whale Interactions with Trap/Pot Gear (88 FR 7362; Published February 3, 2023; hereafter referred to as the 2023 Emergency Rule).

Within the EA for the 2023 Emergency Rule, NMFS presented several figures intended to represent North Atlantic right whale visual observations collected February-April 2018-2022 within the action area of the emergency rule. As outlined in the figure descriptions, the figures represent North Atlantic right whale visual sightings from the Center for Coastal Studies (CCS) aerial surveys, sightings by the Northeast Fisheries Science Center's (NEFSC) dedicated aerial and shipboard surveys, and opportunistic sightings collected by NEFSC.


Upon further internal review, NMFS realized we inadvertently included dedicated aerial and shipboard surveys and opportunistic sightings collected during May 2018 and May 2019. Additionally, the figures within the 2023 Emergency Rule EA did not include dedicated surveys conducted by the CCS in April 2021.

Within the MRA Wedge area, the original figures inadvertently included 4 sighting location points (representing 6 right whales) from May 2018 and May 2019 observations. The original figures also inadvertently excluded 18 sighting location points (representing 23 right whales) from April 2021 observations (See Correction Figure A).


This correction applies to three figures within the EA, Figure 2 (Chapter 3), Figure 3 (Chapter 2), and Figure 9 (Chapter 6). All of the figures have been corrected in this version (dated March 31 2023) to remove visual sightings from May 2018 and 2019 and updated to include CCS aerial survey visual sightings collected in April 2021 (See Correction Figure B).

The error in the original figures do not change any conclusions within the 2023 EA analyses, nor does it alter the Finding of No Significant Impact (FONSI) determination. The figures were used for qualitative representations of empirical sightings data, not quantitative analysis of risk to right whales. The empirical data differs from the systematic survey data incorporated into the habitat density model that is used in the Large Whale Decision Support Tool to estimate relative risk and risk reduction.

Correction Figure A: Below is an illustrative map showing the differences between the original published Figure 3 and the updated Figure 3 in this correction. The original Figure 3 of the Environmental Assessment (EA) for the 2023 Emergency Rule to Reduce Right Whale Interactions with Trap/Pot Gear incorrectly includes North Atlantic right whale visual sightings from opportunistic and dedicated surveys conducted May 2018 and May 2019 (represented by hollow circles). Additionally, the original Figure 3 is missing North Atlantic right whale visual sightings from Center for Coastal Studies (CCS) aerial surveys conducted April 2021 (represented by yellow crosses). Massachusetts Restricted Area is shaded in dark gray, and MRA Wedge is represented by the shaded white area.



Correction Figure B: Below is a figure showing the side by side comparison between the original published Figure 3 (left panel) and the updated Figure 3 (right panel) in this correction. The left panel shows the original Figure 3 published in the Environmental Assessment (EA) for the 2023 Emergency Rule. This figure inadvertently includes North Atlantic right whale visual sightings from opportunistic and dedicated surveys conducted May 2018 and May 2019. Additionally, the original Figure 3 is missing North Atlantic right whale visual sightings from Center for Coastal Studies (CCS) aerial surveys conducted April 2021. The panel on the right, the updated Figure 3, shows the intended representations of North Atlantic right whale visual observations collected February-April 2018-2022 within the action area of the 2023 Emergency Rule. In both panels, Massachusetts Restricted Area is shaded in dark gray, and MRA Wedge is represented by the shade white area.



Chapter 1 Appendices

Appendix 1.1 Letters of Concern

Following are three letters NMFS received regarding the overlap of North Atlantic right whales and fishing gear in unrestricted federal waters surround by Massachusetts Restricted Area.



The Commonwealth of Massachusetts

Division of Marine Fisheries

251 Causeway Street, Suite 400, Boston, MA 02114
p: (617) 626-1520 | f: (617) 626-1509
www.mass.gov/marinefisheries



CHARLES D. BAKER
Governor

KARYN E. POLITO
Lt. Governor

KATHLEEN A. THEOHARIDES
Secretary

RONALD S. AMIDON
Commissioner

DANIEL J. MCKIERNAN
Director

January 7, 2022

Michael Pentony
Regional Administrator
NOAA Fisheries GARFO
55 Great Republic Drive
Gloucester, MA 01930

RE: Seasonal entanglement risk for North Atlantic Right Whales in the EEZ west of Stellwagen Bank

Dear Mr. Pentony,

I am writing to inform you of an emerging entanglement risk to North Atlantic right whales (NARW) that occurs in a certain zone of federal waters sandwiched between the state and federal closures.

As you are aware, NOAA Fisheries created the Massachusetts Restricted Area (MRA) in 2015 to reduce the risk of entanglement risk to the large aggregations of NARW that occur there seasonally. This closure to fixed fishing gear included MA state waters within Cape Cod Bay and adjacent federal waters around Stellwagen Bank from February 1st through April 30th of each year. DMF immediately created analogous state regulations closing the area to fixed fishing gear.

Since 2016, DMF has also added dynamic management to the state waters portion of the MRA by extending the closure into the month of May when aerial surveillance shows that right whales remain present. In addition to this action, since the beginning of the closure, DMF has engaged in efforts, with assistance from the Massachusetts Environmental Police, to retrieve abandoned gear in the closure annually to ensure that the entanglement risk to right whales is effective as intended.

Since the advent of the MRA closure in 2015, seasonal usage of state and federal waters outside of Cape Cod Bay increased in certain areas and times where fixed gear fishing was allowed. Recent sighting data indicate that NARW stay for a longer time period than they have historically, and these whales are increasingly observed in state and adjacent federal waters in Massachusetts Bay and north to the NH state line. These changes in distribution increased the entanglement risk to NARW along the MA coastal waters. In response to these changes in entanglement risk, as well as continued declines in the population status of NARW, in 2021 DMF closed MA state waters from southeastern Cape Cod north the NH border to lobster fishing from February 1st to May 15th (Figure 1).




Figure 1. DMF trap/gear closure, February 1 – May 15

The National Marine Fisheries Service then mirrored the northern extension of the closure, known as Massachusetts North Restricted Area, in their Phase 1 amendment to the Atlantic Large Whale Take Reduction Plan in September of 2021. The Massachusetts North closure only runs through April 30 each year under the federal plan (Figure 2).




Figure 2. Map of Massachusetts Restricted Area

The increasing presence of NARW in these northern areas is not exclusive to state waters. In recent years, aerial surveillance conducted by the Center for Coastal Studies (CCS) has documented the presence of right whales in both open and closed portions of the waters north of Cape Cod Bay. The map below depicting gear and whales from 2018 demonstrates the necessity for DMF's northern extension of the state waters closure implemented in 2021 (Figure 3).




Figure 3. Right whale sightings in 2018 and buoy lines documented on April 23, 2018

However, the implementation of the Massachusetts North Restricted Area has created a gap between the closed areas between state waters of Massachusetts Bay and the northern federal waters portion of the original Massachusetts Restricted Area (Figure 2 and 4). Federally permitted vessels can continue to fish with persistent buoy lines in these areas adjacent to MA state waters during the closure period, and this area lies beyond the jurisdiction of the Commonwealth.




Figure 4. Right whale sightings in 2021 and buoy lines documented in April and May 2021

Given the current dire status of the NARW population and the need for continued reductions in entanglement risk we wanted to ensure that NOAA Fisheries was aware of this issue. We feel that continued overlap of persistent buoy lines with aggregations of NARW pose an entanglement threat and we are concerned that any future NARW entanglement in waters off the Massachusetts coast could threaten the opportunity of MA-based fishers to participate in fixed gear fisheries.

Sincerely,

Daniel J. McKiernan, Director

CC: Marine Fisheries Advisory Commission

Colleen Coogan
Lead, Marine Mammal Sea Turtle Team
NMFS, Greater Atlantic Region
Colleen.Coogan@NOAA.gov

January 5, 2022

Dear Colleen,

We are writing to ask the Agency to re-evaluate entanglement risk to right whales in the federal waters adjacent to the Massachusetts Bay Restricted Area. We intended to address this issue during the January Atlantic Large Whale Take Reduction Team (ALWTRT or Team) meeting but unfortunately that meeting was canceled. As a result, we ask NMFS to share this letter with the Team and to expeditiously seek additional input from them prior to the February 1 start of the 2022 Massachusetts Bay Restricted Area period.

As the attachments to this letter show, when the Massachusetts Department of Marine Fisheries (MADMF) expands its state water restrictions to the north (Scituate to the New Hampshire border) on February 1, a wedge of unprotected federal waters will be created parallel to the Massachusetts Bay Restricted Area. We are concerned that redistributed gear from both restricted areas could increase entanglement risk in the wedge. Any entanglement will further impact Massachusetts fishermen who are arguably already the most regulated portion of the industry when it comes to reducing risk to right whales.

In an effort to assess whether this area may pose unintentional risk to right whales, we made a public records request to MADMF for past sightings and gear data. Specifically, we requested:

- Data on whether and how much gear was set in this area in 2021, during the restricted season (Feb 1 – May 15).
- For comparison, any data from 2020 prior to the expansion of the MADMF state waters expansion to better understand if a shift in effort had occurred.
- Data on right whale detections (visual and acoustical) for these areas in the spring (2015-2021).
- The overlap of gear and right whales (detected visually or acoustically) in this area between Feb 1 and May 15 of 2021.

MADMF provided a series of maps (attached to this document) with the following caveats:

- Because of how gear is documented during surveys, data on gear should be viewed as more qualitative than quantitative. Therefore, the most accurate way to view the gear data are comparing gear seen in a single day.
- To view potential overlap, a single day of gear was overlaid with multiple years of right whale sightings from 2018 and 2021.
- Sightings of right whales provided used NARWC data on right whale sightings for 2015-2020.

Understanding that past aerial survey effort has been focused on Cape Cod Bay (not this area), the maps are qualitative rather than quantitative and not effort corrected. We also recognize that the gear data are only occasionally collected and represent aggregations rather than individual buoys. Still, it appears

that the risk in this area is not negligible. As a result, we ask NMFS to analyze the following issues and to share the results with the Team:

1. Whether the gear aggregations represent potentially wet stored gear or actively fished gear between February 1st- May 15th (or the end of the restriction period)?
2. If wet stored, is that the result of a lack of land-based storage areas for fishermen using this area? If so, can NMFS and MADMF provide alternative land-based locations in support of the industry and to reduce entanglement risk to right whales?
3. Does NMFS intend to survey this area for gear and right whales during the upcoming restricted period? If so, how often will visual surveys be conducted?
4. How likely will it be for gear set in this area to be properly marked prior to February 1, 2021? Does NMFS and/or MADMF have a gear marking monitoring plan in place and will results be shared with members of the ALWTRT?
5. Should an entanglement occur in this area during the restricted period, does NMFS have a plan to prevent further risk to right whales?

As stated previously, we ask that you share our concerns with the Team and seek their additional input and suggestions as to how to prevent inadvertent risk during the upcoming restricted period. Our intention is not to further impact Massachusetts fishermen, but rather to ensure that all of the efforts put forward by them to date, are not thwarted by increasing gear in a small area where right whales are likely to be present.

Thank you in advance for your consideration,


Regina Asmutis-Silvia
Whale and Dolphin Conservation


Erica Fuller
Conservation Law Foundation


CC:

Bob Glen (robert.glenn@mass.gov)
Bennett Brooks (bbrooks@cbi.org)
Marisa Trego (marisa.trego@noaa.gov)

ATTACHMENTS: Maps provided by MADMF









Dear ALWTRT Members,

I am supporting the concerns identified in the Asmutis/ Fuller letter to the ALWTRT dated January 5, 2022. Their concerns are supported by earlier work looking at gear and whales in the Stellwagen Bank National Marine Sancturay (SBNMS, Wiley et al. 2003, attached). I have provided the MADMS maps included in the Asmutis/Fuller letter and maps from Wiley et al. 2003 for comparison. While the SBNMS boundary is not included in the MADMF maps, it is clear that substantial gear has occurred in the area of concern during periods of right whale occupancy since at least the early 2000's (Wiley et al. 2003) and continues to this day (MADMS maps). Of greater importance might be a comparison of the seasonal movement of gear concentration on and off the southwest corner of the Stellwagen Bank. The extremely high summer concentration of gear on the southwest corner identified in Wiley et al. 2003 is entirely absent during the winter months, with an area to the west exhibiting high gear densities that were absent during the summer months. The accepted reason for this gear migration is not to increase lobster catch during this time, but to move gear to deeper, safer water to reduce winter storm damage, while avoiding the effort and difficulty of moving gear to limited land based storage areas (i.e., winter storage). The imporatrnce of this is that, rather than moving gear to shore when faced with the existing closures, at least some fishermen can be expected to move gear to the nearest areas that remain open, (i.e., the "wedge" descibed in the Asmutid/Fuller letter), thereby increasing entanglement risk in that area during a time when decreased risk is the goal of MADMF, NMFS and ALWTRT. You will also notice that fishing effort is placed along the closure line on the Western Gulf Of Maine Closed Area. I also note that the MADMF maps show a decrease in gear throughout Massachusetts Bay from 2018 – 2021. This is a major contribution by Massachusetts lobster fishermen to the protection of right whales and should be applauded. I hope that this information helps in our understaning and decision-making.

Sincerely,

A handwritten signature in black ink, appearing to read 'David Wiley', followed by a long, wavy horizontal line.

David Wiley, PhD
Research Ecologist
NOAA/Stellwagen Bank National Marine Sanctuary



The Distribution and Density of Commercial Fisheries and Baleen Whales within the Stellwagen Bank National Marine Sanctuary: July 2001–June 2002

PAPER

ABSTRACT

Research in a national marine sanctuary provides the ability to monitor, assess and understand changes in, and threats to, the area. In July 2001, the Stellwagen Bank National Marine Sanctuary undertook a year-long study to quantify and map patterns of human and marine mammal use. Data were collected during monthly standardized shipboard surveys that bisected the Sanctuary at 5 km (2.5 nm) intervals. We used a subset of those data and ArcView's Spatial Analyst program to conduct an analysis of the density and distribution of fixed gear (trap and gillnet) fisheries, mobile gear (otter trawl and scallop dredge) fisheries and baleen whales. We used this to develop a "user geography" of the Sanctuary based on patterns of use and identify high use areas that might pose the risk of environmental damage. We also used ArcView to develop an index of Relative Interaction Potential (RIP) to identify where baleen whales might become entangled in fishing gear, a known threat within the Sanctuary. The RIP identified a number of areas that stood out in terms of entanglement risk. Information from the study will allow managers to identify future changes in Sanctuary use and investigate current areas of intense use for potential harm.

INTRODUCTION

National Marine Sanctuaries (NMS) are Ocean areas of special national significance whose protection and beneficial use require comprehensive management and planning. The primary goal of the NMS program is to protect the designated area's resources. However, multiple uses are allowed if such uses are consistent with the sanctuary's primary goal of resource protection. Because resource protection and resource use are often in conflict, considerable information is needed if legitimate planning and defensible management are to occur, and resources are to be protected in the face of exploitation.

One of the main suites of information needed for successful management and planning is the spatial and temporal distribution of various activities that take place within a sanctuary and the levels at which they occur. Such information can then be used as a baseline against which to measure future changes and to investigate the degree to which such uses might interact with sanctuary resources or other user groups. Unfortunately, few sanctuaries have

such data to guide their decision-making. Most information available to managers is either collected at scales that make its application to sanctuary management questionable or is largely anecdotal. Because sanctuary decisions are often embedded in controversy, such information frequently creates, rather than informs, debate. If good decisions depend on good science, better and more rigorous information must be available to decision-makers (Lubchenco, 1995; Caughley and Gunne, 1996).

The productive waters encompassed by the Stellwagen Bank National Marine Sanctuary (SBNMS or Sanctuary) are home to an impressive array of marine life and are utilized by an equally impressive array of user groups. Public input has indicated high levels of concern over environmental issues such as the potential for habitat degradation by mobile fishing gear and the entanglement of baleen whales in fixed fishing gear. However, few data have been available to help guide Sanctuary management on such topics. In July of 2001, the SBNMS initiated a year-long study with the goal of determining the spatial and temporal distribution of human activity, marine mammals, and selected fish species. In this paper, we used a subset of those data to investigate the spatial and temporal densities of:

1. fixed gear fishing effort (i.e., gillnet and trap fisheries),
2. mobile gear fishing effort (i.e., otter trawl and scallop dredge fisheries), and
3. baleen whales; i.e., humpback (*Megaptera novaeangliae*), right (*Eubalaena glacialis*), fin (*Balaenoptera physalus*), and minke (*Balaenoptera acutorostrata*) whales.

We used those data to depict the "user geography" of the SBNMS and alert managers to areas where intense use or co-occurrence might signal potential harm to sanctuary resources.

METHODS

Study Area—The SBNMS (Figure 1) covers an area of 2,181 km² (842 mi²) in the southwest Gulf of Maine. It is an offshore sanctuary, with its boundary being ~ 5.5 km (3 nm) north of Race Point (Provincetown), MA, ~ 5.5 km (3 nm) southeast of Cape Ann (Gloucester), MA and 46 km (25 nm) east of Boston, MA. The area's main bathymetric feature is Stellwagen Bank, a curved glacial moraine that is almost 37 km (20 nm) in length and over 11 km (6 nm) in

David N. Wiley, Just C.

Moller and Kristin A.

Zilinskas

Stellwagen Bank National

Marine Sanctuary,

Scituate, Massachusetts

David N. Wiley

International Wildlife

Coalition, East Falmouth,

Massachusetts


width at its widest point. Water depths over and around the bank range from 20 to 90 m (65 to 300 ft). To the north of the bank is deeper water (180 m or 600 ft) that rises to ~ 60 m (200 ft) where the Sanctuary border intersects Jeffreys Ledge. There are also numerous smaller bathymetric features. Within this area are seabed types ranging from muddy and sandy bottoms to extensive areas of gravel or small boulder fields (Valentine et al., 2001). The area is home to some of the largest aggregations of baleen whales along the United States' eastern seaboard

(Anon., 1982) and is used extensively by commercial fisheries and recreational interests.

Data Field Collection

Survey Design—To determine the spatial and temporal densities of marine mammals and human activities within the SBNMS, we conducted monthly standardized shipboard surveys along 15 designated tracklines that bisected the Sanctuary in an east/west direction and ran approximately perpendicular to Stellwagen Bank (Figure 1). Track 1 was the southernmost

Figure 1. The Stellwagen Bank National Marine Sanctuary showing survey blocks and tracklines. Monthly survey of tracklines was conducted from July 2001 through June 2002.



line and track 15 was the most northern line. Because the SBNMS is irregularly shaped, tracklines were not of equal length. Track lengths were 37 km (20 nm) for line numbers 1–11, 27 km (14.5 nm) for line numbers 12 and 13, ~17 km (9 nm) for line number 14, and 7 km (4 nm) for line number 15. Total track length was 485 km (262 nm). Tracklines were separated by 5 km (2.5 nm) and survey speed was 12–13 knots. Observations were limited to sea states of ~ Beaufort 4 or less.

Three days were required to complete each month's survey. Trackline coverage was not random. Tracks were grouped into 3 blocks (Figure 1): Block A; tracklines 1–5 (181 km or 97.5 nm of trackline), Block B; tracklines 6–10 (181 km or 97.5 nm of trackline), and Block C; tracklines 11–15 (114 km or 61.5 nm of trackline). Each block required one survey day to complete (including transit time). The order in which blocks were surveyed was determined by prevailing weather conditions and the previous month's survey pattern. Because of weather conditions, survey days were not consecutive, nor were all tracks surveyed in every month (see results).

Observation Platforms—Two observations platforms were used in the study: the *F/V Wavelength* and the *M/V AndyLynn*. The *F/V Wavelength* was a 10 m (32 ft) lobster style boat with an elevated (6 m) "tuna tower" from which observations were made. The *F/V Wavelength* was used for the July and August 2001 surveys. All other surveys were conducted from the *M/V AndyLynn*, a 20 m (65 ft) party fishing boat. Observations from the *M/V AndyLynn* were made from the upper bridge, which was 6 m above the water.

Sighting Categories—We grouped sightings into five major categories. These were: Marine Mammals, Fish, Commercial Fishing Vessels, Vessels (including commercial shipping), and Fixed Fishing Gear. Each category consisted of an assortment of identifiers likely to be encountered during surveys. For example, the Commercial Fishing Vessel category consisted of Stern Trawler¹, Side Trawler², Scallop Dredge, Gillnet Boat, Lobster Boat, Longline Boat, Unidentified, and Other. A vessel's identity was inferred from its observed deck configuration. To confirm identifications, we photographed vessels whenever possible. Differentiation was made between vessels transiting an area and those actively engaged in their trade at the time of observation. Marine mammal and fish identification was made from characteristic field markings.

Data Collection—Data were collected using line transect methodology (Burnham et al., 1980). The data collection team consisted of

three people: two observers and a recorder/observer. Briefly, two observers each searched a 90° portion of a 180° field. The 180° field consisted of a semicircle extending from midship on the starboard side of the survey vessel, forward to midship on the port side of the survey vessel. When not recording, the recorder acted as a roving observer covering the full 180° arc. Observations were made with the unaided eye and with 7 x 50 power binoculars. Observer stations were rotated at the end of each trackline or at ~30 min intervals during adverse weather conditions (e.g. some winter surveys).

Sightings were recorded on hand-held computers using a data collection program similar to that described in Garrett-Logan and Smith (1997) and provided by the NOAA Fisheries³. At each sighting, the recorder documented its time (to the nearest second), identity, estimated radial angle from the ship's heading, estimated distance, number of objects in the sighting (high, low and best estimate), and behavior. For stationary or slow moving surface objects (e.g., fixed gear surface markers or slow moving vessels) radial angles were taken at right angles to the vessel's heading (90° or 270°). For ephemeral or fast moving objects (e.g., marine mammals or sport vessels) radial angles were recorded at the time of sighting. Distances were determined by visually estimating the distance to the sighted objects with the aid of range finding binoculars. Estimates to larger objects were aided by the use of radar, which also provided constant feedback for determining the accuracy of visual estimates to smaller objects that failed to appear on radar (e.g., surface buoys or marine mammals).

Data Processing and GIS Analyses

Determination of the latitude and longitude for sighted objects—The latitude and longitude of sighted objects were calculated using the bearing and range of the object from the ship's location. The ship's location, heading and speed were recorded at five-second intervals using a laptop computer containing "The Cap'n" software⁴ interfaced with a Garmin GPS 48 Navigator. Synchronized time stamps from the hand-held computers (containing sightings) and the laptop (containing vessel latitude/longitude) were used to find the five-second-interval location closest to the time of sighting. This provided the ship's heading and location at the time of the sighting. Deviations by the ship away from the trackline's true east/west orientation were mathematically corrected for and a true bearing to the sighted object was calculated using the radial angle to the sighting. The latitude and longitude of the sighted object were then calculated using the ship's position and the range and corrected bearing to the object.

For analyses, we used a subset of the sighting categories and collapsed those we used into three broad groupings: (1) Baleen Whales, (2) Fixed Fishing Gear, and (3) Mobile Fishing Gear. The Baleen Whale category consisted of humpback, right, fin, and minke whales. The Fixed Fishing Gear category consisted of bullet buoys, high flyers, floatballs, and various combinations of the three (e.g., highflyer with a floatball). For the Fixed Fishing Gear category, we also included the location of individual gillnet and lobster (trap fishing) boats. This was necessary because the use of a particular style of surface buoy is not necessarily unique to either the gillnet or trap fishery. The inclusion of these fishing vessels provided insight into which fishery the buoys were likely to belong. The Mobile Fishing Gear category included stern trawlers, side trawlers and scallop dredges. Stern trawlers with cable in the water, but nets on their net-reels were considered scallop dredges. Only vessels active in their trade at the time of observation were included in the study.

GIS Spatial and Temporal Density Analysis—

To provide an indication of the relative abundance and distribution of the various categories, we grouped sightings into 12-month and seasonal time periods. Seasons were: summer; July, August and September, fall; October, November and December, winter; January, February and March, and spring; April, May and June. Within these periods, all tracklines were not equally surveyed. To correct for differences in effort, we partitioned sightings into strips of 2.5 km (1.25 nm) on either side of a trackline (the effective search area during a survey). For each time period, we divided each sighting within these strips by the number of times the trackline was surveyed, thereby calculating a sightings/month value for each object. We did not correct for differences in sighting probabilities relating to distance from the trackline or sea state. The resulting data were investigated using Geographic Information Systems (GIS) technology (ArcView 8.2) by converting them into a personal geodatabase format with feature classes created from the individual records. These were georeferenced to conform to the Massachusetts State Plane coordinate system (North American Datum, 1983, Lambert Conformal Conic projection) for compatibility with datasets from other sources.

ArcView's Spatial Analyst extension was used to create density surfaces that identified where sightings (e.g., fishing vessels or whales) were concentrated and provided a prediction of their distribution (ESRI, 2001). Density surfaces were created using the Kernel Density function. Values were calculated in square kilometers with an output raster cell size of 100 m²

and a search radius of 5000 m. It is important to note that the resulting densities are greatly dependent on the search radius chosen. For example, larger search radii can link sightings over a larger area, but "dilute" heavy, localized concentrations. Smaller search radii can provide a more accurate quantification of localized densities, but reduce the analysis' ability to provide a broad understanding of patterns over a wider area. Our choice of a 5000 m search area was a compromise between these factors, with an emphasis on the goals of identifying the broader patterns of uses occurring within the Sanctuary and providing a baseline against which future changes could be measured.

Once the density areas were calculated, the range of density values was divided into ten equal interval classes. The relative large number of classes was selected in order to provide a better visualization of the data range and the areas of different use concentrations. For a category's seasonal maps, we used the season with the greatest range of densities as the basis for creating the classes for all other seasons. This allowed densities and patterns to be compared among seasons. However, as explained above, within each density surface are areas of higher and lower concentrations than reported in the accompanying class boundaries. For all maps, we provided the sightings data from which the density contours were calculated. In some cases a single sighting contained multiple objects. This was particularly common for surface buoys in the northwest portion of the Sanctuary and on Stellwagen Bank's southwest corner in the summer when gear aggregations were extremely dense.

*GIS Interaction Potential Analysis—*To investigate the potential for interaction between the Baleen Whale and Fixed Fishing Gear categories, we developed an index of Relative Interaction Potential (RIP). To derive the RIP, we created a matrix of five-minute grid cells that covered the SBNMS. The grid matrix was generated using the ArcInfo's *Generate* command and *Fishnet* option. Within each grid cell, we multiplied the total number of sighted objects within the two categories being investigated⁶. This resulted in a range of numbers for each grid cell that represented the potential for interaction. For example, if a grid cell had no whales (a zero value) and any number of fixed gear, the resulting value is zero or no probability of interaction. The same result would occur for any number of whales and no fixed gear. At the other extreme, if a grid cell had a large number of whales and a large number of fixed gear, a large index value would be calculated representing a much higher potential for interaction. To normalize the result, the index values were divided by the area within the grid cells.

For comparisons, we aggregated grid cell values into quartiles.

RIPs were calculated for the 12-month period and on a seasonal basis. For seasonal RIPs, we used the season with the greatest range of interaction potentials as the basis for creating the four classes upon which other seasons were based. This allowed RIPs to be compared among seasons. For greater visualization, we provided the sightings data from which the RIPs were calculated.

RESULTS

Survey Effort—For the twelve-month period July 2001–June 2002, a total of 5,700 km (3,078 nm) miles of trackline were available to be surveyed, of which 4,460 km (2,408 nm) miles (78%) were completed. Tracks 1–3 had the greatest coverage (92%) and tracks 11–14 had the least (58%) (Table 1). By month, survey coverage was greatest in July, October, April, and June (100%) and least in February (30%) (Table 1). For the entire survey area, the number of track-miles surveyed did not differ significantly by season (ANOVA $F=2.86$, $P=0.104$). However, considerably less trackline was covered during the winter months. Percent trackline coverage by season was: summer; 83%, fall; 87%, winter; 51%, and spring; 92%.

Sightings—The analyses is based on 6,526 sightings of 9,991 objects (a sighting could contain multiple objects). The totals by category were: Fixed Gear; 4,963 sightings of 6,130 surface buoys, Gillnet Boats; 55 sightings of 56

boats, Lobster Boats; 100 sightings of 101 boats, Mobile Fishing Vessels; 187 sightings of 189 vessels, and Baleen Whales; 352 sightings of 414 animals.

Spatial and Temporal Density—Fixed gear fishing effort (gillnet and trap fisheries)

Twelve-month summary—Numerically, fixed fishing gear was the dominant human use of the SBNMS and it occurred throughout the Sanctuary (Figure 2). Density surfaces ranged from a high of 1.73–1.92 surface bouys/km²/month around the southwest corner of Stellwagen Bank and the northwest section of the Sanctuary off Cape Ann, to lows of 0.0–0.19 surface bouys/km²/month, primarily in the southeastern section of the Sanctuary. The dense areas coincided with the presence of trap fishing vessels, suggesting concentrations of fishing gear targeting lobster or, in some cases, crab.

In general, the density of fixed fishing gear was greatest in the western portions of the Sanctuary and diminished to the east. The presence of trap fishing vessels was also greatest in the western portions of the Sanctuary, suggesting that much of this activity was associated with the lobster/crab fishery. While the level of fixed fishing activity decreased to the east, substantial levels of use still occurred there. These levels were highest (~0.2–0.6 surface bouys/km²/month) in an area northeast of Stellwagen Bank and along a line delineating the Western Gulf of Maine Closed Area⁶ (WGMCA), an area closed to groundfishing

Table 1. Coverage of survey tracklines by month, season and year. Dark blocks signify surveyed tracklines and white blocks signify tracks that were not surveyed. Each month's survey consisted of 15 tracklines totalling 475km (256nm). Surveys occurred within the Stellwagen Bank National Marine Sanctuary from July 2001 through June 2002.

		Track Line																	
		Block 1					Block 2					Block 3							
		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	Percent of Survey Area Completed by Month		
Summer	July '01																100		
	August																78		
	September																	70	
Fall	October																100		
	November																		85
	December																	77	
Winter	January '02																38		
	February																30		
	March																85		
Spring	April																100		
	May																	76	
	June																100		
Percent of Track Surveyed for the Year		92	92	92	83	83	75	83	83	83	83	58	58	58	58	75			

(e.g., Atlantic cod (*Gadhus morhua*)). These areas coincided with the presence of gillnet fishing vessels, indicating that this fishery occurred primarily in the eastern and northern portions of the Sanctuary. With the exception of the southwest corner of Stellwagen Bank, there was a tendency for fixed gear not to be associated with the shoal water of Stellwagen Bank itself.

Seasonal summary—There were substantial seasonal changes in the level and distribution of surface buoys indicating the presence/absence of fixed fishing gear (Figure 3). The densest aggregation occurred during the summer months around the southwest corner of Stellwagen Bank (3-4 surface bouys/km²/month). This aggregation persisted at reduced levels in the spring and fall (~1 surface

Figure 2. The density and distribution of surface buoys within the Stellwagen Bank National Marine Sanctuary from July 2001 through June 2002. Each point represents the sighting of one or more surface buoys. Surface buoys are indicators of fixed fishing gear (trap or gillnet) "sets" that can extend thousands of meters along the seafloor. Two surface buoys equals one set. Trap and gillnet sets cannot be unambiguously differentiated by surface buoys. Sightings of actively fishing lobster (trap) and gillnet vessel are provided as an aid to determining the type of gear in an area.





Figure 3. The seasonal density and distribution of surface buoys within the Stellwagen Bank National Marine Sanctuary from July 2001 through June 2002. Each point represents the sighting of one or more surface buoys. Surface buoys are indicators of fixed fishing gear (trap or gillnet) "sets" that can extend thousands of meters along the seafloor. Two surface buoys equals one set. Trap and gillnet sets cannot be unambiguously differentiated by surface buoys. Sightings of actively fishing lobster (trap) and gillnet vessel are provided as an aid to determining the type of gear in an area.




bouys/km²/month), but was absent during the winter months. Another dense seasonal aggregation occurred in the northwest section of the Sanctuary during the spring, winter and fall (~1–2 surface bouys/km²/month). A third concentration occurred during the winter in an area to the west of Stellwagen Bank. Excluding the high use area around the southwest corner of Stellwagen Bank, fixed gear was most abundant in the Sanctuary during the fall, winter and spring and was primarily associated with trap vessels.

Spatial and Temporal Density—Mobile fishing effort

Twelve-month summary—There were two major concentrations of mobile fishing vessels (Figure 4). The densest aggregation (0.048–0.052 vessels/km²/month) occurred in the southeast section of the Sanctuary. The primary vessels associated with that area were scallop dredges, although substantial numbers of stern and eastern trawlers also worked the area. A second aggregation occurred over a broad area covering

Figure 4. The density and distribution of mobile fishing vessels (stern dragger, eastern dragger and scallop dredge) within the Stellwagen Bank National Marine Sanctuary from July 2001 through June 2002. Each point represents the sighting of an active fishing vessel.



the Sanctuary's northwest quarter and consisted primarily of stern and eastern trawlers. Monthly densities in this region ranged up to 0.036 vessels/km²/month. With the exception of the heavily used portion in the southeast corner, mobile vessel's made less use of the Sanctuary's eastern section and the shallower area on top of Stellwagen Bank proper.

Seasonal Summary—The major use areas identified in the 12-month summary were retained on a seasonal basis (Figure 5). The southeast segment was used in all seasons, with scallop vessels most prevalent in winter and summer. Stern and eastern trawlers remained active in the northwest section, with a tendency to move further offshore in the spring.

Spatial and Temporal Density—Baleen Whales

Twelve-month summary—The highest use area for baleen whales was around the southwest corner of Stellwagen Bank (0.11–0.12 whale/km²/month), followed by the area around Jeffreys Ledge (~0.08–0.09 whale/km²/month) (Figure 6). Other areas of concentration occurred around the southeast and northwest corners of Stellwagen Bank (0.05–0.06 whale/km²/month and 0.02–0.04 whale/km²/month, respectively).

Seasonal Summary—The greatest concentrations of baleen whales occurred during the summer months around the southwest corner of Stellwagen Bank (0.36–0.39 whale/km²/month) and Jeffreys Ledge (0.25–0.28 whale/km²/month) (Figure 7). Other high use areas by season were: fall; southeast corner of Stellwagen Bank (0.13–0.16 whale/km²/month), winter; a small area in the Sanctuary's northeast quarter (0.36–0.39 whale/km²/month)⁷ and spring; Jeffreys Ledge (0.13–0.16 whale/km²/month/month).

Interaction Potential—Baleen Whales and Fixed Fishing Gear

Twelve-month summary—The highest potential for interaction between baleen whales and fixed fishing gear (top quartile 5-minute blocks) were the areas around the southwest and northwest corners of Stellwagen Bank (Figure 8). These areas consisted of six, 5-minute blocks around the Bank's southwest corner and three 5-minute blocks around the northwest corner. Second-level interaction areas were located in the northern portion of the Sanctuary along the southern border of Jeffreys Ledge (three 5-minute blocks), a one 5-minute block section in the southeast portion of the Sanctuary, and a five, 5-minute block area that was contiguous with the high RIP areas of the northwest and southwest corners of the Bank. The highest

RIPs occurred around the southwest corner of the Bank.

Seasonal summary—The greatest areas of top ranked RIPs occurred during the spring and summer around the southwest and northwest corners of Stellwagen Bank, with each season possessing five top ranked index areas (RIP = 3.34–98.28) (Figure 9). The fall exhibited two top ranked index areas, one on western Jeffreys Ledge and one in the most southwestern portion of the Sanctuary. There were no top ranked RIP areas during the winter season. The southeastern section of the Sanctuary consistently exhibited the lowest RIPs in all seasons.

DISCUSSION

Fulfilling a sanctuary's dual mandate of multiple use and resource protection requires an understanding of how human activities are conducted, how those activities might impact the environment, and where and at what levels they occur. To that end, we provide a brief description of each fishery and its reported potential environmental impacts. We then use the survey results to describe patterns of use and identify areas that might be at risk of harm. We also used the RIP index to identify where baleen whales might be at the greatest risk of entanglement and suggest ways to mitigate such interactions.

We offer a number of caveats to this discussion. First, there is no known metric equating the density of fishing effort with environmental harm and the degree to which Sanctuary resources might be impacted, if at all, is unknown. Second, the reported fishing effort and distribution must be viewed with knowledge of the concurrent fisheries management regime, such as the patchwork of closures implemented by the New England Fisheries Management Council (NEFMC) to reduce groundfishing effort (Table 2) and the year-round Western Gulf of Maine Closed Area. Changes in fisheries management will undoubtedly change current fishing patterns. Finally, even long-term monitoring data are more powerful in explaining the past than predicting the future (Bondrup-Nielson and Herman, 1995). Our data provide a valuable snapshot of occurrences within the SBNMS from July 2001–June 2002. The degree to which they reflect previous or future occurrences is unknown.

DESCRIPTION OF FISHERY TYPES

Fixed Gear Fisheries

Trap Fishery—Trap fisheries employ a passive methodology in that traps sit on the seabed and

Figure 5. The seasonal density and distribution of mobile fishing vessels (stern dragger, eastern dragger and scallop dredge) within the Stellwagen Bank National Marine Sanctuary from July 2001 through June 2002. Each point represents the sighting of an active fishing vessel.





Figure 6. The density and distribution of baleen whales; i.e., humpback (*Megaptera novaeangliae*), right (*Eubalaena glacialis*), fin (*Balaenoptera physalus*), and minke (*Balaenoptera acutorostrata*) whales, within the Stellwagen Bank National Marine Sanctuary from July 2001 through June 2002. Each point represents the sighting of one or more whales.



use bait (usually dead fish) to attract lobsters, and to a lesser extent crabs, to the traps. Traps are wire or wooden cages that typically measure 91 cm by 53 cm by 34 cm (36 in by 21 in by 13.5 in), although some can be larger. Traps are often fished in "trawls" consisting of a number of traps leading off a common "ground line". In the area around the SBNMS, trawls typically consist of ~ 25 traps spaced 30-55 m (100-

180 ft) apart (W. Hoffman, Massachusetts Division of Marine Fisheries, Boston, MA, Pers. Comm.) Therefore a single trawl can be over 1,219 m (4,000 ft) in length. Ground lines along the length of the trawl characteristically consist of buoyant polypropylene line that can float more than 5 m (16 ft) above the bottom (McKiernan et al., 2002). On each end of a trawl, a "buoy line" runs from the gear to a

Figure 7. The seasonal density and distribution of baleen whales; i.e., humpback (*Megaptera novaeangliae*), right (*Eubalaena glacialis*), fin (*Balaenoptera physalus*), and minke (*Balaenoptera acutorostrata*) whales, within the Stellwagen Bank National Marine Sanctuary from July 2001 through June 2002. Each point represents the sighting of one or more whales.





Figure 8. Relative Interaction Potential (RIP) index showing the potential for interaction between baleen whales and fixed fishing gear, by 5-minute square area. The index was calculated by multiplying the total number of fixed gear surface buoys within a 5-minute square by the total number of whales sighted in that square. Results were compared by quartile. Data were collected from July 2001 through June 2002.



buoy visible at the surface (i.e., the surface buoy). It is important to note that the surface buoy counts provided in our results represent unseen fishing gear on the seafloor. As described above, two surface buoys might indicate the presence of over 1,219 m (4000 ft) of lobster gear. Since 1990, the lobster fishery has ranked first in landed fish value for New England waters (Pol and Carr, 2000).

Environmental Issues of Trap Fisheries— Lobster/crab traps are a passive fishing gear that has minimal impact on the seabed. In addition, they pose minimal threat to small cetaceans such as porpoise and dolphins, or seabirds. The fishery also has a number of mechanisms that allow it to achieve substantial selectivity. For example, traps are fitted with an escape panel along a lower edge that allows

Figure 9. Seasonal Relative Interaction Potential (RIP) indexes showing the potential for interaction between baleen whales and fixed fishing gear, by 5-minute square area. The index was calculated by multiplying the total number of fixed gear surface buoys within a 5-minute square by the total number of whales sighted in that square. Class ranges were developed by taking the quartiles for the season with the greatest range in RIP values (summer) and applying them to all other seasons. Data were collected from July 2001 through June 2002.




Table 2. Groundfish closures within the Stellwagen Bank National Marine Sanctuary by month. Closures were instituted by the New England Fishery Management Council to recovery depleted groundfish stocks.

Area	Closure Dates
Entire Sanctuary	1 – 30 April
North of 42° 30' 00"	1 – 31 May
South of 42° 30' 00"	1 October – 30 November 1 – 28 February 1 – 31 March

sub-legal size animals to pass through it. In addition, because lobsters are live captured, immatures or females brooding eggs can be returned to the water unharmed. Traps are also fitted with corrodible links which cause lost traps to fall apart, limiting the time spent as ghost fishing gear.

A drawback to the fishery is its interaction with baleen whales. Right, humpback, fin, and minke whales are all known to become entangled in the buoy lines running from traps to the surface or in the groundlines floating off the bottom between traps (Waring et al., 2001, Kenney and Hartley, 2001). This issue is most severe for the highly endangered right whale. Unless anthropogenic mortality in this species is reduced, it is projected to become extinct within ~ 200yrs (Caswell et al., 1999) and lobster gear has been identified as a major threat to the species (50CFR229.32). As a result, NOAA Fisheries has promulgated the Atlantic Large Whale Take Reduction Plan to reduce the incidental take of baleen whales in the lobster fishery, with a focus on right whale protection (50CFR229.32).

Distribution and Seasonality of Trap Fishing Gear—The trap fishery was focused on the western half of the Sanctuary. A dense aggregation of traps existed in the Sanctuary's most northwestern section (just off the coast of Cape Ann, MA) with areas of decreasing density radiating out from that hub. Trap fishing was the dominant commercial fishery in the southwestern portion of the Sanctuary, with a particularly dense area located on the southwest corner of Stellwagen Bank.

With the exception of the dense aggregation of traps located on the southwest corner of Stellwagen Bank, traps were at a Sanctuary minimum during the summer months. This is because lobsters, the fishery's main target, are concentrated in shallow, near-shore waters during that season. As lobsters move offshore in the fall, fishermen follow them into the deeper waters west of Stellwagen Bank. By winter, a substantial portion of the fishery is focused in that area and the water immediately west of the Sanctuary's western boarder. This concentra-

tion is due to the lack of lobster in the near-shore waters and because traps in shallow water (<~ 25 m or 80 ft) are vulnerable to destruction caused by winter storms (W. Adler, President, Massachusetts Lobstermen's Assoc. Marshfield, MA, pers. comm.). In springtime fishermen reverse the process, following lobster from the deeper waters of the Sanctuary back to near-shore waters.

The trap fishery that focused on the southwest corner of Stellwagen Bank was the exception to this trend. This fishery targeted crab and was at its peak during the summer months, when fishing densities were among the highest observed anywhere within the SBNMS. In the fall, this fishery shifted slightly west, as fishermen targeted the more profitable lobsters in the storm-safe deeper waters west of the Bank, and some numbers of them likely remained through the winter. In the spring, the disappearance of lobsters from deep water and a reduction in storm frequency and severity accompanied the re-establishment of the fishery on the Bank's southwest corner.

Potential areas of concern—Potential areas of concern for the trap fishery are covered under the section on Interaction Between Fixed Gear and Baleen Whales.

Gillnet Fishery—Gillnets are comprised of thin, transparent, monofilament webbing stretched between a buoyant "float line" running along the top on the net and a heavy "lead line" running along the bottom. Tension between the buoyant float line and the heavy lead line causes the webbing to rise from the seabed to a height of 2.5 to 3.6 m (8 to 12 ft). If flatfish (e.g., flounder) are targeted, the float line and lead line are tied together, limiting the height to ~ 1 m (3 ft). A single net is ~ 91 m (300 ft) long and nets are joined together into "strings". In the Gulf of Maine, net strings range between 458 m (1500 ft) and 2,292 m (7,500 ft) in length (Read, 1994). Each end of a string is marked on the surface with a buoy (usually a "high flyer") that is attached to the gear by a line also used for hauling. Strings of gillnets are often set in a zigzag or even circular pattern, with small weights along the lead line acting as pivot points. As with the trap fishery, it is important to note that an observation of two surface buoys can indicate the presence of hundreds or thousands of meters of netting on the seafloor below them. The landed value and ranking of New England's gillnet fleet has varied greatly since its resurgence in the 1970's. Pol and Carr (2000) ranked gillnetting fourth in landed value in 1997, the most recent year of analysis.

Environmental issues of the Gillnet Fishery—As a passive fishing gear, gillnets have mini-

ment in and out of the Sanctuary. The potential areas of concern involving gillnet interactions with baleen whales are covered under the section on Interaction Between Fixed Gear and Baleen Whales.

num impact upon the seabed. An additional positive attribute is that they can be size selective, allowing undersized fish to pass through the webbing uncaught (Hamley, 1975). However, gillnets are relatively unselective in terms of the species that become entrapped in them (see Perrin et al., 1994). For example, almost all marine mammals frequenting the SBNMS are vulnerable to incidental kill in gillnets (e.g., Kraus, 1990; Read, 1994; Wiley et al., 1995; Waring et al., 2001). Seabirds and marine turtles can also be incidentally caught during fishing operations.

Several attempts have been made to reduce the kill of non-target species in gillnets. This includes the use of acoustic devices to deter harbor porpoise (*Phocoena phocoena*) from nets during some portions of the year when porpoise are in the Sanctuary. This mitigation attempt has also raised concerns. If acoustic deterrents are aversive to harbor porpoise instead of simply alerting them to the presence of nets, they could act as a barrier to porpoise movement. As with the lobster fishery, the gillnet fishery is subject to the Atlantic Large Whale Take Reduction Plan to reduce the incidental take of baleen whales. Specific information on that plan can be found in 50CFR229.32.

Distribution and Seasonality of Gillnet

Fishing Gear—Gillnetting was most prevalent in the northern and eastern portions of the Sanctuary, and was the dominant fishing activity in the Sanctuary's northeast quarter. The densest aggregation of gillnet activity occurred south of Jeffreys Ledge along a line formed by the Western Gulf of Maine Closed Area. A second concentration of gillnet activity occurred within a broad area along the northeast flank of Stellwagen Bank and another in the northwest section of the Sanctuary off Cape Ann. On a seasonal basis, fewer gillnet boats were observed in the spring than in other seasons.

Some gillnet vessels and unidentified fixed gear were observed in the Western Gulf of Maine Closed Area, mostly during the summer and fall. Unidentified fixed gear could indicate illegally fished gillnets or could belong to legally operating lobster or hagfish boats. The inability to differentiate between legal and illegal gear presents a substantial management problem within this section of the Sanctuary.

Potential areas of concern—If acoustic deterrent devices on gillnets act as a deterrent to harbor porpoise movements, the most likely area of impact would be along the northern boarder of the Western Gulf of Maine Closed Area, where a concentration of pingered gillnets could potentially impede porpoise move-

Mobile Gear Fisheries

Mobile gear fisheries consisted of otter trawls and scallop dredges. There was also a single observation of a hydraulic clam dredge operating in the west central part of the sanctuary. Because many of the sea floor impacts of these fisheries are similar their environmental issues will be discussed jointly.

Otter Trawl Fishery—Otter trawlers or "draggers" target primarily groundfish by towing a large conical net along the seabed (Von Brandt, 1984). The net opening is maintained by the action of a buoyant "headrope" (on the top), a weighted "footrope" (on the bottom), and the spreading affect of heavy trawl "doors" (up to 450 kg or ~1,000 lbs) on either side of the net's mouth. The resistance of the doors moving through the water maintains a net opening width of 15 to 25 m (50–80 ft) (Carrothers, 1981).

Fish are captured by the forward motion of the net along the bottom, which causes fish to enter the net's mouth and collect in the anterior "codend". Fish capture is facilitated by the movement of the footrope along the bottom that disturbs bottom dwelling fish and forces them up into the path of net. The footrope can be modified with rollers or other devices that provide fishermen with access to rocky or uneven bottom (Carr and Milliken, 1998). From 1950 through 1990, trawlers ranked first in landed fish value for New England and second from 1990–1997 (Pol and Carr, 2000).

Scallop Dredge Fishery—A scallop dredge consists of a ~5 m (15 ft) wide rigid metal box trailing a bag of metal rings. The weight of the dredge (up to 700 kg or 1500 lbs) and the angle of the forward cutting bar force the dredge to dig a few centimeters (1-2 in) into the seabed. The forward motion of the cutting bar dislodges scallops from the bottom causing them to pass over the bar and collect in the trailing chain bag. Scallop vessels usually tow two dredges simultaneously at speeds under ~ 5 knots (Rango and McSherry, 2001). Scallop dredges are considered "dry" dredges in that they do not use water jets or suction in the capture process. From 1950–1997, scallop dredges ranked third and occasionally second (1950 and 1980) in landing values for New England's commercial fisheries (Pol and Carr, 2000).

Environmental Issues of Mobile Gear

Fisheries—The issues of trawl and/or dredge impact on bottom habitat and benthic fauna, and the associated impact on marine biodiversi-

ty and the recruitment of commercial stocks, is hotly debated. Numerous authors have documented at least short-term impacts to the seabed and/or benthic fauna (see reviews in Jennings and Kaiser, 1998; Turner et al., 1999; and DeAlteris et al., 2000). In general, mobile gears were found to disrupt bottom substrate, suspend fine sediments and remove or damage large epifaunal invertebrates, often in only a single pass (Fresse et al., 1999). However, these impacts must also be measured against natural disturbances to the seabed caused by forces such as storm activity (DeAlteris et al., 1999) and the long-term environmental impact of mobile gear is not understood. An additional issue is the bycatch of non-target species or size classes during the fishing process.

Distribution and Seasonality of the Otter Trawl Fishery—The broadest and densest area of otter trawl activity occurred in the most northwest section of the Sanctuary, off the coast of Cape Ann, MA. Another focus of trawl activity was in the Sanctuary's most southeastern area off the tip of Cape Cod, MA. Smaller pockets of trawling occurred just south of the Stellwagen Bank's southwest corner and on Jeffreys Ledge along the northern border of the Sanctuary.

The distribution of otter trawl activity showed indications of distinct seasonality. The concentration of activity off Cape Ann persisted throughout the winter, summer and fall, but disappeared in the spring when almost all trawling was focused around the northwest corner of Stellwagen Bank and a second area just to the northeast of that area. These areas harbored little or no fishing in other seasons. Similarly, an area around the southeast corner of Stellwagen Bank was fished heavily in the spring, but was much reduced in the winter. The area just south of Stellwagen Bank's southwest corner was fished primarily in the fall and an area just south of Stellwagen Bank's northwest corner was fished primarily during the summer. No otter trawl activity was observed in the Western Gulf of Maine Closed Area.

Distribution and Seasonality of the Scallop Dredge Fishery—The scallop dredge fishery showed distinct geographic fidelity, being confined primarily to the southeastern portion of the Sanctuary. Based on vessel density, this locale exhibited greater use than any other area targeted by the mobile gear sector. A far lesser area of scallop dredge activity extended from that area in a broad swath across Stellwagen Bank and up the mid section of its western slope. Low levels of scallop dredge activity were observed in the northwest section of the Sanctuary and no scallop vessels were observed in the northeast section. No scallop

dredge activity was observed in the Western Gulf of Maine Closed Area.

Seasonality in the scallop fishery was pronounced, with the greatest effort in the winter and the least in the spring. However, these patterns are complicated by the ability of stern and eastern rigged trawlers to be involved in the scallop fishery, but assigned to the trawling category.

Potential areas of concern—Based on levels of activity, the greatest areas of concern would be in the vicinity of the southeast corner of Stellwagen Bank, where scallop dredges and otter trawler occurred in relatively high numbers and the northwest section of the Sanctuary where relatively high levels of otter trawling occurred. However, if habitat impact is dependent on substrate type, lesser-used areas might be equally or more negatively impacted than those areas identified only through intensity of use.

Interactions between fixed gear and baleen whales -

Entanglement in fixed gear is an identified mortality threat for most species of baleen whales, and both gillnet and trap fisheries have been implicated (Waring et al., 2001). Since the creation of the SBNMS in 1990, numerous sightings of entangled whales have occurred within its border and whales have been observed becoming entangled in the Sanctuary (e.g., Weinrich, 1999).

The Relative Interaction Potential (RIP) index suggested that the most likely sites of whale entanglement would be Stellwagen Bank's southwest and northwest corners, followed by southern Jeffreys Ledge. The highest RIPs occurred in the summer around the southwest corner of Stellwagen Bank. The analysis' prediction was retroactively corroborated by the sighting of three entangled humpback whales on Stellwagen Bank's southwest corner in late July and August of 2001 (Center for Coastal Studies, Provincetown, MA, unpublished data). While entangled whales can tow fishing gear hundreds of miles, the occurrence of entangled whales within the highest RIP areas strengthens the possibility that at least some of the interactions occurred there. The high RIP values associated with the southwest and northwest corners of Stellwagen Bank and to a lesser extent southern Jeffreys Ledge are also areas where entangled whales are frequently reported, although this is complicated by the fact that the whale watching vessels reporting entanglements are also concentrated in those areas.

In summary RIPs were capable of identifying interaction "hot spots" and could provide managers with the opportunity to manage at scales smaller than the entire sanctuary. In terms

of whale entanglement, managers could use RIPs to target specific areas for actions such as fishery closures, gear modifications, or intensive surveillance to facilitate rescue attempts. They can also be used as a valuable tool to facilitate dialogue and information exchange between interest groups seeking solution to the problem.

CONCLUSION

National Marine Sanctuaries are often in the difficult position of protecting resources while promoting a multiple use philosophy. This can only be accomplished through information that allows decision-makers to understand the abundance and distribution of Sanctuary resources, and the magnitude and distribution of potential interactors. The use of shipboard surveys and GIS analyses can quantify such information and provide important insights for management, such as the co-occurrence of vulnerable resources and potentially harmful human activities. However, it must be emphasized that there is no current metric equating levels of activity with harm. While we have chosen to use the results to explore environmental risk, they could also be used to identify areas where the Sanctuary plays an important economic role in the local community. The fact that zones of intense use can simultaneously be areas of elevated environmental risk and increased economic benefit represents a major challenge to Sanctuary management.

An additional benefit of mapping distributional data is the ability to gain a broader understanding of the Sanctuary by using it as a foundation for soliciting local knowledge. For example, discussions with local lobstermen led to an understanding of the impact that storm activity and water depth had on fishing patterns. Thus, data such as ours can be a tool for initiating important dialogue between the Sanctuary and the public, a concept that lies at the heart of the NMS program. While we have attempted a sample analysis and discussion of our data, we await the complex scrutiny, review and input of the many interested parties that make up the SBNMS community. In this way the explanatory power of our data will be maximized and a deeper understanding of the Sanctuary will evolve.

ACKNOWLEDGEMENTS:

We thank Andy Costa (*M/V AndyLynn*), David Hobson (*F/V Wavelength*), Lisa Conger (New England Aquarium), Elizabeth Pomfret-Wiley (National Marine Fisheries Service), and Sara Baldauf-Wagner (SBNMS) for valuable help during the surveys. David Wiley's research was partially funded by

an Environmental Leadership Award from the Switzer Foundation. This is SBNMS contribution 03-02.

REFERENCES

- Anonymous. 1982. CeTAP: A characterization of marine mammals and turtles in the mid- and North Atlantic areas of the U.S. outer continental shelf. Final Report to the Cetacean and Turtle Assessment Program. Pages 450. University of Rhode Island.
- Bondrup-Nielson. S. and Herman, T. B. 1995. Long-term monitoring of the environment: placebo or placebo! In: *Ecosystem Monitoring and Protected Areas*, eds. S. Bondrup-Nielson, T.B. Herman, J. Willison, and N.W. Munro, pp. 22-26. Wolfville: Science and Management of Protected Areas Association.
- Burnham, K. P. Anderson. D. R. and Laake. J. L. 1980. Estimation of density from line transect sampling of biological populations. *Wildlife Monographs* 72: 1-202.
- Carr. H.A. and Milliken, H. O. 1998. Conservation engineering: options to minimize fishing's impacts to the sea floor. In: eds. E. M. Dorsey and J. Perdersen, pp. 100-103. *Effects of Fishing Gear on the Sea Floor of New England*. Conservation Law Foundation, Boston, MA.
- Carrothers. P. J. 1981. Catch variability due to variations in groundfish otter trawl behavior. In: *Bottom Trawl Surveys*, eds. W. G. Doubleday and D. Rivard, pp. 2547-257. Canadian Special Publication of Fisheries and Aquatic Sciences.
- Caswell. H. M. Fujiwara M. and Brault. S. 1999. Declining survival probability threatens the North Atlantic right whale. *Proceedings of the National Academy of Science* 96: 3308-3313.
- Caughley. G. and Gunne. A. 1996. *Conservation Biology in Theory and Practice*. Cambridge: Blackwell Science. 459 pp.
- DeAlteris. J. L. Skrobe. L. and Lipsky. C. 1999. The significance of seabed disturbance by mobile fishing gear relative to natural processes: A case study in Narragansett Bay, Rhode Island. In: *Fish Habitat: Essential Fish Habitat and Rehabilitation*. ed L. Benaka, pp. 224-237. Bethesda: American Fisheries Society.
- DeAlteris. J. T. Skrobe. L.G. and Castro. K.M. 2000. Effects of mobile bottom fishing gear on biodiversity and habitat in offshore New England waters. *Northeastern Naturalist* 7: 739-394.
- ESRI. 2001. *Using ArcGIS Spatial Analyst*. Redlands, California: Environmental Systems Research Institute.
- Freedman, B., Staicer. C. and Woodley. S. 1995. Ecological monitoring and research in greater ecological reserves: A conceptual framework. In: *Ecosystem Monitoring and Protected Areas*, eds. S. Bondrup-Nielson, T.B. Herman, J. Willison, and N.W. Munro, pp. 68-80. Wolfville: Science and Management of Protected Areas Association.
- Freese. L. Auster. P. Heifetz. J. Heifetz. and Wing. B.L. 1999. Effects of trawling on seafloor habitat and associated invertebrate taxa in the Gulf of Alaska. *Marine Ecology Progress Series* 182: 119-126.
- Garret-Logan. N. and Smith.T. 1997. A hand-held, pen-based computer system for marine mammal surveys. *Marine Mammal Science* 13: 694-700.

- Hamley, J. M. 1975. Review of gillnet selectivity. *Journal of the Fisheries Research Board of Canada* 32: 1943–1969.
- Kraus, S. D. 1990. Rates and potential cause of mortality in North Atlantic right whales. *Marine Mammal Science* 6: 278–290.
- Lubchenco, J. 1995. The role of science in formulating a biodiversity strategy. *Bioscience* 45: s-7–s-9.
- McKiernan, D. Pol. M. and Malkoski, V. 2001. A study of the underwater profiles of lobster trawl groundlines. Massachusetts Division of Marine Fisheries, Boston, MA.
- NMFS. 1999. Taking of marine mammals incidental to commercial fishing operations; Atlantic large whale take reduction plan regulations. Pages 7529–7556. NMFS, Washington, D.C.
- Perrin, W. F. Donovan, G. P. and Barlow, J. eds. 1994. *Gillnets and Cetaceans*. Cambridge: International Whaling Commission. 629 pp.
- Pol. M. and Carr, H. A. 2000. Overview of gear developments and trends in the New England commercial fishing industry. *Northeastern Naturalist* 7: 739–794.
- Rago, P. and McSherry, M. 2001. Spatial distribution of fishing effort for sea scallops: 1998–2000. Conference on the Effects of Fishing Gear on Fish Habitat in the Northeastern U.S., October 23–25, 2001, Boston, MA.
- Read, A. J. 1994. Interactions between cetaceans and gillnet and trap fisheries in the Northwest Atlantic. In: Pages 617 in *Gillnets and Cetaceans*, eds. W.F. Perrin, G. P. Donovan, and J. Barlow, pp. 133–148. Cambridge: International Whaling Commission.
- Tuck, I. D. Hall, S. Robertson, M., R. Armstrong, E. and Basford, D.J. 1998. Effects of physical trawling disturbances in a previously unfished sheltered Scottish sea loch. *Marine Ecology Progress Series* 162: 227–242.
- Turner, S. J. Thrush, S. F. Hewitt, J. E. Cummings, V. J. and Funnell, G. 1999. Fishing impacts and the degradation or loss of habitat structure. *Fisheries Management and Ecology* 6: 401–420.
- Valentine, P. C. et al. 2001. Sea Floor Topography and Backscatter Intensity Images of Stellwagen Bank National Marine Sanctuary off Boston, Massachusetts: U.S. Geological Survey Open-File Report 00-410, U.S. Geological Survey, Reston, VA.
- Von Brandt, A. 1984. *Fish Catching Methods of the World*. England: Fishing News Books Ltd. 418 pp.
- Waring, G. Quintal, J. M. and Swartz, S. 2001. U.S. Atlantic and Gulf of Mexico stock assessments–2001. Pages 310. U.S. Dept. of Comm., NOAA, NMFS, Northeast Fisheries Science Center, Woods Hole, MA.
- Weinrich, M. 1999. Behavior of a humpback whale (*Megaptera novaeangliae*) upon entanglement in a gillnet. *Marine Mammal Science* 15: 211–213.
- Wiley, D. N. Asmutis, R. Pitchford, T. and Gannon, D. 1995. Stranding and mortality of humpback whales, *Megaptera novaeangliae*, in the mid-Atlantic and southeast regions of the United States. *Fishery Bulletin* 93: 196–205.

ENDNOTE

¹Sometimes referred to as western-rigged trawlers.

²Sometimes referred to as eastern-rigged trawlers.

³Northeast Fisheries Science Center, Woods Hole, MA 02543

⁴Nautical Technologies Ltd. 217 Burleigh Road, Bangor, Maine 04401.

⁵Charles Mayo (Center for Coastal Studies, Provincetown, MA 02657) first proposed this method of determining risk for use by the Atlantic Large Whale Take Reduction Team (ALWTRT). The ALWTRT was tasked by the Department of Commerce with identifying ways to reduce the incidental kill of several species of baleen whales in fixed fishing gear.

⁶The Western Gulf of Maine Closed Area is a year-round ground fishing closure created by the New England Fishery Management Council to recover depleted stocks. Gillnetting, otter trawling and scallop dredging are prohibited within the area.

⁷This area was based a concentration of right whales observed on a single survey in March 2002. Excluding those sightings, the highest winter concentration of baleen whales occurred on the southwest corner of Stellwagen Bank (0.13–0.16 whale/km²).



The Commonwealth of Massachusetts

Division of Marine Fisheries

251 Causeway Street, Suite 400, Boston, MA 02114
p: (617) 626-1520 | f: (617) 626-1509
www.mass.gov/marinefisheries



CHARLES D. BAKER
Governor

KARYN E. POLITO
Lt. Governor

BETHANY A. CARD
Secretary

RONALD S. AMIDON
Commissioner

DANIEL J. MCKIERNAN
Director

December 12, 2022

Michael Pentony
Regional Administrator
NOAA Fisheries Greater Atlantic Regional Fisheries Office
55 Great Republic Drive
Gloucester, MA 01930

Dear Mr. Pentony,

I am writing you to share two concerns I have regarding the federal coordination of the Atlantic Large Whale Take Reduction Plan (ALWTRP) rule-making to reduce the risk of serious injury and mortality to the North Atlantic right whale (“NARW”). I hope you can consider and address these concerns this winter and as ALWTRP rule making progresses over the course of the next two years.

1. Spatial Gaps Between State and Federal Trap Gear Closures for the Massachusetts Restricted Area

I wrote to you on this subject on January 7, 2022. In this letter, I expressed my concerns regarding the seasonal entanglement risk for the NARW in the EEZ west of Stellwagen Bank and informed NOAA Fisheries there is a portion of federal waters within the Gulf of Maine that remains open to trap fishing and the use of persistent buoy lines and is sandwiched between Massachusetts’ February 1 – May 15 Commercial Trap Gear Closure to Protect Right Whales [322 CMR 12.04] and the federal February 1 – April 30 Massachusetts Restricted Area Closure.

This spatial gap between state and federal closures poses a substantial and unnecessary entanglement risk to NARW. Having this near-shore area remain open to trap gear fishing and persistent buoy lines when adjacent state and federal waters are closed creates an opportunity for federally permitted vessels to fish or store buoyed trap gear in the area. Since 2018, sightings data indicate that NARW are being increasingly observed in state and federal waters in Massachusetts Bay and north towards the New Hampshire coastline. The combined effect is a documentable seasonal co-occurrence between NARW and buoyed trap gear, particularly during April and May when right whales begin to seasonally migrate out of Cape Cod Bay (Figure 1). I am concerned this continued overlap of buoyed trap gear with aggregations of NARW could result in an entanglement in waters off Massachusetts’ coast that could threaten the viability of Massachusetts’ fixed gear fisheries moving forward.

NOAA Fisheries was responsive when I raised this issue back in January and you pursued an emergency rule to close the so-called Massachusetts Restricted Area Wedge from April 1 – April 30 in 2022. The Massachusetts Restricted Area Wedge was inclusive of those federal waters west of 70° 30' west longitude between 42° 12' N latitude to the south and 42° 39.77' N latitude to the north. I commend you for taking this important action.




Figure 1. Right whale sightings in 2021 and buoy lines documented in April and May 2021 (CCS data)

During the course of 2022, NOAA Fisheries did not pursue interim rule-making to make permanent the emergency closure of the Massachusetts Restricted Area Wedge. As a result, in 2023, this area will again be open to federally permitted vessels to fish or store buoyed trap gear during the late winter and early spring months. I encourage NOAA Fisheries to consider the entanglement risk posed by spatial gaps in seasonal buoyed trap gear closure coverage. Moreover, I strongly support NOAA Fisheries re-closing the Massachusetts Restricted Area Wedge—similar to this past year—for 2023 and 2024, or until the new ALWTRP rules are implemented.

2. Enhanced Coordination in ALWTRP Rule Making Within NOAA Fisheries and with the Councils

As a result of the recent Boasberg decision, NOAA Fisheries has initiated a two-year rule making process to reduce the risk of NARW entanglements in regulated fisheries by 90% coastwide in order to achieve PBR. This presents a substantial and unprecedented conservation challenge. The breadth of this rule-making endeavor is considerable and it expands across

various fisheries and gear types. Moreover, some potential outcomes may have indirect impacts on fisheries not regulated under the ALWTRP. Accordingly, NOAA Fisheries should enhance coordination regarding ALWTRP rule-making efforts between its Protected Resources Division and its Sustainable Fisheries Division and with the New England and Mid-Atlantic Fishery Management Councils.

There is substantial overlap between what the TRT is discussing and considering and the work being conducted by the Sustainable Fisheries Division. In my experience, there can be a disconnect between the two programs. For instance, there are legacy fisheries—where effort and participation is tightly controlled at the state and federal levels (e.g., lobster trap)—that are required to substantially cut how they conduct their fishing activities to address NARW entanglement risk. Meanwhile, there are limited federal controls on the proliferation of new fixed gear fishing effort (e.g., waved whelk pot, black sea bass pot) in the federal zone that increase the presence of persistent buoy lines in the water column and subsequent risk to NARW. This disconnect complicates management and hurts NOAA Fisheries credibility with stakeholders.

To this point, I was encouraged that staff from the Sustainable Fisheries Division attended the recent ALWTRP industry scoping meeting with the southern New England gillnet fleet. There is overlap between the management of the skate, monkfish, and groundfish fisheries in the region and the management of this gillnet fishery with regards to NARW entanglement risk. Having staff from both divisions present made for a more robust and informed dialogue. More deliberate coordination among NOAA Fisheries staff is necessary and appropriate to comprehensively address the robust challenge the TRT currently faces.

Similarly, there should be vigorous coordination between NOAA Fisheries and the New England and Mid-Atlantic Fishery Management Councils. There are certain management measures that may achieve risk reduction that are outside the purview of the TRT and require Council action. For instance, the southern New England monkfish and skate gillnet fleet expressed interest in addressing latent effort as a means of reducing entanglement risk and this would require the Council to amend the relevant fishery management plans.

However, the most important place for coordination between the ALWTRP rule making process and the Councils is with regards to the potential use of on-demand buoy line systems (“ropeless fishing”) and or alternatively, using only one buoy line on multi-trap trawls. If these types of trap fishing activities are going to be authorized or mandated in the federal zone, there will be a proliferation of trap gear without surface markings. This substantially increases the likelihood of gear conflicts and poses a significant additional safety risk to commercial fishers whose gear may become hung-up on this unmarked gear. To avoid such gear conflicts, I anticipate the New England and Mid-Atlantic Fishery Management Councils are going to have to take actions across a variety of federally managed fisheries prosecuted by mobile gear to require vessels be equipped with technology to determine the presence unbuoyed trap gear.

On a similar but unrelated matter, the draft Sturgeon Action Plan to reduce bycatch in gillnet fisheries warrants similar coordination across NOAA Fisheries, and with the Councils, as well as the Atlantic States Marine Fisheries Commission.

Thank you for your time reviewing my concerns. Please let me know if there is any way for the Massachusetts Division of Marine Fisheries to further assist NOAA Fisheries in meeting this critical and considerable management challenge.

Best regards,

A handwritten signature in black ink, reading "Daniel J. McKiernan". The signature is fluid and cursive, with the first name "Daniel" and last name "McKiernan" clearly legible.

Daniel J. McKiernan, Director
Massachusetts Division of Marine Fisheries

cc: Massachusetts Marine Fisheries Advisory Commission; Massachusetts Department of Fish and Game; New England Fishery Management Council; Mid-Atlantic Fishery Management Council

Enc: January 7, 2022 letter from DMF to GARFO



The Commonwealth of Massachusetts

Division of Marine Fisheries

251 Causeway Street, Suite 400, Boston, MA 02114
p: (617) 626-1520 | f: (617) 626-1509
www.mass.gov/marinefisheries



CHARLES D. BAKER
Governor

KARYN E. POLITO
Lt. Governor

BETHANY A. CARD
Secretary

RONALD S. AMIDON
Commissioner

DANIEL J. MCKIERNAN
Director

December 12, 2022

Michael Pentony
Regional Administrator
NOAA Fisheries Greater Atlantic Regional Fisheries Office
55 Great Republic Drive
Gloucester, MA 01930

Dear Mr. Pentony,

I am writing you to share two concerns I have regarding the federal coordination of the Atlantic Large Whale Take Reduction Plan (ALWTRP) rule-making to reduce the risk of serious injury and mortality to the North Atlantic right whale (“NARW”). I hope you can consider and address these concerns this winter and as ALWTRP rule making progresses over the course of the next two years.

1. Spatial Gaps Between State and Federal Trap Gear Closures for the Massachusetts Restricted Area

I wrote to you on this subject on January 7, 2022. In this letter, I expressed my concerns regarding the seasonal entanglement risk for the NARW in the EEZ west of Stellwagen Bank and informed NOAA Fisheries there is a portion of federal waters within the Gulf of Maine that remains open to trap fishing and the use of persistent buoy lines and is sandwiched between Massachusetts’ February 1 – May 15 Commercial Trap Gear Closure to Protect Right Whales [322 CMR 12.04] and the federal February 1 – April 30 Massachusetts Restricted Area Closure.

This spatial gap between state and federal closures poses a substantial and unnecessary entanglement risk to NARW. Having this near-shore area remain open to trap gear fishing and persistent buoy lines when adjacent state and federal waters are closed creates an opportunity for federally permitted vessels to fish or store buoyed trap gear in the area. Since 2018, sightings data indicate that NARW are being increasingly observed in state and federal waters in Massachusetts Bay and north towards the New Hampshire coastline. The combined effect is a documentable seasonal co-occurrence between NARW and buoyed trap gear, particularly during April and May when right whales begin to seasonally migrate out of Cape Cod Bay (Figure 1). I am concerned this continued overlap of buoyed trap gear with aggregations of NARW could result in an entanglement in waters off Massachusetts’ coast that could threaten the viability of Massachusetts’ fixed gear fisheries moving forward.

NOAA Fisheries was responsive when I raised this issue back in January and you pursued an emergency rule to close the so-called Massachusetts Restricted Area Wedge from April 1 – April 30 in 2022. The Massachusetts Restricted Area Wedge was inclusive of those federal waters west of 70° 30' west longitude between 42° 12' N latitude to the south and 42° 39.77' N latitude to the north. I commend you for taking this important action.




Figure 1. Right whale sightings in 2021 and buoy lines documented in April and May 2021 (CCS data)

During the course of 2022, NOAA Fisheries did not pursue interim rule-making to make permanent the emergency closure of the Massachusetts Restricted Area Wedge. As a result, in 2023, this area will again be open to federally permitted vessels to fish or store buoyed trap gear during the late winter and early spring months. I encourage NOAA Fisheries to consider the entanglement risk posed by spatial gaps in seasonal buoyed trap gear closure coverage. Moreover, I strongly support NOAA Fisheries re-closing the Massachusetts Restricted Area Wedge—similar to this past year—for 2023 and 2024, or until the new ALWTRP rules are implemented.

2. Enhanced Coordination in ALWTRP Rule Making Within NOAA Fisheries and with the Councils

As a result of the recent Boasberg decision, NOAA Fisheries has initiated a two-year rule making process to reduce the risk of NARW entanglements in regulated fisheries by 90% coastwide in order to achieve PBR. This presents a substantial and unprecedented conservation challenge. The breadth of this rule-making endeavor is considerable and it expands across

various fisheries and gear types. Moreover, some potential outcomes may have indirect impacts on fisheries not regulated under the ALWTRP. Accordingly, NOAA Fisheries should enhance coordination regarding ALWTRP rule-making efforts between its Protected Resources Division and its Sustainable Fisheries Division and with the New England and Mid-Atlantic Fishery Management Councils.

There is substantial overlap between what the TRT is discussing and considering and the work being conducted by the Sustainable Fisheries Division. In my experience, there can be a disconnect between the two programs. For instance, there are legacy fisheries—where effort and participation is tightly controlled at the state and federal levels (e.g., lobster trap)—that are required to substantially cut how they conduct their fishing activities to address NARW entanglement risk. Meanwhile, there are limited federal controls on the proliferation of new fixed gear fishing effort (e.g., waved whelk pot, black sea bass pot) in the federal zone that increase the presence of persistent buoy lines in the water column and subsequent risk to NARW. This disconnect complicates management and hurts NOAA Fisheries credibility with stakeholders.

To this point, I was encouraged that staff from the Sustainable Fisheries Division attended the recent ALWTRP industry scoping meeting with the southern New England gillnet fleet. There is overlap between the management of the skate, monkfish, and groundfish fisheries in the region and the management of this gillnet fishery with regards to NARW entanglement risk. Having staff from both divisions present made for a more robust and informed dialogue. More deliberate coordination among NOAA Fisheries staff is necessary and appropriate to comprehensively address the robust challenge the TRT currently faces.

Similarly, there should be vigorous coordination between NOAA Fisheries and the New England and Mid-Atlantic Fishery Management Councils. There are certain management measures that may achieve risk reduction that are outside the purview of the TRT and require Council action. For instance, the southern New England monkfish and skate gillnet fleet expressed interest in addressing latent effort as a means of reducing entanglement risk and this would require the Council to amend the relevant fishery management plans.

However, the most important place for coordination between the ALWTRP rule making process and the Councils is with regards to the potential use of on-demand buoy line systems (“ropeless fishing”) and or alternatively, using only one buoy line on multi-trap trawls. If these types of trap fishing activities are going to be authorized or mandated in the federal zone, there will be a proliferation of trap gear without surface markings. This substantially increases the likelihood of gear conflicts and poses a significant additional safety risk to commercial fishers whose gear may become hung-up on this unmarked gear. To avoid such gear conflicts, I anticipate the New England and Mid-Atlantic Fishery Management Councils are going to have to take actions across a variety of federally managed fisheries prosecuted by mobile gear to require vessels be equipped with technology to determine the presence unbuoyed trap gear.

On a similar but unrelated matter, the draft Sturgeon Action Plan to reduce bycatch in gillnet fisheries warrants similar coordination across NOAA Fisheries, and with the Councils, as well as the Atlantic States Marine Fisheries Commission.

Thank you for your time reviewing my concerns. Please let me know if there is any way for the Massachusetts Division of Marine Fisheries to further assist NOAA Fisheries in meeting this critical and considerable management challenge.

Best regards,

A handwritten signature in black ink, reading "Daniel J. McKiernan". The signature is fluid and cursive, with a long horizontal stroke at the end.

Daniel J. McKiernan, Director
Massachusetts Division of Marine Fisheries

cc: Massachusetts Marine Fisheries Advisory Commission; Massachusetts Department of Fish and Game; New England Fishery Management Council; Mid-Atlantic Fishery Management Council

Enc: January 7, 2022 letter from DMF to GARFO



The Commonwealth of Massachusetts

Division of Marine Fisheries

251 Causeway Street, Suite 400, Boston, MA 02114
p: (617) 626-1520 | f: (617) 626-1509
www.mass.gov/marinefisheries



CHARLES D. BAKER
Governor

KARYN E. POLITO
Lt. Governor

KATHLEEN A. THEOHARIDES
Secretary

RONALD S. AMIDON
Commissioner

DANIEL J. MCKIERNAN
Director

January 7, 2022

Michael Pentony
Regional Administrator
NOAA Fisheries GARFO
55 Great Republic Drive
Gloucester, MA 01930

RE: Seasonal entanglement risk for North Atlantic Right Whales in the EEZ west of Stellwagen Bank

Dear Mr. Pentony,

I am writing to inform you of an emerging entanglement risk to North Atlantic right whales (NARW) that occurs in a certain zone of federal waters sandwiched between the state and federal closures.

As you are aware, NOAA Fisheries created the Massachusetts Restricted Area (MRA) in 2015 to reduce the risk of entanglement risk to the large aggregations of NARW that occur there seasonally. This closure to fixed fishing gear included MA state waters within Cape Cod Bay and adjacent federal waters around Stellwagen Bank from February 1st through April 30th of each year. DMF immediately created analogous state regulations closing the area to fixed fishing gear.

Since 2016, DMF has also added dynamic management to the state waters portion of the MRA by extending the closure into the month of May when aerial surveillance shows that right whales remain present. In addition to this action, since the beginning of the closure, DMF has engaged in efforts, with assistance from the Massachusetts Environmental Police, to retrieve abandoned gear in the closure annually to ensure that the entanglement risk to right whales is effective as intended.

Since the advent of the MRA closure in 2015, seasonal usage of state and federal waters outside of Cape Cod Bay increased in certain areas and times where fixed gear fishing was allowed. Recent sighting data indicate that NARW stay for a longer time period than they have historically, and these whales are increasingly observed in state and adjacent federal waters in Massachusetts Bay and north to the NH state line. These changes in distribution increased the entanglement risk to NARW along the MA coastal waters. In response to these changes in entanglement risk, as well as continued declines in the population status of NARW, in 2021 DMF closed MA state waters from southeastern Cape Cod north the NH border to lobster fishing from February 1st to May 15th (Figure 1).




Figure 1. DMF trap/gear closure, February 1 – May 15

The National Marine Fisheries Service then mirrored the northern extension of the closure, known as Massachusetts North Restricted Area, in their Phase 1 amendment to the Atlantic Large Whale Take Reduction Plan in September of 2021. The Massachusetts North closure only runs through April 30 each year under the federal plan (Figure 2).




Figure 2. Map of Massachusetts Restricted Area

The increasing presence of NARW in these northern areas is not exclusive to state waters. In recent years, aerial surveillance conducted by the Center for Coastal Studies (CCS) has documented the presence of right whales in both open and closed portions of the waters north of Cape Cod Bay. The map below depicting gear and whales from 2018 demonstrates the necessity for DMF's northern extension of the state waters closure implemented in 2021 (Figure 3).




Figure 3. Right whale sightings in 2018 and buoy lines documented on April 23, 2018

However, the implementation of the Massachusetts North Restricted Area has created a gap between the closed areas between state waters of Massachusetts Bay and the northern federal waters portion of the original Massachusetts Restricted Area (Figure 2 and 4). Federally permitted vessels can continue to fish with persistent buoy lines in these areas adjacent to MA state waters during the closure period, and this area lies beyond the jurisdiction of the Commonwealth.




Figure 4. Right whale sightings in 2021 and buoy lines documented in April and May 2021

Given the current dire status of the NARW population and the need for continued reductions in entanglement risk we wanted to ensure that NOAA Fisheries was aware of this issue. We feel that continued overlap of persistent buoy lines with aggregations of NARW pose an entanglement threat and we are concerned that any future NARW entanglement in waters off the Massachusetts coast could threaten the opportunity of MA-based fishers to participate in fixed gear fisheries.

Sincerely,

Daniel J. McKiernan, Director

CC: Marine Fisheries Advisory Commission



The Commonwealth of Massachusetts

Division of Marine Fisheries

251 Causeway Street, Suite 400, Boston, MA 02114

p: (617) 626-1520 | f: (617) 626-1509

www.mass.gov/marinefisheries



CHARLES D. BAKER
Governor

KARYN E. POLITO
Lt. Governor

BETHANY A. CARD
Secretary

RONALD S. AMIDON
Commissioner

DANIEL J. MCKIERNAN
Director

January 4, 2023

Michael Pentony
Regional Administrator
NOAA Fisheries Greater Atlantic Regional Fisheries Office
55 Great Republic Drive
Gloucester, MA 01930

Dear Michael,

On December 12 2022, I made a request to NOAA Fisheries to repeat the seasonal closure enacted last winter to close the Massachusetts Restricted Area Wedge (“wedge”)—a spatial gap between the state and federal seasonal trap gear closures in Massachusetts Bay. This area west of Stellwagen Bank is a magnet for trap gear for fishers who opt not to bring their gear home for the season and instead move gear out of the Massachusetts Seasonal Trap Gear Closure and the Massachusetts Restricted Area and into the wedge. The co-occurrence of this gear with the seasonal presence of right whales in the wedge, particularly during the spring months, creates an elevated risk of entanglement.

Your agency’s action to close this area was much appreciated by the Commonwealth, although the timing of your 2022 action was delayed and began on April 1—a full two months after the adjacent areas were already closed. A February 1 start date for this closure would have been preferred to rid the area of persistent buoy lines and further reduce entanglement risk when right whales are present. I had anticipated after last winter’s closure of the area that NOAA would be in a position to enact this as a final rule. Unfortunately, due to a variety of circumstances, a permanent closure to the wedge has not transpired and it has become an ephemeral rule necessitating annual renewal.

Much has transpired over the past few weeks with the enactment of the Congressional Omnibus Appropriations action affecting your agency’s ability to further regulate the American Lobster and Jonah Crab fisheries through modifications to the Take Reduction Plan. I suspect you and the policy and legal experts are still determining the impacts of the Congressional action on the agency’s action plan to conserve right whales. With all this considered, as well as the intense work NOAA Fisheries and DMF have dedicated over the past several years to the development of additional risk reduction measures, I do not want to see this issue fall through the bureaucratic cracks.


DMF believes this closure is still warranted and should be pursued by NOAA Fisheries. In fact, the budgetary language provides authorization for you to continue with this action:

(b) EXCEPTION.—The provisions of subsection (a) shall not apply to an existing emergency rule, or any action taken to extend or make final an emergency rule that is in place on the date of enactment of this Act, affecting lobster and Jonah crab.

You and NOAA Fisheries staff have DMF's full support for a permanent adoption of a closure of the wedge annually from February through May, or as long as the adjacent areas remain closed. Please let me know if you believe this rulemaking is warranted and executable given the circumstances created by the Omnibus Appropriations action. We are already receiving inquiries from lobster fishers who are making business decisions to remove or leave gear in the wedge.

In closing, if you need any assistance on this matter my staff and I are available to assist.

Sincerely,

A handwritten signature in black ink, reading "Daniel J. McKiernan". The signature is fluid and cursive, with a long horizontal stroke at the end.

Daniel J. McKiernan, Director

Cc: Janet Coit, Assistant Administrator for Fisheries
Samuel Rauch, Deputy Assistant Administrator
Marine Fisheries Advisory Commission members

Chapter 3 Appendices

Appendix 3.1 Community Profiles

Appendix Community Profiles

Rockport, MA

Where is Rockport located?

Rockport is a town with a population of 6,952 and classified by the census as falling within an urbanized area. Rural to urban is really a continuum. Increasing urbanization indicates that a community has more jobs overall, more kinds of jobs, and more services like hospitals, social workers and job training centers. However, increasing urbanization can also mean greater pressure to transform working waterfronts for alternative uses, such as hotels or tourist shops.




Involvement in Fisheries

What species are landed in Rockport?

The landings associated with a fishing community tell us what species are important to that community. The diversity of species caught also is indicative of a community's ability to adapt to changing environmental conditions (e.g. populations of specific fish stocks) or changes in fishing regulations that restrict access to resources.

*Groundfish includes cod, winter fl., witch fl., yellowtail fl., am.plaice, haddock, white hake, redfish, pollock.

**Whiting includes red hake, ocean pout, black whiting, whiting.




What are the characteristics of the fishing vessels in Rockport?

The number of fishing vessels in a given port provides a sense of the scale of fishing in that port. Where a large port may serve as the homeport for hundreds of vessels, a smaller one may only have a handful. The number of vessels also may provide a rough sense of the number of fishing-related jobs (e.g. crew positions, jobs in shoreside industries) available in a given location.

Size also matters. Larger vessels can travel farther offshore and stay out for longer periods more easily than smaller vessels. These differences also affect family life. Smaller dayboat fishermen tend to return home every day whereas fishermen on larger vessels may be away from home for weeks on long and distant fishing expeditions.

Apart from the lobster fleet, smaller boats also tend to catch a broader range of species where their larger counterparts are more specialized (e.g. limited access scallop boats and herring pair trawlers). All these characteristics help illuminate the potential impacts of regulatory changes on a given community.

Number of Vessels by Size (2014)




Demographic Attributes

Educational Attainment

The level of educational attainment in a community is associated with issues important for community development, such as income and poverty levels, unemployment rates, and local participation in community activities.

Educational Attainment



How do people make a living in Rockport?

Just as the range of fish species harvested by town residents speaks to their ability to adapt to environmental change, the diversity in local occupations indicates the ability of a community to adapt to economic changes, including changes in the local fishing economy. Is there one predominant industry, for instance, or is there a range of economic opportunities? How many occupations are available that offer incomes similar to fishing or require skills and education common to the average fisherman? How many jobs are available that would provide a working environment that fishermen would be comfortable with?


Unemployment Rate: 5.2%

National Rate: **7.9%***

The unemployment rate in a community is one indicator of the level of opportunity that may exist for fishermen who lose their jobs to find alternative ways of making a living. The unemployment rate may also indicate the desirability of fishing in the face of other opportunities.

*Source: U.S. Department of Labor, **Bureau of Labor Statistics**

Occupations by Industry



Median Household Income: \$70,625.00

National Average: **\$51,914.00** (2011)

Individuals in Rockport living in poverty: 3.7%

The poverty threshold for an individual is defined by the US Census for 2010 as \$11,139. The percentage of a town's population living under this economic threshold is an indicator of the residents' ability to adjust to loss of income and job opportunities in fishing-related and other local industries.

Age structure of residents


Age structure provides potential indications of many broader community issues and institutions. A large number of older residents may be associated with a retirement community or an out-migration of young people. For many fishing communities, an aging population can indicate gentrification, a process that may affect fishermen's access to the waterfront. In some remote coastal communities, people in their late teens or early twenties may leave to look for work or pursue an education outside of their community. A very large population of young people, on the other hand, may indicate the presence of universities or a military base.

Median age: **51.2**

National median: **37.2**



Population pyramid for Rockport, year 2010

Source: www.census.gov



Ethnicity and Race

These factors give a sense of the cultural context of the community, and the relationship of fishing families and groups to the community in which they live. Is this community racially and ethnically diverse? In the northeast region, ethnic diversity in coastal communities tends to be higher in the Mid Atlantic than in New England, though there are significant exceptions in some fishing ports. Moreover, certain ethnic groups have long been associated with fishing in various specific ports throughout the region.



Language and Marginalization

Fishing regulations can be complex. Documents are rarely translated from English into other languages. Lack of strong English language skills could affect participants' ability to engage effectively in the fisheries management process. While these numbers correspond to the overall community in Rockport they may indicate a population needing assistance in integrating their needs and concerns into the process.

Foreign Born: **3.9%**

National Average: **12.7%**

Speak English less than very well: **1.1%**

National Average: **8.7%**

Social Indicators

Social indicators are quantitative measures that describe the well-being of communities and are used to describe social phenomena over time. Below are a series of indices for Rockport that provide measures of fishing engagement and reliance, and social vulnerability. An index combines variables of interest and are used to evaluate community well-being in terms of social, economic and psychological welfare.

Fishing engagement and reliance indices portray the importance or level of dependence of commercial or recreational fishing to coastal communities. The indices include: Commercial Engagement, Commercial Reliance, Recreational Engagement and Recreational Reliance.

Social vulnerability indices represent social factors that can shape either an individual or community's ability to adapt to change. These factors exist within all communities regardless of the importance of fishing. The indices include: Poverty, Population Composition, and Personal Disruption.

Gentrification Pressure indices characterize those factors that, over time may indicate a threat to the viability of a commercial or recreational working waterfront, including infrastructure. The indices include: Retire Migration, Urban Sprawl, Natural Amenities and Housing Disruption.

The factor scores for each index are normalized so that zero is the mean. Therefore, a higher value implies more engagement or reliance upon fishing or higher social vulnerability or vulnerability to gentrification. Learn more about the [social indicators for fishing communities](#).

Social Indicators



Gloucester, MA

Where is Gloucester located?

Gloucester is a town with a population of 28,789 and classified by the census as falling within an urbanized area. Rural to urban is really a continuum. Increasing urbanization indicates that a community has more jobs overall, more kinds of jobs, and more services like hospitals, social workers and job training centers. However, increasing urbanization can also mean greater pressure to transform working waterfronts for alternative uses, such as hotels or tourist shops.



Involvement in Fisheries


What species are landed in Gloucester?

The landings associated with a fishing community tell us what species are important to that community. The diversity of species caught also is indicative of a community's ability to adapt to changing environmental conditions (e.g. populations of specific fish stocks) or changes in fishing regulations that restrict access to resources.

*Groundfish includes cod, winter fl., witch fl., yellowtail fl., am. plaice, haddock, white hake, redfish, pollock.

**Whiting includes red hake, ocean pout, black whiting, whiting.

Landed Values by Species (2014)



What are the characteristics of the fishing vessels in Gloucester?

The number of fishing vessels in a given port provides a sense of the scale of fishing in that port. Where a large port may serve as the homeport for hundreds of vessels, a smaller one may only have a handful. The number of vessels also may provide a rough sense of the number of fishing-related jobs (e.g. crew positions, jobs in shoreside industries) available in a given location.

Size also matters. Larger vessels can travel farther offshore and stay out for longer periods more easily than smaller vessels. These differences also affect family life. Smaller dayboat fishermen tend to return home every day whereas fishermen on larger vessels may be away from home for weeks on long and distant fishing expeditions.


Apart from the lobster fleet, smaller boats also tend to catch a broader range of species where their larger counterparts are more specialized (e.g. limited access scallop boats and herring pair trawlers). All these characteristics help illuminate the potential impacts of regulatory changes on a given community.

Demographic Attributes


Educational Attainment

The level of educational attainment in a community is associated with issues important for community development, such as income and poverty levels, unemployment rates, and local participation in community activities.

Number of Vessels by Size (2014)



Educational Attainment



How do people make a living in Gloucester?


Just as the range of fish species harvested by town residents speaks to their ability to adapt to environmental change, the diversity in local occupations indicates the ability of a community to adapt to economic changes, including changes in the local fishing economy. Is there one predominant industry, for instance, or is there a range of economic opportunities? How many occupations are available that offer incomes similar to fishing or require skills and education common to the average fisherman? How many jobs are available that would provide a working environment that fishermen would be comfortable with?

Unemployment Rate: **4%**

National Rate: **7.9%***

The unemployment rate in a community is one indicator of the level of opportunity that may exist for fishermen who lose their jobs to find alternative ways of making a living. The unemployment rate may also indicate the desirability of fishing in the face of other opportunities.

*Source: U.S. Department of Labor, [Bureau of Labor Statistics](#)



Median Household Income: **\$60,506.00**

National Average: **\$51,914.00** (2011)

Individuals in Gloucester living in poverty: **7.8%**

The poverty threshold for an individual is defined by the US Census for 2010 as \$11,139. The percentage of a town's population living under this economic threshold is an indicator of the residents' ability to adjust to loss of income and job opportunities in fishing-related and other local industries.

Age structure of residents


Age structure provides potential indications of many broader community issues and institutions. A large number of older residents may be associated with a retirement community or an out-migration of young people. For many fishing communities, an aging population can indicate gentrification, a process that may affect fishermen's access to the waterfront. In some remote coastal communities, people in their late teens or early twenties may leave to look for work or pursue an education outside of their community. A very large population of young people, on the other hand, may indicate the presence of universities or a military base.

Median age: **46.4**

National median: **37.2**

Population pyramid for Gloucester, year 2010


Source: www.census.gov




Ethnicity and Race

These factors give a sense of the cultural context of the community, and the relationship of fishing families and groups to the community in which they live. Is this community racially and ethnically diverse? In the northeast region, ethnic diversity in coastal communities tends to be higher in the Mid Atlantic than in New England, though there are significant exceptions in some fishing ports. Moreover, certain ethnic groups have long been associated with fishing in various specific ports throughout the region.

Ethnicity



Race



Language and Marginalization

Fishing regulations can be complex. Documents are rarely translated from English into other languages. Lack of strong English language skills could affect participants' ability to engage effectively in the fisheries management process. While these numbers correspond to the overall community in Gloucester they may indicate a population needing assistance in integrating their needs and concerns into the process.

Foreign Born: **7%**

National Average: **12.7%**

Speak English less than very well: **4%**

National Average: **8.7%**

Social Indicators

Social indicators are quantitative measures that describe the well-being of communities and are used to describe social phenomena over time. Below are a series of indices for Gloucester that provide measures of fishing engagement and reliance, and social vulnerability. An index combines variables of interest and are used to evaluate community well-being in terms of social, economic and psychological welfare.

Fishing engagement and reliance indices portray the importance or level of dependence of commercial or recreational fishing to coastal communities. The indices include: Commercial Engagement, Commercial Reliance, Recreational Engagement and Recreational Reliance.

Social vulnerability indices represent social factors that can shape either an individual or community's ability to adapt to change. These factors exist within all communities regardless of the importance of fishing. The indices include: Poverty, Population Composition, and Personal Disruption.

Gentrification Pressure indices characterize those factors that, over time may indicate a threat to the viability of a commercial or recreational working waterfront, including infrastructure. The indices include: Retire Migration, Urban Sprawl, Natural Amenities and Housing Disruption.

The factor scores for each index are normalized so that zero is the mean. Therefore, a higher value implies more engagement or reliance upon fishing or higher social vulnerability or vulnerability to gentrification. Learn more about the [social indicators for fishing communities](#).

Social Indicators



Boston, MA

Where is Boston located?

Boston is a town with a population of 617,594 and classified by the census as falling within an urbanized area. Rural to urban is really a continuum. Increasing urbanization indicates that a community has more jobs overall, more kinds of jobs, and more services like hospitals, social workers and job training centers. However, increasing urbanization can also mean greater pressure to transform working waterfronts for alternative uses, such as hotels or tourist shops.




Involvement in Fisheries

What species are landed in Boston?

The landings associated with a fishing community tell us what species are important to that community. The diversity of species caught also is indicative of a community's ability to adapt to changing environmental conditions (e.g. populations of specific fish stocks) or changes in fishing regulations that restrict access to resources.

*Groundfish includes cod, winter fl., witch fl., yellowtail fl., am. plaice, haddock, white hake, redfish, pollock.

**Whiting includes red hake, ocean pout, black whiting, whiting.




What are the characteristics of the fishing vessels in Boston?

The number of fishing vessels in a given port provides a sense of the scale of fishing in that port. Where a large port may serve as the homeport for hundreds of vessels, a smaller one may only have a handful. The number of vessels also may provide a rough sense of the number of fishing-related jobs (e.g. crew positions, jobs in shoreside industries) available in a given location.

Size also matters. Larger vessels can travel farther offshore and stay out for longer periods more easily than smaller vessels. These differences also affect family life. Smaller dayboat fishermen tend to return home every day whereas fishermen on larger vessels may be away from home for weeks on long and distant fishing expeditions.

Apart from the lobster fleet, smaller boats also tend to catch a broader range of species where their larger counterparts are more specialized (e.g. limited access scallop boats and herring pair trawlers). All these characteristics help illuminate the potential impacts of regulatory changes on a given community.

Number of Vessels by Size (2014)




Demographic Attributes

Educational Attainment

The level of educational attainment in a community is associated with issues important for community development, such as income and poverty levels, unemployment rates, and local participation in community activities.

Educational Attainment



How do people make a living in Boston?

Just as the range of fish species harvested by town residents speaks to their ability to adapt to environmental change, the diversity in local occupations indicates the ability of a community to adapt to economic changes, including changes in the local fishing economy. Is there one predominant industry, for instance, or is there a range of economic opportunities? How many occupations are available that offer incomes similar to fishing or require skills and education common to the average fisherman? How many jobs are available that would provide a working environment that fishermen would be comfortable with?


Unemployment Rate: 6.3%

National Rate: 7.9%*

The unemployment rate in a community is one indicator of the level of opportunity that may exist for fishermen who lose their jobs to find alternative ways of making a living. The unemployment rate may also indicate the desirability of fishing in the face of other opportunities.

*Source: U.S. Department of Labor, Bureau of Labor Statistics

Occupations by Industry



Median Household Income: \$50,684.00

National Average: \$51,914.00 (2011)

Individuals in Boston living in poverty: 21.2 % The poverty threshold for an individual is defined by the US Census for 2010 as \$11,139. The percentage of a town's population living under this economic threshold is an indicator of the residents' ability to adjust to loss of income and job opportunities in fishing-related and other local industries.

Age structure of residents

Age structure provides potential indications of many broader community issues and institutions. A large number of older residents may be associated with a retirement community or an out-migration of young people. For many fishing communities, an aging population can indicate gentrification, a process that may affect fishermen's access to the waterfront. In some remote coastal communities, people in their late teens or early twenties may leave to look for work or pursue an education outside of their community. A very large population of young people, on the other hand, may indicate the presence of universities or a military base.

Median age: **30.8**

National median: **37.2**

Population pyramid for Boston, year 2010


Source: www.census.gov




Ethnicity and Race

These factors give a sense of the cultural context of the community, and the relationship of fishing families and groups to the community in which they live. Is this community racially and ethnically diverse? In the northeast region, ethnic diversity in coastal communities tends to be higher in the Mid Atlantic than in New England, though there are significant exceptions in some fishing ports. Moreover, certain ethnic groups have long been associated with fishing in various specific ports throughout the region.

Ethnicity



Race



Language and Marginalization

Fishing regulations can be complex. Documents are rarely translated from English into other languages. Lack of strong English language skills could affect participants' ability to engage effectively in the fisheries management process. While these numbers correspond to the overall community in Portsmouth they may indicate a population needing assistance in integrating their needs and concerns into the process.

Foreign Born: **27.2%**
National Average: **12.7%**
Speak English less than very well: **16.8%**
National Average: **8.7%**

Social Indicators

Social indicators are quantitative measures that describe the well-being of communities and are used to describe social phenomena over time. Below are a series of indices for Portsmouth that provide measures of fishing engagement and reliance, and social vulnerability. An index combines variables of interest and are used to evaluate community well-being in terms of social, economic and psychological welfare.

Fishing engagement and reliance indices portray the importance or level of dependence of commercial or recreational fishing to coastal communities. The indices include: Commercial Engagement, Commercial Reliance, Recreational Engagement and Recreational Reliance.

Social vulnerability indices represent social factors that can shape either an individual or community's ability to adapt to change. These factors exist within all communities regardless of the importance of fishing. The indices include: Poverty, Population Composition, and Personal Disruption.

Gentrification Pressure indices characterize those factors that, over time may indicate a threat to the viability of a commercial or recreational working waterfront, including infrastructure. The indices include: Retire Migration, Urban Sprawl, Natural Amenities and Housing Disruption.

The factor scores for each index are normalized so that zero is the mean. Therefore, a higher value implies more engagement or reliance upon fishing or higher social vulnerability or vulnerability to gentrification. Learn more about the [social indicators for fishing communities](#).

Social Indicators



Cohasset, MA

Where is Cohasset located?

Cohasset is a town with a population of 7,542 and classified by the census as falling within an urbanized area. Rural to urban is really a continuum. Increasing urbanization indicates that a community has more jobs overall, more kinds of jobs, and more services like hospitals, social workers and job training centers. However, increasing urbanization can also mean greater pressure to transform working waterfronts for alternative uses, such as hotels or tourist shops.



Involvement in Fisheries


What species are landed in Cohasset?

The landings associated with a fishing community tell us what species are important to that community. The diversity of species caught also is indicative of a community's ability to adapt to changing environmental conditions (e.g. populations of specific fish stocks) or changes in fishing regulations that restrict access to resources.

*Groundfish includes cod, winter fl., witch fl., yellowtail fl., am. plaice, haddock, white hake, redfish, pollock.

**Whiting includes red hake, ocean pout, black whiting, whiting.

Landed Values by Species (2014)



Number of Vessels by Size (2014)

What are the characteristics of the fishing vessels in Cohasset?

The number of fishing vessels in a given port provides a sense of the scale of fishing in that port. Where a large port may serve as the homeport for hundreds of vessels, a smaller one may only have a handful. The number of vessels also may provide a rough sense of the number of fishing-related jobs (e.g. crew positions, jobs in shoreside industries) available in a given location.

Size also matters. Larger vessels can travel farther offshore and stay out for longer periods more easily than smaller vessels. These differences also affect family life. Smaller dayboat fishermen tend to return home every day whereas fishermen on larger vessels may be away from home for weeks on long and distant fishing expeditions.


Apart from the lobster fleet, smaller boats also tend to catch a broader range of species where their larger counterparts are more specialized (e.g. limited access scallop boats and herring pair trawlers). All these characteristics have the potential to moderate the potential impacts of regulatory changes on a given community.

Demographic Attributes


Educational Attainment

The level of educational attainment in a community is associated with issues important for community development, such as income and poverty levels, unemployment rates, and local participation in community activities.

Number of Vessels by Size (2014)



Educational Attainment



How do people make a living in Cohasset?

Just as the range of fish species harvested by town residents speaks to their ability to adapt to environmental change, the diversity in local occupations indicates the ability of a community to adapt to economic changes, including changes in the local fishing economy. Is there one predominant industry, for instance, or is there a range of economic opportunities? How many occupations are available that offer incomes similar to fishing or require skills and education common to the average fisherman? How many jobs are available that would provide a working environment that fishermen would be comfortable with?


Unemployment Rate: **3.1%**

National Rate: **7.9%***

The unemployment rate in a community is one indicator of the level of opportunity that may exist for fishermen who lose their jobs to find alternative ways of making a living. The unemployment rate may also indicate the desirability of fishing in the face of other opportunities.

*Source: U.S. Department of Labor, [Bureau of Labor Statistics](#)

Occupations by Industry



Median Household Income: **\$114,214.00**

National Average: **\$51,914.00** (2011)

Individuals in Cohasset living in poverty: **1.2 %**

The poverty threshold for an individual is defined by the US Census for 2010 as \$11,139. The percentage of a town's population living under this economic threshold is an indicator of the residents' ability to adjust to loss of income and job opportunities in fishing-related and other local industries.

Age structure of residents


Age structure provides potential indications of many broader community issues and institutions. A large number of older residents may be associated with a retirement community or an out-migration of young people. For many fishing communities, an aging population can indicate gentrification, a process that may affect fishermen's access to the waterfront. In some remote coastal communities, people in their late teens or early twenties may leave to look for work or pursue an education outside of their community. A very large population of young people, on the other hand, may indicate the presence of universities or a military base.

Median age: **43.6**

National median: **37.2**

Population pyramid for Cohasset, year 2010


Source: www.census.gov




Ethnicity and Race

These factors give a sense of the cultural context of the community, and the relationship of fishing families and groups to the community in which they live. Is this community racially and ethnically diverse? In the northeast region, ethnic diversity in coastal communities tends to be higher in the Mid Atlantic than in New England, though there are significant exceptions in some fishing ports. Moreover, certain ethnic groups have long been associated with fishing in various specific ports throughout the region.

Ethnicity



Race



Language and Marginalization

Fishing regulations can be complex. Documents are rarely translated from English into other languages. Lack of strong English language skills could affect participants' ability to engage effectively in the fisheries management process. While these numbers correspond to the overall community in Cohasset they may indicate a population needing assistance in integrating their needs and concerns into the process.

Foreign Born: **3.9%**
National Average: **12.7%**

Speak English less than very well: **0.9%**
National Average: **8.7%**

Social Indicators

Social indicators are quantitative measures that describe the well-being of communities and are used to describe social phenomena over time. Below are a series of indices for Portsmouth that provide measures of fishing engagement and reliance, and social vulnerability. An index combines variables of interest and are used to evaluate community well-being in terms of social, economic and psychological welfare.


Fishing engagement and reliance indices portray the importance or level of dependence of commercial or recreational fishing to coastal communities. The indices include: Commercial Engagement, Commercial Reliance, Recreational Engagement and Recreational Reliance.

Social vulnerability indices represent social factors that can shape either an individual or community's ability to adapt to change. These factors exist within all communities regardless of the importance of fishing. The indices include: Poverty, Population Composition, and Personal Disruption.

Gentrification Pressure indices characterize those factors that, over time may indicate a threat to the viability of a commercial or recreational working waterfront, including infrastructure. The indices include: Retire Migration, Urban Sprawl, Natural Amenities and Housing Disruption.

The factor scores for each index are normalized so that zero is the mean. Therefore, a higher value implies more engagement or reliance upon fishing or higher social vulnerability or vulnerability to gentrification. Learn more about the [social indicators for fishing communities](#).

Social Indicators



Scituate, MA

Where is Scituate located?

Scituate is a town with a population of 18,133 and classified by the census as falling within an urbanized area. Rural to urban is really a continuum. Increasing urbanization indicates that a community has more jobs overall, more kinds of jobs, and more services like hospitals, social workers and job training centers. However, increasing urbanization can also mean greater pressure to transform working waterfronts for alternative uses, such as hotels or tourist shops.




Involvement in Fisheries

What species are landed in Scituate?

The landings associated with a fishing community tell us what species are important to that community. The diversity of species caught also is indicative of a community's ability to adapt to changing environmental conditions (e.g. populations of specific fish stocks) or changes in fishing regulations that restrict access to resources.

*Groundfish includes cod, winter fl., witch fl., yellowtail fl., am. plaice, haddock, white hake, redfish, pollock.

**Whiting includes red hake, ocean pout, black whiting, whiting.



What are the characteristics of the fishing vessels in Scituate?

The number of fishing vessels in a given port provides a sense of the scale of fishing in that port. Where a large port may serve as the homeport for hundreds of vessels, a smaller one may only have a handful. The number of vessels also may provide a rough sense of the number of fishing-related jobs (e.g. crew positions, jobs in shoreside industries) available in a given location.

Size also matters. Larger vessels can travel farther offshore and stay out for longer periods more easily than smaller vessels. These differences also affect family life. Smaller dayboat fishermen tend to return home every day whereas fishermen on larger vessels may be away from home for weeks on long and distant fishing expeditions.


Apart from the lobster fleet, smaller boats also tend to catch a broader range of species where their larger counterparts are more specialized (e.g. limited access scallop boats and herring pair trawlers). All these characteristics help illuminate the potential impacts of regulatory changes on a given community.

Demographic Attributes


Educational Attainment

The level of educational attainment in a community is associated with issues important for community development, such as income and poverty levels, unemployment rates, and local participation in community activities.

Number of Vessels by Size (2014)



Educational Attainment



How do people make a living in Scituate?

Just as the range of fish species harvested by town residents speaks to their ability to adapt to environmental change, the diversity in local occupations indicates the ability of a community to adapt to economic changes, including changes in the local fishing economy. Is there one predominant industry, for instance, or is there a range of economic opportunities? How many occupations are available that offer incomes similar to fishing or require skills and education common to the average fisherman? How many jobs are available that would provide a working environment that fishermen would be comfortable with?


Unemployment Rate: 4%

National Rate: 7.9%*

The unemployment rate in a community is one indicator of the level of opportunity that may exist for fishermen who lose their jobs to find alternative ways of making a living. The unemployment rate may also indicate the desirability of fishing in the face of other opportunities.

*Source: U.S. Department of Labor, Bureau of Labor Statistics

Occupations by Industry



Median Household Income: **\$86,723.00**

National Average: **\$51,914.00** (2011)

Individuals in Scituate living in poverty: **3.1%**

The poverty threshold for an individual is defined by the US Census for 2010 as \$11,139. The percentage of a town's population living under this economic threshold is an indicator of the residents' ability to adjust to loss of income and job opportunities in fishing-related and other local industries.

Age structure of residents

Age structure provides potential indications of many broader community issues and institutions. A large number of older residents may be associated with a retirement community or an out-migration of young people. For many fishing communities, an aging population can indicate gentrification, a process that may affect fishermen's access to the waterfront. In some remote coastal communities, people in their late teens or early twenties may leave to look for work or pursue an education outside of their community. A very large population of young people, on the other hand, may indicate the presence of universities or a military base.

Median age: **45.1**

National median: **37.2**

Population pyramid for Scituate, year 2010


Source: www.census.gov




Ethnicity and Race

These factors give a sense of the cultural context of the community, and the relationship of fishing families and groups to the community in which they live. Is this community racially and ethnically diverse? In the northeast region, ethnic diversity in coastal communities tends to be higher in the Mid Atlantic than in New England, though there are significant exceptions in some fishing ports. Moreover, certain ethnic groups have long been associated with fishing in various specific ports throughout the region.

Ethnicity



Race



Language and Marginalization

Fishing regulations can be complex. Documents are rarely translated from English into other languages. Lack of strong English language skills could affect participants' ability to engage effectively in the fisheries management process. While these numbers correspond to the overall community in Scituate they may indicate a population needing assistance in integrating their needs and concerns into the process.

Foreign Born: **4.6%**
National Average: **12.7%**

Speak English less than very well: **2%**
National Average: **8.7%**

Social Indicators

Social indicators are quantitative measures that describe the well-being of communities and are used to describe social phenomena over time. Below are a series of indices for Portsmouth that provide measures of fishing engagement and reliance, and social vulnerability. An index combines variables of interest and are used to evaluate community well-being in terms of social, economic and psychological welfare.

Fishing engagement and reliance indices portray the importance or level of dependence of commercial or recreational fishing to coastal communities. The indices include: Commercial Engagement, Commercial Reliance, Recreational Engagement and Recreational Reliance.

Social vulnerability indices represent social factors that can shape either an individual or community's ability to adapt to change. These factors exist within all communities regardless of the importance of fishing. The indices include: Poverty, Population Composition, and Personal Disruption.

Gentrification Pressure indices characterize those factors that, over time may indicate a threat to the viability of a commercial or recreational working waterfront, including infrastructure. The indices include: Retire Migration, Urban Sprawl, Natural Amenities and Housing Disruption.

The factor scores for each index are normalized so that zero is the mean. Therefore, a higher value implies more engagement or reliance upon fishing or higher social vulnerability or vulnerability to gentrification. Learn more about the [social indicators for fishing communities](#).

Social Indicators



Plymouth, MA

Where is Plymouth located?

Scituate is a town with a population of 56,468 and classified by the census as falling within an urbanized area. Rural to urban is really a continuum. Increasing urbanization indicates that a community has more jobs overall, more kinds of jobs, and more services like hospitals, social workers and job training centers. However, increasing urbanization can also mean greater pressure to transform working waterfronts for alternative uses, such as hotels or tourist shops.




Involvement in Fisheries

What species are landed in Plymouth?

The landings associated with a fishing community tell us what species are important to that community. The diversity of species caught also is indicative of a community's ability to adapt to changing environmental conditions (e.g. populations of specific fish stocks) or changes in fishing regulations that restrict access to resources.

*Groundfish includes cod, winter fl., witch fl., yellowtail fl., am. plaice, haddock, white hake, redfish, pollock.

**Whiting includes red hake, ocean pout, black whiting, whiting.



What are the characteristics of the fishing vessels in Plymouth?

The number of fishing vessels in a given port provides a sense of the scale of fishing in that port. Where a large port may serve as the homeport for hundreds of vessels, a smaller one may only have a handful. The number of vessels also may provide a rough sense of the number of fishing-related jobs (e.g. crew positions, jobs in shoreside industries) available in a given location.

Size also matters. Larger vessels can travel farther offshore and stay out for longer periods more easily than smaller vessels. These differences also affect family life. Smaller dayboat fishermen tend to return home every day whereas fishermen on larger vessels may be away from home for weeks on long and distant fishing expeditions.


Apart from the lobster fleet, smaller boats also tend to catch a broader range of species where their larger counterparts are more specialized (e.g. limited access scallop boats and herring pair trawlers). All these characteristics help illuminate the potential impacts of regulatory changes on a given community.

Demographic Attributes


Educational Attainment

The level of educational attainment in a community is associated with issues important for community development, such as income and poverty levels, unemployment rates, and local participation in community activities.

Number of Vessels by Size (2014)



Educational Attainment



How do people make a living in Plymouth?


Just as the range of fish species harvested by town residents speaks to their ability to adapt to environmental change, the diversity in local occupations indicates the ability of a community to adapt to economic changes, including changes in the local fishing economy. Is there one predominant industry, for instance, or is there a range of economic opportunities? How many occupations are available that offer incomes similar to fishing or require skills and education common to the average fisherman? How many jobs are available that would provide a working environment that fishermen would be comfortable with?

Unemployment Rate: **5.8%**

National Rate: **7.9%***

The unemployment rate in a community is one indicator of the level of opportunity that may exist for fishermen who lose their jobs to find alternative ways of making a living. The unemployment rate may also indicate the desirability of fishing in the face of other opportunities.

*Source: U.S. Department of Labor, Bureau of Labor Statistics



Median Household Income: **\$74,767.00**

National Average: **\$51,914.00** (2011)

Individuals in Plymouth living in poverty: **6.5%**

The poverty threshold for an individual is defined by the US Census for 2010 as \$11,139. The percentage of a town's population living under this economic threshold is an indicator of the residents' ability to adjust to loss of income and job opportunities in fishing-related and other local industries.


Age structure of residents

Age structure provides potential indications of many broader community issues and institutions. A large number of older residents may be associated with a retirement community or an out-migration of young people. For many fishing communities, an aging population can indicate gentrification, a process that may affect fishermen's access to the waterfront. In some remote coastal communities, people in their late teens or early twenties may leave to look for work or pursue an education outside of their community. A very large population of young people, on the other hand, may indicate the presence of universities or a military base.

Median age: **45.1**

National median: **37.2**


Population pyramid for Plymouth, year 2010
Source: www.census.gov




Ethnicity and Race

These factors give a sense of the cultural context of the community, and the relationship of fishing families and groups to the community in which they live. Is this community racially and ethnically diverse? In the northeast region, ethnic diversity in coastal communities tends to be higher in the Mid Atlantic than in New England, though there are significant exceptions in some fishing ports. Moreover, certain ethnic groups have long been associated with fishing in various specific ports throughout the region.

Ethnicity



Race



Language and Marginalization

Fishing regulations can be complex. Documents are rarely translated from English into other languages. Lack of strong English language skills could affect participants' ability to engage effectively in the fisheries management process. While these numbers correspond to the overall community in Scituate they may indicate a population needing assistance in integrating their needs and concerns into the process.

Foreign Born: **4.8%**
National Average: **12.7%**

Speak English less than very well: **2.6%**
National Average: **8.7%**

Social Indicators

Social indicators are quantitative measures that describe the well-being of communities and are used to describe social phenomena over time. Below are a series of indices for Portsmouth that provide measures of fishing engagement and reliance, and social vulnerability. An index combines variables of interest and are used to evaluate community well-being in terms of social, economic and psychological welfare.

Fishing engagement and reliance indices portray the importance or level of dependence of commercial or recreational fishing to coastal communities. The indices include: Commercial Engagement, Commercial Reliance, Recreational Engagement and Recreational Reliance.

Social vulnerability indices represent social factors that can shape either an individual or community's ability to adapt to change. These factors exist within all communities regardless of the importance of fishing. The indices include: Poverty, Population Composition, and Personal Disruption.

Gentrification Pressure indices characterize those factors that, over time may indicate a threat to the viability of a commercial or recreational working waterfront, including infrastructure. The indices include: Retire Migration, Urban Sprawl, Natural Amenities and Housing Disruption.

The factor scores for each index are normalized so that zero is the mean. Therefore, a higher value implies more engagement or reliance upon fishing or higher social vulnerability or vulnerability to gentrification. Learn more about the [social indicators for fishing communities](#).

Social Indicators



Northeast Fisheries Science Center
Social Sciences Branch

Chapter 4 Appendices

Appendix 4.1 Decision Support Tool Model Runs

Appendix 4.1.1 Baseline Information

The baseline information on right whale habitat density and trap/pot risk within Massachusetts portion of Lobster Management Area 1, including the reduction achieved by the 2021 Final Rule (86 FR 51970).

ModelConfiguration	
1	Model Start Time: 2023-01-10 11:01:13; End Time: 2023-01-10 11:04:42; Duration 00:03:29.4
2	Home Directory: //net/work4/LobsterGroup/Management/RightWhales/DecisionSupportTool
3	Model Version: DST_Beta_V4.1.0.R
4	InputActions: EmergencyRule2023/Ph1v4.0.3_TP_MA_LMA1.csv
5	MapRefDomain: MapRef_3.9.2.Rdata
6	GearMap: FisheryInputs_AllEastCoast_v4.0.1.Rdata
7	Threat Model: ThreatMod_RW_Selectivity_Uncertainty_WithZero.Rdata
8	Whale Habitat Model: Duke_RightWhaleModel_v12_1019.Rdata
9	Whale Vertical Dist File: RightWhaleVerticalDistribution_CSA_V1.1.Rdata
10	Whale Vertical Dist Model: Ensemble
11	Whale Dimensions: WhaleDimensions_AdHoc.Rdata
12	
13	Comment:
14	CoOccurrence: FALSE
15	Run Test Scenario: TRUE
16	Include Ground Gear: TRUE
17	AggregateStrings: FALSE
18	HighResolution: TRUE
19	RelocationCostExp: 1
20	ExpressRedistribution: TRUE
21	RopeStrengthResolution: 500
22	MinGearDensity: 1e-05
23	
24	PrintTables= TRUE
25	PrintDefaultMaps= FALSE, FALSE
26	PrintScenarioMaps= TRUE, FALSE
27	PrintRedistributionMaps= TRUE
28	PrintMapsInHighResolution= FALSE
29	TruncateMaps= 0.995
30	WriteMapSources= TRUE
31	WriteOutputCsv= TRUE
32	WriteDetailedOutput= FALSE
33	PrintSummary= TRUE
34	ArchiveInputSpreadsheet: FALSE

Total Whale Density

	Variable	Month	Default	Scenario	Reduction
1	WhaleDensity	1	27.81	27.81	0 %
2	WhaleDensity	2	34.02	34.02	0 %
3	WhaleDensity	3	64.22	64.22	0 %
4	WhaleDensity	4	90.21	90.21	0 %
5	WhaleDensity	5	21.41	21.41	0 %
6	WhaleDensity	6	0.36	0.36	0 %
7	WhaleDensity	7	0.47	0.47	0 %
8	WhaleDensity	8	0.12	0.12	0 %
9	WhaleDensity	9	0.06	0.06	0 %
10	WhaleDensity	10	0.21	0.21	0 %
11	WhaleDensity	11	1.88	1.88	0 %
12	WhaleDensity	12	7.17	7.17	0 %
13	WhaleDensity	Total	247.95	247.95	0 %

Final Relative Risk Scores – Mean Threat

	Variable	Month	Default	Scenario	Reduction
1	RelativeRisk_Threat	1	2,362.10	1,782.64	24.5 %
2	RelativeRisk_Threat	2	918.22	623.55	32.1 %
3	RelativeRisk_Threat	3	1,890.60	435.72	77 %
4	RelativeRisk_Threat	4	6,258.69	1,108.35	82.3 %
5	RelativeRisk_Threat	5	8,687.93	588.68	93.2 %
6	RelativeRisk_Threat	6	62.25	43.10	30.8 %
7	RelativeRisk_Threat	7	92.14	63.34	31.3 %
8	RelativeRisk_Threat	8	27.65	18.59	32.7 %
9	RelativeRisk_Threat	9	17.54	12.04	31.4 %
10	RelativeRisk_Threat	10	80.84	54.23	32.9 %
11	RelativeRisk_Threat	11	736.28	503.02	31.7 %
12	RelativeRisk_Threat	12	2,205.09	1,697.13	23 %
13	RelativeRisk_Threat	Total	23,339.34	6,930.39	70.3 %

The baseline information on right whale habitat density and trap/pot risk within the Northeast Trap/Pot Management Region, including the reduction achieved by the 2021 Final Rule (86 FR 51970).

ModelConfiguration	
1	Model Start Time: 2023-01-10 10:19:06; End Time: 2023-01-10 10:45:07; Duration 00:26:00.6
2	Home Directory: //net/work4/LobsterGroup/Management/RightWhales/DecisionSupportTool
3	Model Version: DST_Beta_V4.1.0.R
4	InputActions: EmergencyRule2023/Ph1v4.0.3_TP_NE.csv
5	MapRefDomain: MapRef_3.9.2.Rdata
6	GearMap: FisheryInputs_AllEastCoast_v4.0.1.Rdata
7	Threat Model: ThreatMod_RW_Selectivity_Uncertainty_WithZero.Rdata
8	Whale Habitat Model: Duke_RightWhaleModel_v12_1019.Rdata
9	Whale Vertical Dist File: RightWhaleVerticalDistribution_CSA_V1.1.Rdata
10	Whale Vertical Dist Model: Ensemble
11	Whale Dimensions: WhaleDimensions_AdHoc.Rdata
12	
13	Comment:
14	CoOccurrence: FALSE
15	Run Test Scenario: TRUE
16	Include Ground Gear: TRUE
17	AggregateStrings: FALSE
18	HighResolution: TRUE
19	RelocationCostExp: 1
20	ExpressRedistribution: TRUE
21	RopeStrengthResolution: 500
22	MinGearDensity: 1e-05
23	
24	PrintTables= TRUE
25	PrintDefaultMaps= FALSE, FALSE
26	PrintScenarioMaps= TRUE, FALSE
27	PrintRedistributionMaps= TRUE
28	PrintMapsInHighResolution= FALSE
29	TruncateMaps= 0.995
30	WriteMapSources= TRUE
31	WriteOutputCsv= TRUE
32	WriteDetailedOutput= FALSE
33	PrintSummary= TRUE
34	ArchiveInputSpreadsheet: FALSE

Total Whale Density

	Variable	Month	Default	Scenario	Reduction
1	WhaleDensity	1	269.94	269.94	0 %
2	WhaleDensity	2	256.91	256.91	0 %
3	WhaleDensity	3	275.05	275.05	0 %
4	WhaleDensity	4	331.03	331.03	0 %
5	WhaleDensity	5	340.40	340.40	0 %
6	WhaleDensity	6	159.98	159.98	0 %
7	WhaleDensity	7	115.26	115.26	0 %
8	WhaleDensity	8	13.41	13.41	0 %
9	WhaleDensity	9	15.42	15.42	0 %
10	WhaleDensity	10	43.73	43.73	0 %
11	WhaleDensity	11	115.15	115.15	0 %
12	WhaleDensity	12	152.86	152.86	0 %
13	WhaleDensity	Total	2,089.15	2,089.15	0 %

Final Relative Risk Scores – Mean Threat

	Variable	Month	Default	Scenario	Reduction
1	RelativeRisk_Threat	1	13,651.29	7,112.35	47.9 %
2	RelativeRisk_Threat	2	6,474.68	4,018.56	37.9 %
3	RelativeRisk_Threat	3	6,569.32	3,255.01	50.5 %
4	RelativeRisk_Threat	4	12,399.08	4,126.54	66.7 %
5	RelativeRisk_Threat	5	15,373.66	4,463.38	71 %
6	RelativeRisk_Threat	6	8,333.29	5,478.04	34.3 %
7	RelativeRisk_Threat	7	9,647.26	6,302.99	34.7 %
8	RelativeRisk_Threat	8	1,844.90	1,185.29	35.8 %
9	RelativeRisk_Threat	9	1,191.91	754.02	36.7 %
10	RelativeRisk_Threat	10	4,089.43	2,128.30	48 %
11	RelativeRisk_Threat	11	7,567.17	3,954.50	47.7 %
12	RelativeRisk_Threat	12	7,350.29	4,453.16	39.4 %
13	RelativeRisk_Threat	Total	94,492.30	47,232.14	50 %

The baseline information on right whale habitat density and trap/pot risk coastwide, including the reduction achieved by the 2021 Final Rule (86 FR 51970).

ModelConfiguration	
1	Model Start Time: 2023-01-10 09:44:06; End Time: 2023-01-10 10:16:50; Duration 00:32:43.2
2	Home Directory: //net/work4/LobsterGroup/Management/RightWhales/DecisionSupportTool
3	Model Version: DST_Beta_V4.1.0.R
4	InputActions: EmergencyRule2023/Ph1v4.0.3_TP.csv
5	MapRefDomain: MapRef_3.9.2.Rdata
6	GearMap: FisheryInputs_AllEastCoast_v4.0.1.Rdata
7	Threat Model: ThreatMod_RW_Selectivity_Uncertainty_WithZero.Rdata
8	Whale Habitat Model: Duke_RightWhaleModel_v12_1019.Rdata
9	Whale Vertical Dist File: RightWhaleVerticalDistribution_CSA_V1.1.Rdata
10	Whale Vertical Dist Model: Ensemble
11	Whale Dimensions: WhaleDimensions_AdHoc.Rdata
12	
13	Comment:
14	CoOccurrence: FALSE
15	Run Test Scenario: TRUE
16	Include Ground Gear: TRUE
17	AggregateStrings: FALSE
18	HighResolution: TRUE
19	RelocationCostExp: 1
20	ExpressRedistribution: TRUE
21	RopeStrengthResolution: 500
22	MinGearDensity: 1e-05
23	
24	PrintTables= TRUE
25	PrintDefaultMaps= FALSE, FALSE
26	PrintScenarioMaps= TRUE, FALSE
27	PrintRedistributionMaps= TRUE
28	PrintMapsInHighResolution= FALSE
29	TruncateMaps= 0.995
30	WriteMapSources= TRUE
31	WriteOutputCsv= TRUE
32	WriteDetailedOutput= FALSE
33	PrintSummary= TRUE
34	ArchiveInputSpreadsheet: FALSE

Total Whale Density

	Variable	Month	Default	Scenario	Reduction
1	WhaleDensity	1	355.39	355.39	0 %
2	WhaleDensity	2	366.43	366.43	0 %
3	WhaleDensity	3	375.20	375.20	0 %
4	WhaleDensity	4	407.70	407.70	0 %
5	WhaleDensity	5	364.81	364.81	0 %
6	WhaleDensity	6	166.28	166.28	0 %
7	WhaleDensity	7	117.82	117.82	0 %
8	WhaleDensity	8	15.73	15.73	0 %
9	WhaleDensity	9	19.10	19.10	0 %
10	WhaleDensity	10	54.94	54.94	0 %
11	WhaleDensity	11	132.80	132.80	0 %
12	WhaleDensity	12	201.97	201.97	0 %
13	WhaleDensity	Total	2,578.18	2,578.18	0 %

Final Relative Risk Scores – Mean Threat

	Variable	Month	Default	Scenario	Reduction
1	RelativeRisk_Threat	1	14,095.75	7,559.27	46.4 %
2	RelativeRisk_Threat	2	6,767.33	4,292.60	36.6 %
3	RelativeRisk_Threat	3	6,759.06	3,421.45	49.4 %
4	RelativeRisk_Threat	4	12,722.55	4,416.41	65.3 %
5	RelativeRisk_Threat	5	15,487.03	4,577.99	70.4 %
6	RelativeRisk_Threat	6	8,382.55	5,527.84	34.1 %
7	RelativeRisk_Threat	7	9,668.92	6,324.93	34.6 %
8	RelativeRisk_Threat	8	1,862.60	1,203.18	35.4 %
9	RelativeRisk_Threat	9	1,215.45	777.75	36 %
10	RelativeRisk_Threat	10	4,156.58	2,195.72	47.2 %
11	RelativeRisk_Threat	11	7,782.03	4,170.17	46.4 %
12	RelativeRisk_Threat	12	7,908.71	5,013.28	36.6 %
13	RelativeRisk_Threat	Total	96,808.56	49,480.58	48.9 %

The baseline information on right whale habitat density and trap/pot risk within MRA Wedge, Alternative 2 proposed emergency closure area.

ModelConfiguration	
1	Model Start Time: 2023-01-06 17:40:27; End Time: 2023-01-06 17:40:49; Duration 00:00:22.38
2	Home Directory: //net/work4/LobsterGroup/Management/RightWhales/DecisionSupportTool
3	Model Version: DST_Beta_V4.1.0.R
4	InputActions: CrystalsSubfolder/EmergencyRule2023/PrefConst_Baseline.csv
5	MapRefDomain: MapRef_3.9.2.Rdata
6	GearMap: FisheryInputs_AllEastCoast_v4.0.1.Rdata
7	Threat Model: ThreatMod_RW_Selectivity_Uncertainty_WithZero.Rdata
8	Whale Habitat Model: Duke_RightWhaleModel_v12_1019.Rdata
9	Whale Vertical Dist File: RightWhaleVerticalDistribution_CSA_V1.1.Rdata
10	Whale Vertical Dist Model: Ensemble
11	Whale Dimensions: WhaleDimensions_AdHoc.Rdata
12	
13	Comment:
14	CoOccurrence: FALSE
15	Run Test Scenario: TRUE
16	Include Ground Gear: TRUE
17	AggregateStrings: FALSE
18	HighResolution: TRUE
19	RelocationCostExp: 1
20	ExpressRedistribution: TRUE
21	RopeStrengthResolution: 500
22	MinGearDensity: 1e-05
23	
24	PrintTables= TRUE
25	PrintDefaultMaps= TRUE, TRUE
26	PrintScenarioMaps= TRUE, TRUE
27	PrintRedistributionMaps= TRUE
28	PrintMapsInHighResolution= FALSE
29	TruncateMaps= 0.995
30	WriteMapSources= TRUE
31	WriteOutputCsv= TRUE
32	WriteDetailedOutput= FALSE
33	PrintSummary= TRUE
34	ArchiveInputSpreadsheet: FALSE

Total Whale Density

	Variable	Month	Default	Scenario	Reduction
1	WhaleDensity	1	0.04	0.04	0 %
2	WhaleDensity	2	0.04	0.04	0 %
3	WhaleDensity	3	1.43	1.43	0 %
4	WhaleDensity	4	3.30	3.30	0 %
5	WhaleDensity	5	0.13	0.13	0 %
6	WhaleDensity	6	0.00	0.00	0 %
7	WhaleDensity	7	0.00	0.00	0 %
8	WhaleDensity	8	0.00	0.00	0 %
9	WhaleDensity	9	0.00	0.00	0 %
10	WhaleDensity	10	0.00	0.00	0 %
11	WhaleDensity	11	0.00	0.00	0 %
12	WhaleDensity	12	0.00	0.00	0 %
13	WhaleDensity	Total	4.94	4.94	0 %

Final Relative Risk Scores – Mean Threat

	Variable	Month	Default	Scenario	Reduction
1	RelativeRisk_Threat	1	8.23	8.23	0 %
2	RelativeRisk_Threat	2	9.32	9.32	0 %
3	RelativeRisk_Threat	3	346.10	346.10	0 %
4	RelativeRisk_Threat	4	1,029.94	1,029.94	0 %
5	RelativeRisk_Threat	5	25.91	25.91	0 %
6	RelativeRisk_Threat	6	0.32	0.32	0 %
7	RelativeRisk_Threat	7	0.39	0.39	0 %
8	RelativeRisk_Threat	8	0.20	0.20	0 %
9	RelativeRisk_Threat	9	0.17	0.17	0 %
10	RelativeRisk_Threat	10	0.00	0.00	-0.2 %
11	RelativeRisk_Threat	11	0.18	0.18	0 %
12	RelativeRisk_Threat	12	1.29	1.29	0 %
13	RelativeRisk_Threat	Total	1,422.05	1,422.05	0 %

The baseline information on humpback whale habitat density and trap/pot risk within MRA Wedge, Alternative 2 proposed emergency closure area.

ModelConfiguration	
1	Model Start Time: 2023-01-12 12:43:48; End Time: 2023-01-12 12:44:23; Duration 00:00:34.71
2	Home Directory: //net/work4/LobsterGroup/Management/RightWhales/DecisionSupportTool
3	Model Version: DST_Beta_V4.1.0.R
4	InputActions: EmergencyRule2023/PrefConst_Baseline_hback.csv
5	MapRefDomain: MapRef_3.9.2.Rdata
6	GearMap: FisheryInputs_AllEastCoast_v4.0.1.Rdata
7	Threat Model: ThreatMod_RW_Selectivity_Uncertainty_WithZero.Rdata
8	Whale Habitat Model: Duke_HumpbackWhaleModel_v11_0919.Rdata
9	Whale Vertical Dist File: RightWhaleVerticalDistribution_CSA_V1.1.Rdata
10	Whale Vertical Dist Model: Ensemble
11	Whale Dimensions: WhaleDimensions_AdHoc.Rdata
12	
13	Comment:
14	CoOccurrence: FALSE
15	Run Test Scenario: TRUE
16	Include Ground Gear: TRUE
17	AggregateStrings: FALSE
18	HighResolution: TRUE
19	RelocationCostExp: 1
20	ExpressRedistribution: TRUE
21	RopeStrengthResolution: 500
22	MinGearDensity: 1e-05
23	
24	PrintTables= TRUE
25	PrintDefaultMaps= FALSE, FALSE
26	PrintScenarioMaps= TRUE, FALSE
27	PrintRedistributionMaps= TRUE
28	PrintMapsInHighResolution= FALSE
29	TruncateMaps= 0.995
30	WriteMapSources= TRUE
31	WriteOutputCsv= TRUE
32	WriteDetailedOutput= FALSE
33	PrintSummary= TRUE
34	ArchiveInputSpreadsheet: FALSE

Total Whale Density

	Variable	Month	Default	Scenario	Reduction
1	WhaleDensity	1	0.21	0.21	0 %
2	WhaleDensity	2	0.15	0.15	0 %
3	WhaleDensity	3	0.20	0.20	0 %
4	WhaleDensity	4	0.89	0.89	0 %
5	WhaleDensity	5	0.97	0.97	0 %
6	WhaleDensity	6	0.86	0.86	0 %
7	WhaleDensity	7	1.08	1.08	0 %
8	WhaleDensity	8	0.86	0.86	0 %
9	WhaleDensity	9	0.93	0.93	0 %
10	WhaleDensity	10	1.07	1.07	0 %
11	WhaleDensity	11	1.32	1.32	0 %
12	WhaleDensity	12	0.48	0.48	0 %
13	WhaleDensity	Total	9.01	9.01	0 %

Final Relative Risk Scores – Mean Threat

	Variable	Month	Default	Scenario	Reduction
1	RelativeRisk_Threat	1	44.36	44.36	0 %
2	RelativeRisk_Threat	2	36.44	36.44	0 %
3	RelativeRisk_Threat	3	49.22	49.22	0 %
4	RelativeRisk_Threat	4	277.50	277.50	0 %
5	RelativeRisk_Threat	5	194.94	194.94	0 %
6	RelativeRisk_Threat	6	150.13	150.13	0 %
7	RelativeRisk_Threat	7	184.79	184.79	0 %
8	RelativeRisk_Threat	8	194.27	194.27	0 %
9	RelativeRisk_Threat	9	274.15	274.15	0 %
10	RelativeRisk_Threat	10	415.96	415.96	0 %
11	RelativeRisk_Threat	11	544.54	544.54	0 %
12	RelativeRisk_Threat	12	161.08	161.08	0 %
13	RelativeRisk_Threat	Total	2,527.39	2,527.39	0 %

The baseline information on fin whale habitat density and trap/pot co-occurrence (i.e. overlap between whales and gear within MRA Wedge, Alternative 2 proposed emergency closure area.

ModelConfiguration	
1	Model Start Time: 2023-01-12 12:49:52; End Time: 2023-01-12 12:50:22; Duration 00:00:30.25
2	Home Directory: //net/work4/LobsterGroup/Management/RightWhales/DecisionSupportTool
3	Model Version: DST_Beta_V4.1.0.R
4	InputActions: EmergencyRule2023/PrefConst_Baseline_fin.csv
5	MapRefDomain: MapRef_3.9.2.Rdata
6	GearMap: FisheryInputs_AllEastCoast_v4.0.1.Rdata
7	Threat Model: ThreatMod_RW_Selectivity_Uncertainty_WithZero.Rdata
8	Whale Habitat Model: Duke_FinWhaleModel_v12.Rdata
9	Whale Vertical Dist File: RightWhaleVerticalDistribution_CSA_V1.1.Rdata
10	Whale Vertical Dist Model: Ensemble
11	Whale Dimensions: WhaleDimensions_AdHoc.Rdata
12	
13	Comment:
14	CoOccurrence: FALSE
15	Run Test Scenario: TRUE
16	Include Ground Gear: TRUE
17	AggregateStrings: FALSE
18	HighResolution: TRUE
19	RelocationCostExp: 1
20	ExpressRedistribution: TRUE
21	RopeStrengthResolution: 500
22	MinGearDensity: 1e-05
23	
24	PrintTables= TRUE
25	PrintDefaultMaps= FALSE, FALSE
26	PrintScenarioMaps= TRUE, FALSE
27	PrintRedistributionMaps= TRUE
28	PrintMapsInHighResolution= FALSE
29	TruncateMaps= 0.995
30	WriteMapSources= TRUE
31	WriteOutputCsv= TRUE
32	WriteDetailedOutput= FALSE
33	PrintSummary= TRUE
34	ArchiveInputSpreadsheet: FALSE

Total Whale Density

	Variable	Month	Default	Scenario	Reduction
1	WhaleDensity	1	0.81	0.81	0 %
2	WhaleDensity	2	0.38	0.38	0 %
3	WhaleDensity	3	0.34	0.34	0 %
4	WhaleDensity	4	0.76	0.76	0 %
5	WhaleDensity	5	1.11	1.11	0 %
6	WhaleDensity	6	1.13	1.13	0 %
7	WhaleDensity	7	0.96	0.96	0 %
8	WhaleDensity	8	1.46	1.46	0 %
9	WhaleDensity	9	0.55	0.55	0 %
10	WhaleDensity	10	0.29	0.29	0 %
11	WhaleDensity	11	0.48	0.48	0 %
12	WhaleDensity	12	0.79	0.79	0 %
13	WhaleDensity	Total	9.06	9.06	0 %

Final Relative Risk Scores – CoOccurrence

	Variable	Month	Default	Scenario	Reduction
1	RelativeRisk_CoOccurrence	1	1,220.68	1,220.68	0 %
2	RelativeRisk_CoOccurrence	2	676.10	676.10	0 %
3	RelativeRisk_CoOccurrence	3	599.07	599.07	0 %
4	RelativeRisk_CoOccurrence	4	1,751.18	1,751.18	0 %
5	RelativeRisk_CoOccurrence	5	1,647.36	1,647.36	0 %
6	RelativeRisk_CoOccurrence	6	1,460.96	1,460.96	0 %
7	RelativeRisk_CoOccurrence	7	1,237.72	1,237.72	0 %
8	RelativeRisk_CoOccurrence	8	2,472.55	2,472.55	0 %
9	RelativeRisk_CoOccurrence	9	1,210.72	1,210.72	0 %
10	RelativeRisk_CoOccurrence	10	825.89	825.89	0 %
11	RelativeRisk_CoOccurrence	11	1,460.47	1,460.47	0 %
12	RelativeRisk_CoOccurrence	12	1,946.38	1,946.38	0 %
13	RelativeRisk_CoOccurrence	Total	16,509.06	16,509.06	0 %

The baseline information on right whale habitat density and trap/pot risk within MRA Wedge North, Alternative 3 proposed emergency closure area.

ModelConfiguration	
1	Model Start Time: 2023-01-06 17:36:36; End Time: 2023-01-06 17:37:24; Duration 00:00:48.28
2	Home Directory: //net/work4/LobsterGroup/Management/RightWhales/DecisionSupportTool
3	Model Version: DST_Beta_V4.1.0.R
4	InputActions: CrystalsSubfolder/EmergencyRule2023/NonPrefConst_Baseline.csv
5	MapRefDomain: MapRef_3.9.2.Rdata
6	GearMap: FisheryInputs_AllEastCoast_v4.0.1.Rdata
7	Threat Model: ThreatMod_RW_Selectivity_Uncertainty_WithZero.Rdata
8	Whale Habitat Model: Duke_RightWhaleModel_v12_1019.Rdata
9	Whale Vertical Dist File: RightWhaleVerticalDistribution_CSA_V1.1.Rdata
10	Whale Vertical Dist Model: Ensemble
11	Whale Dimensions: WhaleDimensions_AdHoc.Rdata
12	
13	Comment:
14	CoOccurrence: FALSE
15	Run Test Scenario: TRUE
16	Include Ground Gear: TRUE
17	AggregateStrings: FALSE
18	HighResolution: TRUE
19	RelocationCostExp: 1
20	ExpressRedistribution: TRUE
21	RopeStrengthResolution: 500
22	MinGearDensity: 1e-05
23	
24	PrintTables= TRUE
25	PrintDefaultMaps= TRUE, TRUE
26	PrintScenarioMaps= TRUE, TRUE
27	PrintRedistributionMaps= TRUE
28	PrintMapsInHighResolution= FALSE
29	TruncateMaps= 0.995
30	WriteMapSources= TRUE
31	WriteOutputCsv= TRUE
32	WriteDetailedOutput= FALSE
33	PrintSummary= TRUE
34	ArchiveInputSpreadsheet: FALSE

Total Whale Density

	Variable	Month	Default	Scenario	Reduction
1	WhaleDensity	1	6.16	6.16	0 %
2	WhaleDensity	2	6.66	6.66	0 %
3	WhaleDensity	3	2.42	2.42	0 %
4	WhaleDensity	4	4.62	4.62	0 %
5	WhaleDensity	5	3.05	3.05	0 %
6	WhaleDensity	6	0.37	0.37	0 %
7	WhaleDensity	7	0.60	0.60	0 %
8	WhaleDensity	8	0.07	0.07	0 %
9	WhaleDensity	9	0.03	0.03	0 %
10	WhaleDensity	10	1.00	1.00	0 %
11	WhaleDensity	11	4.38	4.38	0 %
12	WhaleDensity	12	1.90	1.90	0 %
13	WhaleDensity	Total	31.27	31.27	0 %

Final Relative Risk Scores – Mean Threat

	Variable	Month	Default	Scenario	Reduction
1	RelativeRisk_Threat	1	1,313.89	1,313.89	0 %
2	RelativeRisk_Threat	2	1,424.45	1,424.45	0 %
3	RelativeRisk_Threat	3	553.68	553.68	0 %
4	RelativeRisk_Threat	4	1,386.82	1,386.82	0 %
5	RelativeRisk_Threat	5	364.88	364.88	0 %
6	RelativeRisk_Threat	6	31.64	31.64	0 %
7	RelativeRisk_Threat	7	41.84	41.84	0 %
8	RelativeRisk_Threat	8	12.44	12.44	0 %
9	RelativeRisk_Threat	9	7.00	7.00	0 %
10	RelativeRisk_Threat	10	198.41	198.41	0 %
11	RelativeRisk_Threat	11	953.48	953.48	0 %
12	RelativeRisk_Threat	12	449.45	449.45	0 %
13	RelativeRisk_Threat	Total	6,737.99	6,737.99	0 %

The baseline information on humpback whale habitat density and trap/pot risk within MRA Wedge North, Alternative 3 proposed emergency closure area.

ModelConfiguration	
1	Model Start Time: 2023-01-12 12:46:17; End Time: 2023-01-12 12:47:09; Duration 00:00:52.54
2	Home Directory: //net/work4/LobsterGroup/Management/RightWhales/DecisionSupportTool
3	Model Version: DST_Beta_V4.1.0.R
4	InputActions: EmergencyRule2023/NonPrefConst_Baseline_hback.csv
5	MapRefDomain: MapRef_3.9.2.Rdata
6	GearMap: FisheryInputs_AllEastCoast_v4.0.1.Rdata
7	Threat Model: ThreatMod_RW_Selectivity_Uncertainty_WithZero.Rdata
8	Whale Habitat Model: Duke_HumpbackWhaleModel_v11_0919.Rdata
9	Whale Vertical Dist File: RightWhaleVerticalDistribution_CSA_V1.1.Rdata
10	Whale Vertical Dist Model: Ensemble
11	Whale Dimensions: WhaleDimensions_AdHoc.Rdata
12	
13	Comment:
14	CoOccurrence: FALSE
15	Run Test Scenario: TRUE
16	Include Ground Gear: TRUE
17	AggregateStrings: FALSE
18	HighResolution: TRUE
19	RelocationCostExp: 1
20	ExpressRedistribution: TRUE
21	RopeStrengthResolution: 500
22	MinGearDensity: 1e-05
23	
24	PrintTables= TRUE
25	PrintDefaultMaps= FALSE, FALSE
26	PrintScenarioMaps= TRUE, FALSE
27	PrintRedistributionMaps= TRUE
28	PrintMapsInHighResolution= FALSE
29	TruncateMaps= 0.995
30	WriteMapSources= TRUE
31	WriteOutputCsv= TRUE
32	WriteDetailedOutput= FALSE
33	PrintSummary= TRUE
34	ArchiveInputSpreadsheet: FALSE

Total Whale Density

	Variable	Month	Default	Scenario	Reduction
1	WhaleDensity	1	3.25	3.25	0 %
2	WhaleDensity	2	2.01	2.01	0 %
3	WhaleDensity	3	2.39	2.39	0 %
4	WhaleDensity	4	17.60	17.60	0 %
5	WhaleDensity	5	32.44	32.44	0 %
6	WhaleDensity	6	29.88	29.88	0 %
7	WhaleDensity	7	32.89	32.89	0 %
8	WhaleDensity	8	27.67	27.67	0 %
9	WhaleDensity	9	31.25	31.25	0 %
10	WhaleDensity	10	32.93	32.93	0 %
11	WhaleDensity	11	18.99	18.99	0 %
12	WhaleDensity	12	8.12	8.12	0 %
13	WhaleDensity	Total	239.39	239.39	0 %

Final Relative Risk Scores – Mean Threat

	Variable	Month	Default	Scenario	Reduction
1	RelativeRisk_Threat	1	623.80	623.80	0 %
2	RelativeRisk_Threat	2	408.19	408.19	0 %
3	RelativeRisk_Threat	3	499.93	499.93	0 %
4	RelativeRisk_Threat	4	5,269.32	5,269.32	0 %
5	RelativeRisk_Threat	5	6,670.04	6,670.04	0 %
6	RelativeRisk_Threat	6	6,792.26	6,792.26	0 %
7	RelativeRisk_Threat	7	6,725.08	6,725.08	0 %
8	RelativeRisk_Threat	8	6,619.17	6,619.17	0 %
9	RelativeRisk_Threat	9	9,019.26	9,019.26	0 %
10	RelativeRisk_Threat	10	11,788.48	11,788.48	0 %
11	RelativeRisk_Threat	11	6,730.38	6,730.38	0 %
12	RelativeRisk_Threat	12	2,107.65	2,107.65	0 %
13	RelativeRisk_Threat	Total	63,253.56	63,253.56	0 %

The baseline information on fin whale habitat density and trap/pot co-occurrence (i.e. overlap between whales and gear within MRA Wedge North, Alternative 3 proposed emergency closure area.

ModelConfiguration	
1	Model Start Time: 2023-01-12 12:47:42; End Time: 2023-01-12 12:48:31; Duration 00:00:49.18
2	Home Directory: //net/work4/LobsterGroup/Management/RightWhales/DecisionSupportTool
3	Model Version: DST_Beta_V4.1.0.R
4	InputActions: EmergencyRule2023/NonPrefConst_Baseline_fin.csv
5	MapRefDomain: MapRef_3.9.2.Rdata
6	GearMap: FisheryInputs_AllEastCoast_v4.0.1.Rdata
7	Threat Model: ThreatMod_RW_Selectivity_Uncertainty_WithZero.Rdata
8	Whale Habitat Model: Duke_FinWhaleModel_v12.Rdata
9	Whale Vertical Dist File: RightWhaleVerticalDistribution_CSA_V1.1.Rdata
10	Whale Vertical Dist Model: Ensemble
11	Whale Dimensions: WhaleDimensions_AdHoc.Rdata
12	
13	Comment:
14	CoOccurrence: FALSE
15	Run Test Scenario: TRUE
16	Include Ground Gear: TRUE
17	AggregateStrings: FALSE
18	HighResolution: TRUE
19	RelocationCostExp: 1
20	ExpressRedistribution: TRUE
21	RopeStrengthResolution: 500
22	MinGearDensity: 1e-05
23	
24	PrintTables= TRUE
25	PrintDefaultMaps= FALSE, FALSE
26	PrintScenarioMaps= TRUE, FALSE
27	PrintRedistributionMaps= TRUE
28	PrintMapsInHighResolution= FALSE
29	TruncateMaps= 0.995
30	WriteMapSources= TRUE
31	WriteOutputCsv= TRUE
32	WriteDetailedOutput= FALSE
33	PrintSummary= TRUE
34	ArchiveInputSpreadsheet: FALSE

Total Whale Density

	Variable	Month	Default	Scenario	Reduction
1	WhaleDensity	1	12.12	12.12	0 %
2	WhaleDensity	2	8.33	8.33	0 %
3	WhaleDensity	3	7.06	7.06	0 %
4	WhaleDensity	4	7.64	7.64	0 %
5	WhaleDensity	5	15.05	15.05	0 %
6	WhaleDensity	6	17.85	17.85	0 %
7	WhaleDensity	7	25.19	25.19	0 %
8	WhaleDensity	8	28.01	28.01	0 %
9	WhaleDensity	9	17.31	17.31	0 %
10	WhaleDensity	10	13.67	13.67	0 %
11	WhaleDensity	11	10.63	10.63	0 %
12	WhaleDensity	12	14.05	14.05	0 %
13	WhaleDensity	Total	176.91	176.91	0 %

Final Relative Risk Scores – CoOccurrence

	Variable	Month	Default	Scenario	Reduction
1	RelativeRisk_CoOccurrence	1	16,219.49	16,219.49	0 %
2	RelativeRisk_CoOccurrence	2	12,356.23	12,356.23	0 %
3	RelativeRisk_CoOccurrence	3	10,563.65	10,563.65	0 %
4	RelativeRisk_CoOccurrence	4	15,885.27	15,885.27	0 %
5	RelativeRisk_CoOccurrence	5	19,163.61	19,163.61	0 %
6	RelativeRisk_CoOccurrence	6	24,095.25	24,095.25	0 %
7	RelativeRisk_CoOccurrence	7	29,118.48	29,118.48	0 %
8	RelativeRisk_CoOccurrence	8	40,916.99	40,916.99	0 %
9	RelativeRisk_CoOccurrence	9	31,003.52	31,003.52	0 %
10	RelativeRisk_CoOccurrence	10	31,246.91	31,246.91	0 %
11	RelativeRisk_CoOccurrence	11	23,440.77	23,440.77	0 %
12	RelativeRisk_CoOccurrence	12	25,584.09	25,584.09	0 %
13	RelativeRisk_CoOccurrence	Total	279,594.26	279,594.26	0 %

Appendix 4.1.2 Alternative 2 Model Runs

Risk reduction presented in this section were run with a constraint for coastwide trap/pot fisheries. For section 6.2 of this EA, these results were then calculated relative to different baselines to reflect only risk reduction in addition to the previous rule based on the data presented in section 4.1.1 of this Appendix.

Gear Reduction (i.e. all lines removed)

February gear reduction in the MRA Wedge, Alternative 2 proposed emergency restricted area

	ModelConfiguration
1	Model Start Time: 2023-01-09 14:46:54; End Time: 2023-01-09 15:19:42; Duration 00:32:47.4
2	Home Directory: //network4/LobsterGroup/Management/RightWhales/DecisionSupportTool
3	Model Version: DST_Beta_V4.1.0.R
4	InputActions: CrystalsSubfolder/EmergencyRule2023/ER_Pref_GR_Feb_TP_wPh1v4.0.3.csv
5	MapRefDomain: MapRef_3.9.2.Rdata
6	GearMap: FisheryInputs_AllEastCoast_v4.0.1.Rdata
7	Threat Model: ThreatMod_RW_Selectivity_Uncertainty_WithZero.Rdata
8	Whale Habitat Model: Duke_RightWhaleModel_v12_1019.Rdata
9	Whale Vertical Dist File: RightWhaleVerticalDistribution_CSA_V1.1.Rdata
10	Whale Vertical Dist Model: Ensemble
11	Whale Dimensions: WhaleDimensions_AdHoc.Rdata
12	
13	Comment:
14	CoOccurrence: FALSE
15	Run Test Scenario: TRUE
16	Include Ground Gear: TRUE
17	AggregateStrings: FALSE
18	HighResolution: TRUE
19	RelocationCostExp: 1
20	ExpressRedistribution: TRUE
21	RopeStrengthResolution: 500
22	MinGearDensity: 1e-05
23	
24	PrintTables= TRUE
25	PrintDefaultMaps= TRUE, TRUE
26	PrintScenarioMaps= TRUE, TRUE
27	PrintRedistributionMaps= TRUE
28	PrintMapsInHighResolution= FALSE
29	TruncateMaps= 0.995
30	WriteMapSources= TRUE
31	WriteOutputCsv= TRUE
32	WriteDetailedOutput= FALSE
33	PrintSummary= TRUE
34	ArchiveInputSpreadsheet: FALSE

Final Relative Risk Scores – Mean Threat

	Variable	Month	Default	Scenario	Reduction
1	RelativeRisk_Threat	1	14,095.75	7,559.27	46.4 %
2	RelativeRisk_Threat	2	6,767.33	4,284.82	36.7 %
3	RelativeRisk_Threat	3	6,759.06	3,421.45	49.4 %
4	RelativeRisk_Threat	4	12,722.55	4,416.41	65.3 %
5	RelativeRisk_Threat	5	15,487.03	4,577.99	70.4 %
6	RelativeRisk_Threat	6	8,382.55	5,527.84	34.1 %
7	RelativeRisk_Threat	7	9,668.92	6,324.93	34.6 %
8	RelativeRisk_Threat	8	1,862.60	1,203.18	35.4 %
9	RelativeRisk_Threat	9	1,215.45	777.75	36 %
10	RelativeRisk_Threat	10	4,156.58	2,195.72	47.2 %
11	RelativeRisk_Threat	11	7,782.03	4,170.17	46.4 %
12	RelativeRisk_Threat	12	7,908.71	5,013.28	36.6 %
13	RelativeRisk_Threat	Total	96,808.56	49,472.80	48.9 %

March gear reduction in the MRA Wedge, Alternative 2 proposed emergency restricted area

ModelConfiguration	
1	Model Start Time: 2023-01-09 14:51:53; End Time: 2023-01-09 15:31:58; Duration 00:40:05.4
2	Home Directory: //net/work4/LobsterGroup/Management/RightWhales/DecisionSupportTool
3	Model Version: DST_Beta_V4.1.0.R
4	InputActions: CrystalsSubfolder/EmergencyRule2023/ER_Pref_GR_Mar_TP_wPh1v4.0.3.csv
5	MapRefDomain: MapRef_3.9.2.Rdata
6	GearMap: FisheryInputs_AllEastCoast_v4.0.1.Rdata
7	Threat Model: ThreatMod_RW_Selectivity_Uncertainty_WithZero.Rdata
8	Whale Habitat Model: Duke_RightWhaleModel_v12_1019.Rdata
9	Whale Vertical Dist File: RightWhaleVerticalDistribution_CSA_V1.1.Rdata
10	Whale Vertical Dist Model: Ensemble
11	Whale Dimensions: WhaleDimensions_AdHoc.Rdata
12	
13	Comment:
14	CoOccurrence: FALSE
15	Run Test Scenario: TRUE
16	Include Ground Gear: TRUE
17	AggregateStrings: FALSE
18	HighResolution: TRUE
19	RelocationCostExp: 1
20	ExpressRedistribution: TRUE
21	RopeStrengthResolution: 500
22	MinGearDensity: 1e-05
23	
24	PrintTables= TRUE
25	PrintDefaultMaps= TRUE, TRUE
26	PrintScenarioMaps= TRUE, TRUE
27	PrintRedistributionMaps= TRUE
28	PrintMapsInHighResolution= FALSE
29	TruncateMaps= 0.995
30	WriteMapSources= TRUE
31	WriteOutputCsv= TRUE
32	WriteDetailedOutput= FALSE
33	PrintSummary= TRUE
34	ArchiveInputSpreadsheet: FALSE

Final Relative Risk Scores – Mean Threat

	Variable	Month	Default	Scenario	Reduction
1	RelativeRisk_Threat	1	14,095.75	7,559.27	46.4 %
2	RelativeRisk_Threat	2	6,767.33	4,292.60	36.6 %
3	RelativeRisk_Threat	3	6,759.06	3,134.79	53.6 %
4	RelativeRisk_Threat	4	12,722.55	4,416.41	65.3 %
5	RelativeRisk_Threat	5	15,487.03	4,577.99	70.4 %
6	RelativeRisk_Threat	6	8,382.55	5,527.84	34.1 %
7	RelativeRisk_Threat	7	9,668.92	6,324.93	34.6 %
8	RelativeRisk_Threat	8	1,862.60	1,203.18	35.4 %
9	RelativeRisk_Threat	9	1,215.45	777.75	36 %
10	RelativeRisk_Threat	10	4,156.58	2,195.72	47.2 %
11	RelativeRisk_Threat	11	7,782.03	4,170.17	46.4 %
12	RelativeRisk_Threat	12	7,908.71	5,013.28	36.6 %
13	RelativeRisk_Threat	Total	96,808.56	49,193.92	49.2 %

April gear reduction in the MRA Wedge, Alternative 2 proposed emergency restricted area

ModelConfiguration	
1	Model Start Time: 2023-01-09 14:43:08; End Time: 2023-01-09 15:29:00; Duration 00:45:52.2
2	Home Directory: //net/work4/LobsterGroup/Management/RightWhales/DecisionSupportTool
3	Model Version: DST_Beta_V4.1.0.R
4	InputActions: CrystalsSubfolder/EmergencyRule2023/ER_Pref_GR_Apr_TP_wPh1v4.0.3.csv
5	MapRefDomain: MapRef_3.9.2.Rdata
6	GearMap: FisheryInputs_AllEastCoast_v4.0.1.Rdata
7	Threat Model: ThreatMod_RW_Selectivity_Uncertainty_WithZero.Rdata
8	Whale Habitat Model: Duke_RightWhaleModel_v12_1019.Rdata
9	Whale Vertical Dist File: RightWhaleVerticalDistribution_CSA_V1.1.Rdata
10	Whale Vertical Dist Model: Ensemble
11	Whale Dimensions: WhaleDimensions_AdHoc.Rdata
12	
13	Comment:
14	CoOccurrence: FALSE
15	Run Test Scenario: TRUE
16	Include Ground Gear: TRUE
17	AggregateStrings: FALSE
18	HighResolution: TRUE
19	RelocationCostExp: 1
20	ExpressRedistribution: TRUE
21	RopeStrengthResolution: 500
22	MinGearDensity: 1e-05
23	
24	PrintTables= TRUE
25	PrintDefaultMaps= TRUE, TRUE
26	PrintScenarioMaps= TRUE, TRUE
27	PrintRedistributionMaps= TRUE
28	PrintMapsInHighResolution= FALSE
29	TruncateMaps= 0.995
30	WriteMapSources= TRUE
31	WriteOutputCsv= TRUE
32	WriteDetailedOutput= FALSE
33	PrintSummary= TRUE
34	ArchiveInputSpreadsheet: FALSE

Final Relative Risk Scores – Mean Threat

	Variable	Month	Default	Scenario	Reduction
1	RelativeRisk_Threat	1	14,095.75	7,559.27	46.4 %
2	RelativeRisk_Threat	2	6,767.33	4,292.60	36.6 %
3	RelativeRisk_Threat	3	6,759.06	3,421.45	49.4 %
4	RelativeRisk_Threat	4	12,722.55	3,559.10	72 %
5	RelativeRisk_Threat	5	15,487.03	4,577.99	70.4 %
6	RelativeRisk_Threat	6	8,382.55	5,527.84	34.1 %
7	RelativeRisk_Threat	7	9,668.92	6,324.93	34.6 %
8	RelativeRisk_Threat	8	1,862.60	1,203.18	35.4 %
9	RelativeRisk_Threat	9	1,215.45	777.75	36 %
10	RelativeRisk_Threat	10	4,156.58	2,195.72	47.2 %
11	RelativeRisk_Threat	11	7,782.03	4,170.17	46.4 %
12	RelativeRisk_Threat	12	7,908.71	5,013.28	36.6 %
13	RelativeRisk_Threat	Total	96,808.56	48,623.27	49.8 %

Closure (i.e. all lines relocated)

February closure in the MRA Wedge, Alternative 2 proposed emergency restricted area

	ModelConfiguration
1	Model Start Time: 2023-01-09 14:37:35; End Time: 2023-01-09 15:05:40; Duration 00:28:05.4
2	Home Directory: //net/work4/LobsterGroup/Management/RightWhales/DecisionSupportTool
3	Model Version: DST_Beta_V4.1.0.R
4	InputActions: CrystalsSubfolder/EmergencyRule2023/ER_Pref_Clos_Feb_TP_wPh1v4.0.3.csv
5	MapRefDomain: MapRef_3.9.2.Rdata
6	GearMap: FisheryInputs_AllEastCoast_v4.0.1.Rdata
7	Threat Model: ThreatMod_RW_Selectivity_Uncertainty_WithZero.Rdata
8	Whale Habitat Model: Duke_RightWhaleModel_v12_1019.Rdata
9	Whale Vertical Dist File: RightWhaleVerticalDistribution_CSA_V1.1.Rdata
10	Whale Vertical Dist Model: Ensemble
11	Whale Dimensions: WhaleDimensions_AdHoc.Rdata
12	
13	Comment:
14	CoOccurrence: FALSE
15	Run Test Scenario: TRUE
16	Include Ground Gear: TRUE
17	AggregateStrings: FALSE
18	HighResolution: TRUE
19	RelocationCostExp: 1
20	ExpressRedistribution: TRUE
21	RopeStrengthResolution: 500
22	MinGearDensity: 1e-05
23	
24	PrintTables= TRUE
25	PrintDefaultMaps= TRUE, TRUE
26	PrintScenarioMaps= TRUE, TRUE
27	PrintRedistributionMaps= TRUE
28	PrintMapsInHighResolution= FALSE
29	TruncateMaps= 0.995
30	WriteMapSources= TRUE
31	WriteOutputCsv= TRUE
32	WriteDetailedOutput= FALSE
33	PrintSummary= TRUE
34	ArchiveInputSpreadsheet: FALSE

Final Relative Risk Scores – Mean Threat

	Variable	Month	Default	Scenario	Reduction
1	RelativeRisk_Threat	1	14,095.75	7,559.27	46.4 %
2	RelativeRisk_Threat	2	6,767.33	4,423.70	34.6 %
3	RelativeRisk_Threat	3	6,759.06	3,421.45	49.4 %
4	RelativeRisk_Threat	4	12,722.55	4,416.41	65.3 %
5	RelativeRisk_Threat	5	15,487.03	4,577.99	70.4 %
6	RelativeRisk_Threat	6	8,382.55	5,527.84	34.1 %
7	RelativeRisk_Threat	7	9,668.92	6,324.93	34.6 %
8	RelativeRisk_Threat	8	1,862.60	1,203.18	35.4 %
9	RelativeRisk_Threat	9	1,215.45	777.75	36 %
10	RelativeRisk_Threat	10	4,156.58	2,195.72	47.2 %
11	RelativeRisk_Threat	11	7,782.03	4,170.17	46.4 %
12	RelativeRisk_Threat	12	7,908.71	5,013.28	36.6 %
13	RelativeRisk_Threat	Total	96,808.56	49,611.68	48.8 %

March closure in the MRA Wedge, Alternative 2 proposed emergency restricted area

ModelConfiguration	
1	Model Start Time: 2023-01-09 14:40:39; End Time: 2023-01-09 15:12:00; Duration 00:31:20.4
2	Home Directory: //net/work4/LobsterGroup/Management/RightWhales/DecisionSupportTool
3	Model Version: DST_Beta_V4.1.0.R
4	InputActions: CrystalsSubfolder/EmergencyRule2023/ER_Pref_Clos_Mar_TP_wPh1v4.0.3.csv
5	MapRefDomain: MapRef_3.9.2.Rdata
6	GearMap: FisheryInputs_AllEastCoast_v4.0.1.Rdata
7	Threat Model: ThreatMod_RW_Selectivity_Uncertainty_WithZero.Rdata
8	Whale Habitat Model: Duke_RightWhaleModel_v12_1019.Rdata
9	Whale Vertical Dist File: RightWhaleVerticalDistribution_CSA_V1.1.Rdata
10	Whale Vertical Dist Model: Ensemble
11	Whale Dimensions: WhaleDimensions_AdHoc.Rdata
12	
13	Comment:
14	CoOccurrence: FALSE
15	Run Test Scenario: TRUE
16	Include Ground Gear: TRUE
17	AggregateStrings: FALSE
18	HighResolution: TRUE
19	RelocationCostExp: 1
20	ExpressRedistribution: TRUE
21	RopeStrengthResolution: 500
22	MinGearDensity: 1e-05
23	
24	PrintTables= TRUE
25	PrintDefaultMaps= TRUE, TRUE
26	PrintScenarioMaps= TRUE, TRUE
27	PrintRedistributionMaps= TRUE
28	PrintMapsInHighResolution= FALSE
29	TruncateMaps= 0.995
30	WriteMapSources= TRUE
31	WriteOutputCsv= TRUE
32	WriteDetailedOutput= FALSE
33	PrintSummary= TRUE
34	ArchiveInputSpreadsheet: FALSE

Final Relative Risk Scores – Mean Threat

	Variable	Month	Default	Scenario	Reduction
1	RelativeRisk_Threat	1	14,095.75	7,559.27	46.4 %
2	RelativeRisk_Threat	2	6,767.33	4,292.60	36.6 %
3	RelativeRisk_Threat	3	6,759.06	3,170.53	53.1 %
4	RelativeRisk_Threat	4	12,722.55	4,416.41	65.3 %
5	RelativeRisk_Threat	5	15,487.03	4,577.99	70.4 %
6	RelativeRisk_Threat	6	8,382.55	5,527.84	34.1 %
7	RelativeRisk_Threat	7	9,668.92	6,324.93	34.6 %
8	RelativeRisk_Threat	8	1,862.60	1,203.18	35.4 %
9	RelativeRisk_Threat	9	1,215.45	777.75	36 %
10	RelativeRisk_Threat	10	4,156.58	2,195.72	47.2 %
11	RelativeRisk_Threat	11	7,782.03	4,170.17	46.4 %
12	RelativeRisk_Threat	12	7,908.71	5,013.28	36.6 %
13	RelativeRisk_Threat	Total	96,808.56	49,229.66	49.1 %

April closure scenario in MRA Wedge, Alternative 2 proposed emergency restricted area

	ModelConfiguration
1	Model Start Time: 2023-01-09 14:08:03; End Time: 2023-01-09 14:38:31; Duration 00:30:28.2
2	Home Directory: //net/work4/LobsterGroup/Management/RightWhales/DecisionSupportTool
3	Model Version: DST_Beta_V4.1.0.R
4	InputActions: CrystalsSubfolder/EmergencyRule2023/ER_Pref_Clos_Apr_TP_wPh1v4.0.3.csv
5	MapRefDomain: MapRef_3.9.2.Rdata
6	GearMap: FisheryInputs_AllEastCoast_v4.0.1.Rdata
7	Threat Model: ThreatMod_RW_Selectivity_Uncertainty_WithZero.Rdata
8	Whale Habitat Model: Duke_RightWhaleModel_v12_1019.Rdata
9	Whale Vertical Dist File: RightWhaleVerticalDistribution_CSA_V1.1.Rdata
10	Whale Vertical Dist Model: Ensemble
11	Whale Dimensions: WhaleDimensions_AdHoc.Rdata
12	
13	Comment:
14	CoOccurrence: FALSE
15	Run Test Scenario: TRUE
16	Include Ground Gear: TRUE
17	AggregateStrings: FALSE
18	HighResolution: TRUE
19	RelocationCostExp: 1
20	ExpressRedistribution: TRUE
21	RopeStrengthResolution: 500
22	MinGearDensity: 1e-05
23	
24	PrintTables= TRUE
25	PrintDefaultMaps= TRUE, TRUE
26	PrintScenarioMaps= TRUE, TRUE
27	PrintRedistributionMaps= TRUE
28	PrintMapsInHighResolution= FALSE
29	TruncateMaps= 0.995
30	WriteMapSources= TRUE
31	WriteOutputCsv= TRUE
32	WriteDetailedOutput= FALSE
33	PrintSummary= TRUE
34	ArchiveInputSpreadsheet: FALSE

Final Relative Risk Scores – Mean Threat

	Variable	Month	Default	Scenario	Reduction
1	RelativeRisk_Threat	1	14,095.75	7,559.27	46.4 %
2	RelativeRisk_Threat	2	6,767.33	4,292.60	36.6 %
3	RelativeRisk_Threat	3	6,759.06	3,421.45	49.4 %
4	RelativeRisk_Threat	4	12,722.55	3,622.45	71.5 %
5	RelativeRisk_Threat	5	15,487.03	4,577.99	70.4 %
6	RelativeRisk_Threat	6	8,382.55	5,527.84	34.1 %
7	RelativeRisk_Threat	7	9,668.92	6,324.93	34.6 %
8	RelativeRisk_Threat	8	1,862.60	1,203.18	35.4 %
9	RelativeRisk_Threat	9	1,215.45	777.75	36 %
10	RelativeRisk_Threat	10	4,156.58	2,195.72	47.2 %
11	RelativeRisk_Threat	11	7,782.03	4,170.17	46.4 %
12	RelativeRisk_Threat	12	7,908.71	5,013.28	36.6 %
13	RelativeRisk_Threat	Total	96,808.56	48,686.62	49.7 %

Appendix 4.1.3 Alternative 3 Model Runs

Risk reduction presented in this section were run with a constraint for coastwide trap/pot fisheries. For section 6.2 of this EA, these results were then calculated relative to different baselines to reflect only risk reduction in addition to the previous rule based on the data presented in section 4.1.1 of this Appendix.

Gear Reduction (i.e. all lines removed)

February gear reduction in the MRA Wedge North, Alternative 3 proposed emergency restricted area

ModelConfiguration	
1	Model Start Time: 2023-01-09 13:31:43; End Time: 2023-01-09 14:24:35; Duration 00:52:52.8
2	Home Directory: //net/work4/LobsterGroup/Management/RightWhales/DecisionSupportTool
3	Model Version: DST_Beta_V4.1.0.R
4	InputActions: CrystalsSubfolder/EmergencyRule2023/ER_NonPref_GR_Feb_TP_wPh1v4.0.3.csv
5	MapRefDomain: MapRef_3.9.2.Rdata
6	GearMap: FisheryInputs_AllEastCoast_v4.0.1.Rdata
7	Threat Model: ThreatMod_RW_Selectivity_Uncertainty_WithZero.Rdata
8	Whale Habitat Model: Duke_RightWhaleModel_v12_1019.Rdata
9	Whale Vertical Dist File: RightWhaleVerticalDistribution_CSA_V1.1.Rdata
10	Whale Vertical Dist Model: Ensemble
11	Whale Dimensions: WhaleDimensions_AdHoc.Rdata
12	
13	Comment:
14	CoOccurrence: FALSE
15	Run Test Scenario: TRUE
16	Include Ground Gear: TRUE
17	AggregateStrings: FALSE
18	HighResolution: TRUE
19	RelocationCostExp: 1
20	ExpressRedistribution: TRUE
21	RopeStrengthResolution: 500
22	MinGearDensity: 1e-05
23	
24	PrintTables= TRUE
25	PrintDefaultMaps= TRUE, TRUE
26	PrintScenarioMaps= TRUE, TRUE
27	PrintRedistributionMaps= TRUE
28	PrintMapsInHighResolution= FALSE
29	TruncateMaps= 0.995
30	WriteMapSources= TRUE
31	WriteOutputCsv= TRUE
32	WriteDetailedOutput= FALSE
33	PrintSummary= TRUE
34	ArchiveInputSpreadsheet: FALSE

Final Relative Risk Scores – Mean Threat

	Variable	Month	Default	Scenario	Reduction
1	RelativeRisk_Threat	1	14,095.75	7,559.27	46.4 %
2	RelativeRisk_Threat	2	6,767.33	3,257.32	51.9 %
3	RelativeRisk_Threat	3	6,759.06	3,421.45	49.4 %
4	RelativeRisk_Threat	4	12,722.55	4,416.41	65.3 %
5	RelativeRisk_Threat	5	15,487.03	4,577.99	70.4 %
6	RelativeRisk_Threat	6	8,382.55	5,527.84	34.1 %
7	RelativeRisk_Threat	7	9,668.92	6,324.93	34.6 %
8	RelativeRisk_Threat	8	1,862.60	1,203.18	35.4 %
9	RelativeRisk_Threat	9	1,215.45	777.75	36 %
10	RelativeRisk_Threat	10	4,156.58	2,195.72	47.2 %
11	RelativeRisk_Threat	11	7,782.03	4,170.17	46.4 %
12	RelativeRisk_Threat	12	7,908.71	5,013.28	36.6 %
13	RelativeRisk_Threat	Total	96,808.56	48,445.31	50 %

March gear reduction in the MRA Wedge North, Alternative 3 proposed emergency restricted area

ModelConfiguration	
1	Model Start Time: 2023-01-09 14:05:36; End Time: 2023-01-09 15:08:17; Duration 01:02:24
2	Home Directory: //net/work4/LobsterGroup/Management/RightWhales/DecisionSupportTool
3	Model Version: DST_Beta_V4.1.0.R
4	InputActions: CrystalsSubfolder/EmergencyRule2023/ER_NonPref_GR_Mar_TP_wPh1v4.0.3.csv
5	MapRefDomain: MapRef_3.9.2.Rdata
6	GearMap: FisheryInputs_AllEastCoast_v4.0.1.Rdata
7	Threat Model: ThreatMod_RW_Selectivity_Uncertainty_WithZero.Rdata
8	Whale Habitat Model: Duke_RightWhaleModel_v12_1019.Rdata
9	Whale Vertical Dist File: RightWhaleVerticalDistribution_CSA_V1.1.Rdata
10	Whale Vertical Dist Model: Ensemble
11	Whale Dimensions: WhaleDimensions_AdHoc.Rdata
12	
13	Comment:
14	CoOccurrence: FALSE
15	Run Test Scenario: TRUE
16	Include Ground Gear: TRUE
17	AggregateStrings: FALSE
18	HighResolution: TRUE
19	RelocationCostExp: 1
20	ExpressRedistribution: TRUE
21	RopeStrengthResolution: 500
22	MinGearDensity: 1e-05
23	
24	PrintTables= TRUE
25	PrintDefaultMaps= TRUE, TRUE
26	PrintScenarioMaps= TRUE, TRUE
27	PrintRedistributionMaps= TRUE
28	PrintMapsInHighResolution= FALSE
29	TruncateMaps= 0.995
30	WriteMapSources= TRUE
31	WriteOutputCsv= TRUE
32	WriteDetailedOutput= FALSE
33	PrintSummary= TRUE
34	ArchiveInputSpreadsheet: FALSE

Final Relative Risk Scores – Mean Threat

	Variable	Month	Default	Scenario	Reduction
1	RelativeRisk_Threat	1	14,095.75	7,559.27	46.4 %
2	RelativeRisk_Threat	2	6,767.33	4,292.60	36.6 %
3	RelativeRisk_Threat	3	6,759.06	2,981.20	55.9 %
4	RelativeRisk_Threat	4	12,722.55	4,416.41	65.3 %
5	RelativeRisk_Threat	5	15,487.03	4,577.99	70.4 %
6	RelativeRisk_Threat	6	8,382.55	5,527.84	34.1 %
7	RelativeRisk_Threat	7	9,668.92	6,324.93	34.6 %
8	RelativeRisk_Threat	8	1,862.60	1,203.18	35.4 %
9	RelativeRisk_Threat	9	1,215.45	777.75	36 %
10	RelativeRisk_Threat	10	4,156.58	2,195.72	47.2 %
11	RelativeRisk_Threat	11	7,782.03	4,170.17	46.4 %
12	RelativeRisk_Threat	12	7,908.71	5,013.28	36.6 %
13	RelativeRisk_Threat	Total	96,808.56	49,040.33	49.3 %

April gear reduction in the MRA Wedge North, Alternative 3 proposed emergency restricted area

	ModelConfiguration
1	Model Start Time: 2023-01-09 11:58:35; End Time: 2023-01-09 12:48:28; Duration 00:49:52.2
2	Home Directory: //net/work4/LobsterGroup/Management/RightWhales/DecisionSupportTool
3	Model Version: DST_Beta_V4.1.0.R
4	InputActions: CrystalsSubfolder/EmergencyRule2023/ER_NonPref_GR_Apr_TP_wPh1v4.0.3.csv
5	MapRefDomain: MapRef_3.9.2.Rdata
6	GearMap: FisheryInputs_AllEastCoast_v4.0.1.Rdata
7	Threat Model: ThreatMod_RW_Selectivity_Uncertainty_WithZero.Rdata
8	Whale Habitat Model: Duke_RightWhaleModel_v12_1019.Rdata
9	Whale Vertical Dist File: RightWhaleVerticalDistribution_CSA_V1.1.Rdata
10	Whale Vertical Dist Model: Ensemble
11	Whale Dimensions: WhaleDimensions_AdHoc.Rdata
12	
13	Comment:
14	CoOccurrence: FALSE
15	Run Test Scenario: TRUE
16	Include Ground Gear: TRUE
17	AggregateStrings: FALSE
18	HighResolution: TRUE
19	RelocationCostExp: 1
20	ExpressRedistribution: TRUE
21	RopeStrengthResolution: 500
22	MinGearDensity: 1e-05
23	
24	PrintTables= TRUE
25	PrintDefaultMaps= TRUE, TRUE
26	PrintScenarioMaps= TRUE, TRUE
27	PrintRedistributionMaps= TRUE
28	PrintMapsInHighResolution= TRUE
29	TruncateMaps= 0.995
30	WriteMapSources= TRUE
31	WriteOutputCsv= TRUE
32	WriteDetailedOutput= FALSE
33	PrintSummary= TRUE
34	ArchiveInputSpreadsheet: FALSE

Final Relative Risk Scores – Mean Threat

	Variable	Month	Default	Scenario	Reduction
1	RelativeRisk_Threat	1	14,095.75	7,559.27	46.4 %
2	RelativeRisk_Threat	2	6,767.33	4,292.60	36.6 %
3	RelativeRisk_Threat	3	6,759.06	3,421.45	49.4 %
4	RelativeRisk_Threat	4	12,722.55	3,303.40	74 %
5	RelativeRisk_Threat	5	15,487.03	4,577.99	70.4 %
6	RelativeRisk_Threat	6	8,382.55	5,527.84	34.1 %
7	RelativeRisk_Threat	7	9,668.92	6,324.93	34.6 %
8	RelativeRisk_Threat	8	1,862.60	1,203.18	35.4 %
9	RelativeRisk_Threat	9	1,215.45	777.75	36 %
10	RelativeRisk_Threat	10	4,156.58	2,195.72	47.2 %
11	RelativeRisk_Threat	11	7,782.03	4,170.17	46.4 %
12	RelativeRisk_Threat	12	7,908.71	5,013.28	36.6 %
13	RelativeRisk_Threat	Total	96,808.56	48,367.57	50 %

Closure (i.e. all lines relocated)

February closure in the MRA Wedge North, Alternative 3 proposed emergency restricted area

ModelConfiguration	
1	Model Start Time: 2023-01-09 11:47:01; End Time: 2023-01-09 12:37:11; Duration 00:50:10.2
2	Home Directory: //net/work4/LobsterGroup/Management/RightWhales/DecisionSupportTool
3	Model Version: DST_Beta_V4.1.0.R
4	InputActions: CrystalsSubfolder/EmergencyRule2023/ER_NonPref_Clos_Feb_TP_wPh1v4.0.3.csv
5	MapRefDomain: MapRef_3.9.2.Rdata
6	GearMap: FisheryInputs_AllEastCoast_v4.0.1.Rdata
7	Threat Model: ThreatMod_RW_Selectivity_Uncertainty_WithZero.Rdata
8	Whale Habitat Model: Duke_RightWhaleModel_v12_1019.Rdata
9	Whale Vertical Dist File: RightWhaleVerticalDistribution_CSA_V1.1.Rdata
10	Whale Vertical Dist Model: Ensemble
11	Whale Dimensions: WhaleDimensions_AdHoc.Rdata
12	
13	Comment:
14	CoOccurrence: FALSE
15	Run Test Scenario: TRUE
16	Include Ground Gear: TRUE
17	AggregateStrings: FALSE
18	HighResolution: TRUE
19	RelocationCostExp: 1
20	ExpressRedistribution: TRUE
21	RopeStrengthResolution: 500
22	MinGearDensity: 1e-05
23	
24	PrintTables= TRUE
25	PrintDefaultMaps= TRUE, TRUE
26	PrintScenarioMaps= TRUE, TRUE
27	PrintRedistributionMaps= TRUE
28	PrintMapsInHighResolution= TRUE
29	TruncateMaps= 0.995
30	WriteMapSources= TRUE
31	WriteOutputCsv= TRUE
32	WriteDetailedOutput= FALSE
33	PrintSummary= TRUE
34	ArchiveInputSpreadsheet: FALSE

Final Relative Risk Scores – Mean Threat

	Variable	Month	Default	Scenario	Reduction
1	RelativeRisk_Threat	1	14,095.75	7,559.27	46.4 %
2	RelativeRisk_Threat	2	6,767.33	3,649.28	46.1 %
3	RelativeRisk_Threat	3	6,759.06	3,421.45	49.4 %
4	RelativeRisk_Threat	4	12,722.55	4,416.41	65.3 %
5	RelativeRisk_Threat	5	15,487.03	4,577.99	70.4 %
6	RelativeRisk_Threat	6	8,382.55	5,527.84	34.1 %
7	RelativeRisk_Threat	7	9,668.92	6,324.93	34.6 %
8	RelativeRisk_Threat	8	1,862.60	1,203.18	35.4 %
9	RelativeRisk_Threat	9	1,215.45	777.75	36 %
10	RelativeRisk_Threat	10	4,156.58	2,195.72	47.2 %
11	RelativeRisk_Threat	11	7,782.03	4,170.17	46.4 %
12	RelativeRisk_Threat	12	7,908.71	5,013.28	36.6 %
13	RelativeRisk_Threat	Total	96,808.56	48,837.26	49.6 %

March closure in the MRA Wedge North, Alternative 3 proposed emergency restricted area

	ModelConfiguration
1	Model Start Time: 2023-01-09 15:35:15; End Time: 2023-01-09 16:07:42; Duration 00:32:26.4
2	Home Directory: //net/work4/LobsterGroup/Management/RightWhales/DecisionSupportTool
3	Model Version: DST_Beta_V4.1.0.R
4	InputActions: CrystalsSubfolder/EmergencyRule2023/ER_NonPref_Clos_Mar_TP_wPh1v4.0.3.csv
5	MapRefDomain: MapRef_3.9.2.Rdata
6	GearMap: FisheryInputs_AllEastCoast_v4.0.1.Rdata
7	Threat Model: ThreatMod_RW_Selectivity_Uncertainty_WithZero.Rdata
8	Whale Habitat Model: Duke_RightWhaleModel_v12_1019.Rdata
9	Whale Vertical Dist File: RightWhaleVerticalDistribution_CSA_V1.1.Rdata
10	Whale Vertical Dist Model: Ensemble
11	Whale Dimensions: WhaleDimensions_AdHoc.Rdata
12	
13	Comment:
14	CoOccurrence: FALSE
15	Run Test Scenario: TRUE
16	Include Ground Gear: TRUE
17	AggregateStrings: FALSE
18	HighResolution: TRUE
19	RelocationCostExp: 1
20	ExpressRedistribution: TRUE
21	RopeStrengthResolution: 500
22	MinGearDensity: 1e-05
23	
24	PrintTables= TRUE
25	PrintDefaultMaps= TRUE, TRUE
26	PrintScenarioMaps= TRUE, TRUE
27	PrintRedistributionMaps= TRUE
28	PrintMapsInHighResolution= FALSE
29	TruncateMaps= 0.995
30	WriteMapSources= TRUE
31	WriteOutputCsv= TRUE
32	WriteDetailedOutput= FALSE
33	PrintSummary= TRUE
34	ArchiveInputSpreadsheet: FALSE

Final Relative Risk Scores – Mean Threat

	Variable	Month	Default	Scenario	Reduction
1	RelativeRisk_Threat	1	14,095.75	7,559.27	46.4 %
2	RelativeRisk_Threat	2	6,767.33	4,292.60	36.6 %
3	RelativeRisk_Threat	3	6,759.06	3,230.76	52.2 %
4	RelativeRisk_Threat	4	12,722.55	4,416.41	65.3 %
5	RelativeRisk_Threat	5	15,487.03	4,577.99	70.4 %
6	RelativeRisk_Threat	6	8,382.55	5,527.84	34.1 %
7	RelativeRisk_Threat	7	9,668.92	6,324.93	34.6 %
8	RelativeRisk_Threat	8	1,862.60	1,203.18	35.4 %
9	RelativeRisk_Threat	9	1,215.45	777.75	36 %
10	RelativeRisk_Threat	10	4,156.58	2,195.72	47.2 %
11	RelativeRisk_Threat	11	7,782.03	4,170.17	46.4 %
12	RelativeRisk_Threat	12	7,908.71	5,013.28	36.6 %
13	RelativeRisk_Threat	Total	96,808.56	49,289.89	49.1 %

April closure scenario in MRA Wedge North, Alternative 3 proposed emergency restricted area

ModelConfiguration	
1	Model Start Time: 2023-01-09 11:40:28; End Time: 2023-01-09 12:29:36; Duration 00:49:08.4
2	Home Directory: //net/work4/LobsterGroup/Management/RightWhales/DecisionSupportTool
3	Model Version: DST_Beta_V4.1.0.R
4	InputActions: CrystalsSubfolder/EmergencyRule2023/ER_NonPref_Clos_Apr_TP_wPh1v4.0.3.csv
5	MapRefDomain: MapRef_3.9.2.Rdata
6	GearMap: FisheryInputs_AllEastCoast_v4.0.1.Rdata
7	Threat Model: ThreatMod_RW_Selectivity_Uncertainty_WithZero.Rdata
8	Whale Habitat Model: Duke_RightWhaleModel_v12_1019.Rdata
9	Whale Vertical Dist File: RightWhaleVerticalDistribution_CSA_V1.1.Rdata
10	Whale Vertical Dist Model: Ensemble
11	Whale Dimensions: WhaleDimensions_AdHoc.Rdata
12	
13	Comment:
14	CoOccurrence: FALSE
15	Run Test Scenario: TRUE
16	Include Ground Gear: TRUE
17	AggregateStrings: FALSE
18	HighResolution: TRUE
19	RelocationCostExp: 1
20	ExpressRedistribution: TRUE
21	RopeStrengthResolution: 500
22	MinGearDensity: 1e-05
23	
24	PrintTables= TRUE
25	PrintDefaultMaps= TRUE, TRUE
26	PrintScenarioMaps= TRUE, TRUE
27	PrintRedistributionMaps= TRUE
28	PrintMapsInHighResolution= TRUE
29	TruncateMaps= 0.995
30	WriteMapSources= TRUE
31	WriteOutputCsv= TRUE
32	WriteDetailedOutput= FALSE
33	PrintSummary= TRUE
34	ArchiveInputSpreadsheet: FALSE

Final Relative Risk Scores – Mean Threat

	Variable	Month	Default	Scenario	Reduction
1	RelativeRisk_Threat	1	14,095.75	7,559.27	46.4 %
2	RelativeRisk_Threat	2	6,767.33	4,292.60	36.6 %
3	RelativeRisk_Threat	3	6,759.06	3,421.45	49.4 %
4	RelativeRisk_Threat	4	12,722.55	3,706.10	70.9 %
5	RelativeRisk_Threat	5	15,487.03	4,577.99	70.4 %
6	RelativeRisk_Threat	6	8,382.55	5,527.84	34.1 %
7	RelativeRisk_Threat	7	9,668.92	6,324.93	34.6 %
8	RelativeRisk_Threat	8	1,862.60	1,203.18	35.4 %
9	RelativeRisk_Threat	9	1,215.45	777.75	36 %
10	RelativeRisk_Threat	10	4,156.58	2,195.72	47.2 %
11	RelativeRisk_Threat	11	7,782.03	4,170.17	46.4 %
12	RelativeRisk_Threat	12	7,908.71	5,013.28	36.6 %
13	RelativeRisk_Threat	Total	96,808.56	48,770.27	49.6 %



UNITED STATES DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
NATIONAL MARINE FISHERIES SERVICE
GREATER ATLANTIC REGIONAL FISHERIES OFFICE
55 Great Republic Drive
Gloucester, MA 01930

DATE: January 24, 2023

MEMORANDUM FOR: Michael Pentony
Regional Administrator

FROM: Marianne Ferguson
NEPA Policy Analyst

THROUGH: Timothy Cardiasmenos
NEPA Coordinator

CARDIASMENOS.TI
MOTHY.APOSTLE.1
267850247

Digitally signed by
CARDIASMENOS.TIMOTHY.APOS
TLE.1267850247
Date: 2023.01.24 16:51:33 -05'00'

SUBJECT: Final Clearance of the Environmental Assessment (EA) and
Finding of No Significant Impact (FONSI) for the Proposal to
Promulgate a 2023 Emergency Rule to Reduce Right Whale
Interactions with Trap/Pot Gear

Thank you for the opportunity to review the subject documents. All comments provided on the subject EA have been adequately addressed. National Environmental Policy Act (NEPA) staff reviewed the FONSI for the subject EA. We have determined that it complies with the requirements of NEPA and recommend you concur by signing below. We have no further comment on the EA or FONSI statement.

cc: Marisa Trego, PRD Jennifer Anderson, PRD
Jean Higgins, PRD Jennifer Goebel, PRD
Mark Capone, GCNE Colleen Coogan, PRD
Sam Duggan, GCNE
Scott Farley, GCNE

1. I concur. **Michael Pentony** Digitally signed by Michael Pentony
Date: 2023.01.25 13:22:39 -05'00' _____ Date

2. I do not concur. _____ Date

Attachments



FINDING OF NO SIGNIFICANT IMPACT
Under the National Environmental Policy Act
2023 Emergency Final Rule to
Reduce Right Whale Interactions with Trap/Pot Gear

I. Purpose of Finding of No Significant Impact (FONSI): The National Environmental Policy Act (NEPA) requires the preparation of an Environmental Impact Statement (EIS) for any proposal for a major federal action significantly affecting the quality of the human environment. 42 U.S.C. § 4332(C). The Council on Environmental Quality (CEQ) Regulations direct agencies to prepare a Finding of No Significant Impact (FONSI) when an action not otherwise excluded will not have a significant impact on the human environment. 40 CFR §§ 1500.4(b), 1500.5(b), & 1501.6. To evaluate whether a significant impact on the human environment is likely, the CEQ regulations direct agencies to analyze the potentially affected environment and the degree of the effects of the proposed action. 40 CFR § 1501.3(b). In doing so, agencies should consider the geographic extent of the affected area (i.e., national, regional, or local), the resources located in the affected area (40 CFR § 1501.3(b)(1)), and whether the project is considered minor or small-scale (NAO 216-6A CM, Appendix A-2). In considering the degree of effect on these resources, agencies should examine, as appropriate, short- and long-term effects, beneficial and adverse effects, and effects on public health and safety, as well as effects that would violate laws for the protection of the environment (40 CFR § 1501.3(b)(2)(i)-(iv); NAO 216-6A CM Appendix A-2 - A-3), and the magnitude of the effect (e.g., negligible, slight, moderate, high). CEQ identifies specific criteria for consideration. 40 CFR § 1501.3(b)(2)(i)-(iv). Each criterion is discussed below with respect to the proposed action and considered individually as well as in combination with the others.

In preparing this FONSI, we reviewed the **Environmental Assessment (EA) of a Proposal to Promulgate a 2023 Emergency Rule to Reduce Right Whale Interactions with Trap/Pot Gear** which evaluates the affected area, the scale and geographic extent of the proposed action, and the degree of effects on those resources (including the duration of impact, and whether the impacts were adverse and/or beneficial and their magnitude). The EA is hereby incorporated by reference. 40 CFR § 1501.6(b).

II. Approach to Analysis:

The proposed action is not considered to meaningfully contribute to a significant impact based on the scale and degree of impact. The alternatives for this emergency closure of the waters between state and federal portions of the Massachusetts Restricted Area (MRA) from February 1 to April 30, 2023 are expected to have minimal impacts on their own because the emergency closure area is small relative to the entire MRA, minimal economic impacts are expected, and any indirect effects are likely beneficial for the environment. The emergency closure would be of limited duration (i.e. February through April) further reducing impact.

The proposed action will not meaningfully contribute to significant impacts to specific resources. The Atlantic Large Whale Take Reduction Plan (Plan) was developed pursuant to section 118(f) of the MMPA to reduce the level of mortality and serious injury of large whales as a result of

trap/pot and gillnet commercial fishing gear. Modifications to the Plan are designed to meet established management goals and objectives. The expected impacts of the proposed action on the Valued Ecosystem Components (VECs) and associated analyses are described in Chapter 6 of the EA.

The Preferred Alternative for the Emergency Rule is expected to have impacts ranging from slight negative to moderate negative on protected species due to the continuation of fishing activity outside of the closure period (Section 6.2). The Preferred Alternative is expected to have impacts ranging from slight negative to negligible on habitat (Section 6.3) and slight negative impacts on human communities (Section 6.4). None of these conclusions, when considered together, are expected to result in any overall significant impact.

The proposed action is not connected to other actions that have caused or may cause effects to the resources in the affected area. Thus, there is no potential for the effects of the proposed action to add to the effects of other projects, such that the effects taken together could be significant, as described in the Cumulative Effects Analysis in Section 6.5.

III. Geographic Extent and Scale of the Proposed Action:

The proposed action establishes an emergency closure off the Atlantic coast to close the gap between the previous MRA and the expanded restricted area which includes Massachusetts State waters, and is therefore local in its geographic extent. The area proposed to be closed to the trap/pot fisheries in the Preferred Alternative is approximately 200 square miles (518 square kilometers) and begins in federal waters east of Cape Ann, is bounded landward by the Massachusetts state waters, east until it intersects with the MRA, and runs north until it intersects the state water boundary. The resources present throughout this region with the potential to be impacted by the Emergency Rule are described in Section 5.2 of the EA. The proposed action and its impacts focus on a small area for the months of February, March, and April. Due to being very narrow in time and geographic extent, the environmental effects for the VECs concerned in this action, focus on a very specific area. These effects are not expected to result in substantial changes to any VECs or specific geographic areas.

IV. Degree of Effect:

- A. The potential for the proposed action to threaten a violation of Federal, state, or local law or requirements imposed for environmental protection.*

The Emergency Rule is not expected to violate federal, state, or local environmental laws.

- B. The degree to which the proposed action is expected to affect public health or safety.*

The Emergency Rule is not expected to significantly affect public health and safety. This action does not modify current requirements in a way that would impact the potential safety of small vessel operators. Fishers were informed of the possibility of this closure and many knew to avoid setting gear here again this year. This impact is not considered significant because we expect that this action will either prevent fishing during this time in this area or allow fishing to continue according to current practices, rather than fish in an unsafe manner. Refer to Section 6.4.2 for a

description of the social impacts of the proposed action.

C. The degree to which the proposed actions is expected to affect a sensitive biological resource, including:

a. Federal threatened or endangered species and critical habitat;

Impacts to protected resources including endangered species from the proposed action are discussed in Section 6.2 of the EA. Subsection 6.2.5 identifies expected moderate to slight negative impacts to ESA-listed species from the alternatives proposed for the Emergency Rule. Specifically, this determination has been made because co-occurrence of ESA-listed species of large whales with buoy lines and associated entanglement risk will likely decline minimally to substantially, depending on the whale species. While entanglement risk is not completely eliminated, the action does significantly reduce risk in the action area for marine mammals. Overall, we consider the Emergency Rule Preferred Alternative to have moderate to slight negative impacts on ESA-listed large whales but is not expected to cause new effects to listed species or critical habitat that were not already considered in the previous ESA Section 7 consultations.

b. stocks of marine mammals as defined in the Marine Mammal Protection Act;

The Atlantic Large Whale Take Reduction Plan (Plan) was developed pursuant to section 118(f) of the Marine Mammal Protection Act to reduce the level of mortality and serious injury of large whales as a result of trap/pot and gillnet commercial fishing gear. In alignment with federal obligations under the Plan, the proposed action is not expected to adversely affect stocks of marine mammals (Section 6.2). Specifically, this determination has been made because the overlap between MMPA-protected large whales with buoy lines and associated entanglement risk is expected to decrease. While entanglement risk is not completely eliminated, the action does reduce risk in the action area for marine mammals. Based on all of this information, this action is not expected to adversely affect stocks of marine mammals as defined in the Marine Mammal Protection Act.

c. essential fish habitat identified under the Magnuson–Stevens Fishery Conservation and Management Act;

The proposed action is not expected to cause substantial damage to EFH as defined under the MSA and identified in the American lobster and Jonah crab FMP. Because trap/pot gear will be removed from areas where the habitat is currently disturbed, the Preferred Alternative is expected to reduce the impact of lobster and Jonah crab fishing on the habitat VEC. Operational trap/pot gear is not considered to cause long-term benthic impacts, but the removal of traps may decrease benthic community disturbance, protect local community structure, and may increase local lobster and/or Jonah crab abundance (Uhrin 2016). The impacts of this action's alternatives are all expected to be negligible to slight negative.

d. bird species protected under the Migratory Bird Treaty Act;

Information about seabird interactions with trap/pot gear is limited. However, there is no known evidence of substantial impacts to bird species, including those protected under the Migratory

Bird Treaty Act, from the Northeast trap/pot fisheries in the past. The Preferred Alternative will remove gear from the closure areas and, as a result, this action would only reduce risks to these species.

e. national marine sanctuaries or monuments;

The Stellwagen Bank National Marine Sanctuary and Northeast Canyons and Seamounts Marine National Monument occur within the broad management areas of the Plan, however they are not within the area considered for the emergency closure. As a result, the Emergency Rule is not expected to have any substantial effects on national marine sanctuaries or monuments.

f. vulnerable marine or coastal ecosystems, including, but not limited to, shallow or deep coral ecosystems;

The Preferred Alternative is not expected to have significant impacts on the natural or physical environment, including vulnerable marine or coastal ecosystems. The Preferred Alternative will remove gear from the emergency closure area and is not likely to result in a significant redistribution of gear and any gear that is relocated is not expected to be moved to areas not already heavily impacted by commercial fishing. No additional trap/pot fishing will take place near the continental slope/shelf break where deep sea corals may be found in and around the submarine canyons. The Preferred Alternative is not expected to alter American lobster or Jonah crab fishing patterns relative to this protected area or in any other manner that would lead to adverse impacts on deep sea coral or other vulnerable marine or coastal ecosystems.

g. biodiversity or ecosystem functioning (e.g., benthic productivity, predator-prey relationships, etc.)

The impacts of the Emergency Rule on biodiversity and ecosystem functioning have not been assessed; however, the impacts to components of the ecosystem (i.e., protected species, habitat) have been considered. The proposed action is not expected to have a substantial impact on biodiversity and ecosystem function within the affected area. Additionally, the proposed action is not expected to substantially alter fishing methods or activities or fishing effort or the spatial and/or temporal distribution of current fishing effort outside of the emergency closure area.

D. The degree to which the proposed action is reasonably expected to affect a cultural resource: properties listed or eligible for listing on the National Register of Historic Places; archeological resources (including underwater resources); and resources important to traditional cultural and religious tribal practice.

The Emergency Rule is not likely to affect districts, sites, highways, structures, or objects listed in or eligible for listing in the National Register of Historic Places or cause loss or destruction of significant scientific, cultural, or historical resources. Although there are shipwrecks present in areas where fishing occurs, including some registered on the National Register of Historic Places, vessels try to avoid fishing too close to wrecks due to the possible loss or hang ups of fishing gear. Therefore, it is not likely that the Emergency Rule would adversely affect the historic resources.

- E. The degree to which the proposed action has the potential to have a disproportionately high and adverse effect on the health or the environment of minority or low-income communities, compared to the impacts on other communities (EO 12898).*

The proposed action is not expected to have a disproportionately high and adverse effect on the health or the environment of minority or low-income communities, compared to the impacts on other communities. As described in Section 5.3 and 6.4.2 while some communities expected to be affected by this proposed amendment may have social vulnerabilities that exceed the environmental justice (EJ) thresholds and, therefore, may constitute areas of concern, significant EJ issues are not expected to arise as a result of this proposed amendment. It is anticipated that the impacts from the proposed regulations may impact minorities or the poor, but not through discriminatory application of these regulations. However, it should be noted that some communities are not able to be analyzed as census data are not available to create the indices. There are no known claims for customary usage or subsistence consumption of American lobster or Jonah crab, the primary fishery impacted by this action, by any population including tribes or indigenous groups.

- F. The degree to which the proposed action is likely to result in effects that contribute to the introduction, continued existence, or spread of noxious weeds or nonnative invasive species known to occur in the area or actions that may promote the introduction, growth, or expansion of the range of the species.*

The Emergency Rule would not result in the introduction or spread of non-indigenous species. The Emergency Rule will not result in U.S. vessels leaving regional waters, or result in foreign vessels operating in U.S. waters.

- G. The potential for the proposed action to cause an effect to any other physical or biological resources where the impact is considered substantial in magnitude (e.g., irreversible loss of coastal resources such as marshland or seagrass) or over which there is substantial uncertainty or scientific disagreement.]*

The proposed action is not expected to cause a substantial effect to any other physical or biological resource, nor is there substantial uncertainty or scientific disagreement on the impacts of the proposed actions. As described in Chapter 6, it is not anticipated that there is potential for the proposed actions to cause an effect to any other physical or biological resource where the impact is considered substantial because the proposed actions are not expected to change the way the Northeast trap/pot fisheries are prosecuted.

V. Other Actions Including Connected Actions :

There are no other connected actions (40 CFR § 1501.9(e)(1)). Any other future actions within the Plan would be developed, analyzed, and implemented independently of the proposed action. The Cumulative Effects Analysis in Section 6.5 of the EA discusses other beneficial and adverse actions that are occurring or reasonably certain to occur, and that affect the same resources as the Proposed Action. This section of the EA demonstrates that the effects of these collective actions,

for each resource analyzed, do not result in synergistically significant impacts, either positive or negative.

VI. Mitigation and Monitoring:

NMFS anticipates continued implementation of existing conservation and mitigation measures which are summarized in Section 3.2.1.2 and Appendix 2.1 of the 2021 Final Environmental Impact Statement associated with the 2021 modifications to the Atlantic Large Whale Take Reduction Plan (86 FR 51970). NMFS has not identified additional practicable mitigation measures that would be effective at avoiding, minimizing, or mitigating adverse effects likely to be caused by the Proposed Action. Therefore, NMFS is not proposing or adopting any mitigation measures.

DETERMINATION

The CEQ NEPA regulations, 40 CFR § 1501.6, direct an agency to prepare a FONSI when the agency, based on the EA for the proposed action, determines not to prepare an EIS because the action will not have significant effects. In view of the information presented in this document and the analysis contained in the supporting EA prepared for a 2023 Emergency Rule to Reduce Right Whale Interactions with Trap/Pot Gear, it is hereby determined that the 2023 Emergency Final Rule will not significantly impact the quality of the human environment. The 2023 Emergency Final Rule is hereby incorporated by reference. In addition, all beneficial and adverse impacts of the proposed action, as well as mitigation measures, have been evaluated to reach the conclusion of no significant impacts. Accordingly, preparation of an EIS for this action is not necessary.

**Michael
Pentony**

Digitally signed by Michael
Pentony
Date: 2023.01.25 13:24:09
-05'00'

Regional Administrator
GARFO, NMFS, NOAA

Date

REGULATORY IMPACT REVIEW OF: 2023 EMERGENCY FINAL RULE TO REDUCE RIGHT WHALE INTERACTIONS WITH TRAP/POT GEAR

1. Basis for and Purpose of the Rule

The Atlantic Large Whale Take Reduction Plan (Plan) was developed pursuant to section 118(f) of the Marine Mammal Protection Act (MMPA), to reduce the level of mortality and serious injury of large whales as a result of trap/pot and gillnet commercial fishing gear. After the 1994 amendments to the MMPA, National Marine Fisheries Service (NMFS) created the Atlantic Large Whale Take Reduction Team in 1996 and developed the first Plan which published its implementing regulations on July 22, 1997 (62 FR 39157). The Team consists of stakeholders representing state and federal government agencies, fishing industry, conservation organizations, and researchers. For a more detailed management history of the Plan and management of fishery interactions, please see the Final Environmental Impact Statement (FEIS) accompanying the 2021 amendment to the Plan (NMFS 2021).

One measure included in the 2021 Final Rule (86 FR 51970, September 17, 2021) has left a critical gap in protection where North Atlantic right whales (*Eubalaena glacialis*, hereafter referred to as right whale) distribution information identifies a high risk of overlap between right whales and buoy lines. Right whale monthly distribution data identifies risk in unrestricted waters encapsulated on three sides by the expanded Massachusetts Restricted Area (MRA) during the months of February, March, and April. The 2021 expansion of the geographic extent of the MRA under the Plan to include Massachusetts state waters north to the New Hampshire border (Figure 1) mirrors the Massachusetts State 2021 modification of the state water closure (322 CMR 12.04(2)).

Outside of the boundaries of the MRA, approximately 200 square miles (518 square kilometers) of federal waters remain open to trap/pot fishing between state and federal waters of the closure, creating a wedge (“MRA Wedge”) where 2021 and 2022 data indicates that trap/pot gear is concentrated during the months of February, March, and April (Figure 1 and Figure 2). During aerial surveys in April 2021 and February and March of 2022, the Center for Coastal Studies (CCS) and Northeast Fisheries Science Center (NEFSC) observed right whales and aggregated fishing gear within the wedge. Fishermen that used this area are likely to be using the remaining open waters to fish trap/pot gear and also could be staging their gear in preparation for the opening of the federal waters portion of the MRA on May 1. The high gear density observed in this area outside of the MRA has created an area of high risk of right whale entanglement. In early 2022, NMFS received letters and emails from Massachusetts Division of Marine Fisheries (MA DMF), Stellwagen Bank National Marine Sanctuary, and non-governmental organizations expressing concerns about this gap in restricted waters and the heightened risk of entanglement for right whales. After reviewing available information and due to the high risk of entanglement in this relatively small area, NMFS issued an emergency rule prohibiting trap/pot fishery buoy

lines between federal and state waters within the MRA for the month of April in 2022 (87 FR 11590; March 2, 2022).

In letters dated December 12, 2022 and January 4, 2023, MA DMF again expressed the concern that the continued overlap of persistent buoy lines with aggregations of right whales poses an entanglement threat. MA DMF also expressed the concern that any right whale entanglement in waters off the Massachusetts coast could threaten the opportunity of Massachusetts-based fishermen to participate in fixed-gear fisheries.

The purpose of this Emergency Rule, consistent with the March 2022 emergency rule, is to reduce the acute risk of right whales to entanglement with trap/pot fisheries in waters adjacent to the existing MRA where there is high overlap between right whale aggregations and buoy lines. There is an urgent need to prevent any mortality or serious injury of right whales in the U.S. commercial fisheries because any take is above the Potential Biological Removal (PBR) of 0.7 serious injuries or mortalities a year, established under the Marine Mammal Protection Act (MMPA) for this population. Implementing an emergency restriction to fishing with buoy lines in this area will address a critical gap in entanglement risk where there is a particularly high chance of entanglement in 2023 that was not addressed in recent modifications to the Plan, while a rule implementing this area as a permanent seasonal closure to buoy lines is in development.

Figure 1: The Massachusetts Restricted Area (green shaded area), Massachusetts state waters (crosshatched), and MRA Wedge (simple hatched gray shaded area) are represented. Federal and state waters within the Massachusetts Restricted Area are seasonally closed to the use of vertical buoy lines from February 1 through April 30. Massachusetts State regulations prohibit trap/pot fishing in any waters under the jurisdiction of the Commonwealth from February 1 through May 15, but can be extended the seasonal closure past May 15 in the continued presence of North Atlantic Right whales or rescinded after April 30 in their absence (322 CMR 12.04(2)). The MRA Wedge is closed by this emergency rule in Alternative 2 (Preferred Alternative). MRA Wedge would remain closed to trap/pot fishing with buoy lines from February 1, 2023 through April 30, 2023.





Figure 2: The Massachusetts Restricted Area “Wedge” closed by this rule is represented in pale gray. Fishing gear observed by the Center for Coastal Studies (CCS) on April 19, 2021 (white circles), April 28, 2021 (encircled “x”), February 6, 2022 (open diamonds), and March 11, 2022 (black diamonds), were selected as representative snapshots of fishing gear present in survey areas. Surveys concentrate on Cape Cod Bay; surveyors rarely fly north of mid Cape Ann, offshore Rockport, MA. These maps are used for qualitative not quantitative comparison, and differ from Decision Support Tool data.



1.1 Affected Fisheries

As required by the MMPA, NMFS maintains a List of Fisheries that places each commercial fishery into one of three categories. Fisheries are categorized according to the level of mortality and serious injury of marine mammals that occurs incidental to that fishery. The categorization of a fishery in the List of Fisheries determines whether participants in that fishery are subject to

certain provisions of the MMPA such as registration, observer coverage, and take reduction plan requirements. Individuals fishing in Category I or II fisheries must comply with requirements of any applicable take reduction plan.

Category I fisheries are associated with frequent incidental mortality and serious injury of marine mammals. These fisheries have a mortality and serious injury rate of 50 percent or more of a stock's potential biological removal rate. Category II fisheries are associated with occasional incidental mortality and serious injury of marine mammals, and have a serious injury/mortality rate of more than 1 percent but less than 50 percent of a stock's PBR. Category III fisheries rarely cause serious injury or mortality to marine mammals. Category III fisheries have a serious injury/mortality rate of 1 percent or less of a stock's PBR (NOAA 2002).

The List of Fisheries indicates which fisheries NMFS may regulate under the Plan. Specific fisheries were initially identified for inclusion under the Plan based on documented whale interactions. In 1996, NMFS announced its intention to regulate the Gulf of Maine, U.S. mid-Atlantic lobster trap/pot fishery, U.S. mid-Atlantic coastal gillnet fishery, New England multispecies sink-gillnet fishery, and Southeastern U.S. Atlantic shark gillnet fishery (61 FR 40819-40821).

This list has evolved since 1996, reflecting both changes in nomenclature and modification of the Plan to address additional fisheries. This Emergency Rule focuses on trap/pot fisheries within the vicinity of the emergency closure in LMA 1, particularly Massachusetts permitted lobster and Jonah crab vessels that are most likely to be impacted by this measure during the months the emergency closure would be in effect.

1.2 Regulatory Alternatives

The scope of this analysis is limited to two alternatives that would temporarily modify the Plan to create an emergency restricted area to supplement the existing MRA during the months of February, March, and April 2023. This analysis only affects a small portion of federal waters within Lobster Management Area 1 (LMA 1) in the northeast portion of the waters covered under the Plan. The Alternatives were selected based on surveys conducted by the CCS and the NEFSC that observed right whales and/or fixed fishing gear adjacent to the MRA in April 2021 and February and March 2022, as well as modeling conducted using a Decision Support Tool (DST). The DST is described in Appendix 3.1 Volume II of the 2021 FEIS (NMFS 2021) and the analyses in this document are further documented in section 4.1 of the associated Environmental Assessment (EA).


Alternative 1: No Action (Status Quo)

Alternative 1, “No Action,” leaves the current Plan intact with no regulatory changes proposed. This includes the restricted areas implemented by the Final Rule on September 17, 2021 (86 FR 51970) that went into effect October 18, 2021 and requirements for minimum traps per trawl and weak inserts throughout the buoy line that went into effect May 1, 2022.

Alternative 2: Preferred

Alternative 2, the Preferred Alternative, implements an emergency closure that restricts the use of persistent trap/pot gear buoy lines from February 1, 2023 through April 30, 2023. The restricted area is approximately 200 square miles (518 square kilometers) and begins in federal waters east of Cape Ann. It is bounded landward by the Massachusetts state waters and south along the 70°30' W longitude line until it intersects with the MRA at the 42°12'N latitude line, and runs west along that line until it intersects the state water boundary (MRA Wedge; Figure 3). Authorizations for fishing without buoy lines using ropeless gear in the MRA Wedge during this time must be obtained through an Exempted Fishing Permit.

Figure 3: The area closed by this emergency rule in the Preferred Alternative, MRA Wedge, is the shaded green area. The existing MRA is the hatched area. The boundary of MRA Wedge would remain closed to trap/pot fishing with buoy lines from February 1, 2023 through April 30, 2023.




Alternative 3: Non-Preferred Alternative

Alternative 3, the Non-Preferred Alternative also implements an emergency closure that restricts the use of persistent trap/pot gear buoy lines from February 1, 2023 through April 30, 2023. The closed area is approximately 1,300 square miles (3,367 square kilometers) and extends the northern LMA 1 MRA boundaries up to the New Hampshire border at 42°52.58' N (MRA

Wedge North; Figure 4.2). Authorizations for fishing without buoy lines using ropeless gear in the MRA Wedge North during this time must be obtained through an Exempted Fishing Permit.

Figure 4: The area closed by this emergency rule in the Non-Preferred Alternative, MRA Wedge North to New Hampshire, is the shaded purple area. The existing MRA is the hatched area. The boundary of MRA Wedge North would remain closed to trap/pot fishing with buoy lines from February 1, 2023 through April 30, 2023.



2. Regulatory Impact Review

2.1. Economic Baseline for Comparison

The baseline for the economic analysis is Alternative 1, which requires no action. The number of fishing vessels and fisheries landing data used for this economic analysis are the averages from 2017 to 2021. For Alternative 2, it is estimated that 26-31 trap/pot vessels in the Northeast Region would be affected by the modification of the Plan. Alternative 3 would affect 53-66 vessels.

2.2. *Time Horizon*

The rule will become effective on February 1, or upon publication if later than February 1, and will only be in effect for the months of February, March and April, 2023. Therefore, the costs and benefits will only be estimated for that period.

2.3. *Benefit-Cost Framework*

Benefit-cost analysis (BCA) is the preferred method for analyzing the consequences of a regulatory action such as modifying the requirements of the Plan. BCA is a well-established procedure for assessing the "best" course or scale of action, where "best" is that course which maximizes net benefits (i.e., benefits minus costs). Because BCA assesses the value of an activity in net benefit terms, it requires that a single metric, most commonly dollars, be used to gauge both benefits and costs. The data and economic models necessary to estimate costs may be difficult or costly to gather and develop, and a comprehensive analysis of the costs associated with a regulatory action is not always feasible. Nonetheless, the principle is straightforward, and it is generally possible in practice to develop a monetary estimate of at least some portion of regulatory costs. This is the case for costs stemming from temporary changes to the Plan, which would impose additional restrictions on commercial fishing operations.

Assessing the benefits of a temporary change to the Plan in a BCA framework is also straightforward in principle, but much more difficult in practice. To the extent that new regulations would reduce the risk that whales will die or suffer a serious injury as a result of entanglement in commercial fishing gear, the Emergency Rule would produce real benefits. Ideally, these benefits would be measured first by a biological metric, and then by a dollar metric. A biological metric could take the form of the percentage of risk reduction, the associated expected decrease in extinction risk, increase in the annual growth of the population, or similar measures. Reduced mortalities and reduced nonfatal injuries would matter; decreased extinction risk would matter independently.

The value of protecting right whales might not be adequately or fully captured by people's willingness to pay to protect whales. Their own welfare loss might be deemed relevant, regardless of that figure. In the Plan, moreover, the data required to complete such an analysis are not available. Estimation of the economic benefits attributable to each of the regulatory alternatives that NMFS is considering would require a more detailed understanding of the biological impacts of each measure than current models can provide. It also would require more extensive research than economists have conducted to date on the relationship between conservation and restoration of these species and associated economic values.

In the absence of the information required to conduct a full BCA, the discussion that follows presents a quantitative indicator of the potential impact of each alternative. It then presents estimates of the costs attributable to each alternative. As discussed later in this Regulatory Impact Review, the analysis uses this information to evaluate the cost of each percent of risk reduction of the regulatory alternatives under consideration, where risk is a product of co-occurrence combined with gear strength. Because the alternatives vary with respect to the benefits they would achieve, it is not possible to identify a superior option based on cost per unit

of risk reduction alone. Nonetheless, the cost for each percent of risk reduction figures provide a useful means of comparing the relative impacts of the regulatory provisions that each alternative incorporates.

2.4. *Economic Analysis of Alternatives*

2.4.1. **Benefits of the Alternatives**

Although it is difficult to calculate the monetary benefits of whale protection from this short-term Emergency Rule, we could estimate the effects on risk reduction for each Alternative.

Under Alternative 1 (No Action), the current Plan management regime consisting of time/area closures, minimum trap per trawl requirements, use of weak links in the surface system, and gear marking requirements remains in place. Therefore, the restricted areas included in the 2021 Final Rule are considered part of the status quo for this proposed action. Under Alternative 1, high negative impacts are expected because there would be a risk of entanglement due to the present number of buoy lines that would remain in the water when right whales are abundant in the area.

Trap/pot gears were observed in the LMA1 waters adjacent to the MRA in April 2021, and February and March 2022 by CCS and NEFSC aerial surveys. During these months, it is estimated that 188.45 right whales inhabit the Massachusetts portion of LMA 1, half of these in the month of April, confirming the high density aggregations essential to right whale foraging and survival within Massachusetts and Cape Cod Bay. No gear should be present in Massachusetts state waters during the emergency closure since it went into effect in 2021.

The DST estimates that the total number of right whales within the MRA Wedge is up to 5 whales within the 200 square miles (518 square kilometers) area from February through April. Within 1,297 square miles (3,359 square kilometers) MRA Wedge North, the right whale density from February through April is estimated to be 14 (Table 1). However, this whale density is likely an underestimate and is far below the number of whales sighted during April 2021 and February and March 2022 by the CCS and NEFSC (Figure 2).

Table 1: Mean large whale density for each Alternative in February through April.

Area	Area		Right Whale			Humpback Whale			Fin Whale		
	Sq Mi	Sq km	Feb	Mar	Apr	Feb	Mar	Apr	Feb	Mar	Apr
MRA Wedge (Preferred)	200	518	0.04	1.4	3.3	0.2	0.2	0.9	0.4	0.3	0.8
MRA Wedge North (Non-Preferred)	1,297	3,359	6.7	2.4	4.6	2	2.4	17.6	8.3	7.1	7.6

Removing lines provides benefits to right whales present during the months of February, March, and April. If lines are removed from the restricted areas, the MRA Wedge emergency closure reduces the annual risk by 13.2 to 16.6 percent, and the MRA Wedge North would reduce the

risk by 22.3 to 37.4 percent. Reducing the risk that a right whale encounters buoy rope reduces the potential for an entanglement incident that could lead to mortality or serious injury.

2.4.2. Costs of the Alternatives

We analyzed three alternatives for this Emergency Rule. Alternative 1 leaves the provisions of the Plan unchanged, and thus would have no economic impacts relative to current regulatory requirements. Alternative 2, the Preferred Alternative, proposes to temporarily expand the Massachusetts Restricted Area (MRA) to include the MRA Wedge northward to Cape Ann, Massachusetts from February to April 2023. It would add 200 square miles (518 square kilometers) to the current MRA and bring short-term negative economic impacts to a number of lobster vessels in Southern Essex County, Suffolk County, Norfolk County and Northern Plymouth County. Alternative 3, the Non-Preferred Alternative, expands the boundary of the MRA from February to April 2023 to include a larger area of approximately 1,297 square miles (3,359 square kilometers) that is bounded landward by the Massachusetts state waters, north at 42°52.58', seaward at 69°45', and south along the northern border of the MRA. Alternative 3 would impact additional vessels in Northern Essex County compared to Alternative 2.

The following section gives an overview of the analytic approach and results of economic impacts.

Analytic Approach

Vessels that fished within the proposed restricted area have two options to comply with the Emergency Rule: relocate their traps to the northern waters adjacent to the closure and keep fishing, or bring their traps back to dock and suspend fishing activity. VTR gear distribution data from the past few seasons show that at least half of the traps were placed at the southern portion of the restricted areas proposed in Alternative 2 and Alternative 3. Because these traps are about 20 to 30 miles from the northern boundary of the proposed restricted area, they would be difficult to relocate, as the distance is beyond the normal range of fishing vessels.

For this analysis, we evaluated two scenarios for the economic impacts on lobster vessels: we assume half of the vessels would relocate their traps, and the other half would stop fishing. For relocated vessels, the cost differences come from reduced revenue on the new ground, and extra operating costs to move gear. For vessels that stop fishing, the cost differences include lost revenue, gear moving costs, and saved operating costs from not fishing. The lower and higher end of cost estimates come from the range of lost revenue of the relocated vessels, and a range of gear moving costs for all vessels (see details in the following section).

To estimate catch impacts of the Emergency Rule, we first used the VTR data for 2017-2021 to identify the vessels impacted by each Alternative by using their self-reported fishing coordinates. Although the VTR coordinates only represent the general location of the vessels, it is the best available data for spatial analysis. We then determined the number of vessels and their landing weight for both lobster and Jonah crab. And finally, we calculated the landing value by multiplying the weight and price. The monthly average prices were calculated from NMFS dealer data for 2017-2021. All final values are adjusted to 2021 U.S. dollars by using GDP deflator from U.S. Bureau of Economic Analysis (BEA 2022).

It should be noted that federal permitted fishing vessels that only carry lobster permits are not required to submit VTRs. In order to determine the total number of vessels fishing in this area, we divided the VTR landing value by the percent of VTR vessel coverage. NMFS federal permit data show that from 2017 to 2021, about 41 percent of Massachusetts federal lobster vessels in LMA 1 do not have VTR requirement, which means the landing value from VTR data need to be divided by 59 percent.

Another factor that needs to be considered is the operating cost savings from vessels that stop fishing. Vessel operating costs usually include fuel, bait, ice, fresh water, food and other incidentals. Labor costs are not included because many nearshore vessels are owner-operated, and mates are often paid based on landings rather than by the hour. These costs only occur when the vessel goes on a fishing trip. If a vessel does not fish, then these costs should be considered as savings. We use VTR data to determine the total number of fishing days, and then we apply an average daily operating cost to estimate the total savings.

For the operating costs of transporting gear back to the dock, or to resume fishing outside the restricted area, we assume that fishermen need three to six days to move all their traps around, and multiply that by the daily operating costs based on the average annual operating costs and fishing days for lobster vessels. The detailed results are presented in the next section.

Analysis Results

Vessel Lost Revenue

The emergency restricted area would be in place from February to April. During these months, few vessels were actively fishing and the landings were relatively low compared to summer/fall season. In Table 6.4, we list all lobster and Jonah crab vessels and landing values during February, March, and April from 2017 to 2021 for Alternative 2 and Alternative 3. We also provide the adjusted value by dividing the average value by 59 percent, as not all vessels were reporting their trips. We estimated that about 30 vessels would be affected by Alternative 2, with a total lobster and Jonah crab landing value of \$272,500 (Table 2). Alternative 3 would impact over 50 vessels with a total landing value of \$698,800 (Table 3).

Table 2: The number of affected vessels and landing values in the Alternative 2 from 2017-2021 (in 2021 \$).

	February		March		April	
	Number of vessels	Landing Value	Number of vessels	Landing Value	Number of vessels	Landing Value
2017	18	\$44,672	18	\$37,343	24	\$99,552
2018	25	\$130,445	18	\$64,155	19	\$144,306
2019	16	\$46,591	14	\$35,915	20	\$80,831
2020	19	\$47,206	12	\$22,222	14	\$33,499
2021	13	\$61,224	15	\$43,883	12	\$47,748
Average	18	\$66,028	15	\$40,704	18	\$81,187
Adjusted Average	31	\$112,004	26	\$69,046	30	\$137,719

Notes: 1. Landing values include both lobster and Jonah crab.

2. Both vessel number and landings are from federal VTR data. Based on federal vessel permit data, only 59 percent of Massachusetts federal lobster vessels are required to submit VTR, so the final number are adjusted proportionally to reflect the whole lobster fleet.

Table 3: The number of affected vessels and landing values in the Alternative 3 from 2017-2021 (in 2021 \$).

	February		March		April	
	Number of vessels	Landing Value	Number of vessels	Landing Value	Number of vessels	Landing Value
2017	32	\$144,973	31	\$83,673	39	\$163,309
2018	48	\$488,671	35	\$264,741	39	\$391,033
2019	37	\$194,738	31	\$155,475	35	\$179,161
2020	42	\$250,343	32	\$99,482	31	\$102,029
2021	35	\$266,417	26	\$175,252	26	\$143,211
Average	39	\$269,028	31	\$155,725	34	\$195,749
Adjusted Average	66	\$456,358	53	\$264,159	58	\$332,052

Notes: 1. Landing values include both lobster and Jonah crab.

2. Both vessel number and landings are from federal VTR data. Based on federal vessel permit data, only 59 percent of Massachusetts federal lobster vessels are required to submit VTR, so the final number is adjusted proportionally

to reflect the whole lobster fleet.

Vessel Operating Cost Savings

Vessels that decide to stop fishing during closure months could save some operating costs. We estimated the vessel operating costs based on the cost surveys conducted by the Social Science Branch (SSB) of the Northeast Fisheries Science Center for fishing years 2011, 2012, and 2015. Survey data show that the average annual operating costs for lobster vessels in the Northeast Region is about \$50,365 (in 2021 dollars). Table 4 displays the potential cost savings. We calculate the percentage of trips in each month, and then assign the operating cost to each month based on the trip percentage. At the end, we multiply the cost per vessel and the affected vessel number to get the total cost saving for each month.

Table 4: Cost savings for vessels that stop fishing during closure months (in 2021 \$)

		Affected vessel number	Annual cost per vessel	Closure month trip %	Monthly cost per vessel	Total cost
Alt. 2	Feb	15	\$50,365	4.77%	\$2,403	\$37,092
	Mar	13	\$50,365	3.31%	\$1,669	\$21,806
	Apr	15	\$50,365	4.10%	\$2,067	\$31,210
Alt. 3	Feb	33	\$50,365	4.77%	\$2,403	\$79,075
	Mar	26	\$50,365	3.31%	\$1,669	\$43,894
	Apr	29	\$50,365	4.10%	\$2,067	\$59,614

Notes: 1. We assume that half of the vessels would stop fishing.

2. Annual cost per vessel is based on SSB survey results.

3. Closure month trip percentage is from VTR data.

Final Results

We estimated that 26 to 31 vessels would be affected by Alternative 2 and over 50 vessels affected by Alternative 3. For Alternative 2, the total costs including gear transportation costs and lost revenue range from \$339,000 to \$608,000 for February to April. For vessels moving their gear to new fishing grounds, the total costs are around \$139,000 to \$278,000, about \$3,200 to \$6,400 per vessel; for vessels that stop fishing, the total costs are around \$200,000 to \$331,000, about \$4,600 to \$7,600 per vessel (Table 5). For Alternative 3, the total cost ranges from \$898,000 to \$1,453,000. Total costs for vessels moving their gear to new fishing grounds

range from \$290,000 to \$581,000, about \$3,300 to \$6,600 per vessel. Total costs for vessels that stop fishing are from \$608,000 to \$872,000, about \$6,900 to \$9,900 per vessel (Table 6).

Table 5: Economic impacts of Alternative 2 by month (in 2021\$)

	Feb		March		April		Total	
	Lower	Higher	Lower	Higher	Lower	Higher	Lower	Higher
Relocating costs (half vessels)								
Lost revenue	\$2,800	\$5,600	\$1,726	\$3,452	\$3,443	\$6,886	\$7,969	\$15,938
Gear moving	\$46,310	\$92,619	\$39,185	\$78,370	\$45,292	\$90,583	\$130,786	\$261,572
Sum	\$49,110	\$98,219	\$40,911	\$81,822	\$48,735	\$97,469	\$138,755	\$277,511
Stop fishing costs (half vessels)								
Lost revenue	\$56,002	\$56,002	\$34,523	\$34,523	\$68,860	\$68,860	\$159,385	\$159,385
Gear moving	\$46,310	\$92,619	\$39,185	\$78,370	\$45,292	\$90,583	\$130,786	\$261,572
(Cost savings)	\$37,092	\$37,092	\$21,806	\$21,806	\$31,210	\$31,210	\$90,107	\$90,107
Sum	\$65,219	\$111,529	\$51,903	\$91,088	\$82,942	\$128,233	\$200,064	\$330,850
Total cost	\$114,329	\$209,748	\$92,814	\$172,910	\$131,676	\$225,703	\$338,819	\$608,361

Notes:

1 We estimate lost revenue of the relocating vessels to be between 5 and 10 percent of the total landing value.

2. We estimate gear moving costs to take between 3 and 6 days at \$1,000/day.

Table 6: Economic impacts of Alternative 3 by month (in 2021 \$)

	Feb		March		April		Total	
	Lower	Higher	Lower	Higher	Lower	Higher	Lower	Higher
Relocating costs (half vessels)								
Lost revenue	\$11,409	\$22,818	\$6,604	\$13,208	\$8,301	\$16,603	\$26,314	\$52,628
Gear moving	\$98,726	\$197,452	\$78,879	\$157,758	\$86,512	\$173,025	\$264,117	\$528,234
Sum	\$110,135	\$220,270	\$85,483	\$170,966	\$94,814	\$189,627	\$290,431	\$580,862
Stop fishing costs (half vessels)								
Lost revenue	\$228,179	\$228,179	\$132,079	\$132,079	\$166,026	\$166,026	\$526,285	\$526,285
Gear moving	\$98,726	\$197,452	\$78,879	\$157,758	\$86,512	\$173,025	\$264,117	\$528,234
(Cost savings)	\$79,075	\$79,075	\$43,894	\$43,894	\$59,614	\$59,614	\$182,584	\$182,584
Sum	\$247,829	\$346,555	\$167,064	\$245,943	\$192,924	\$279,437	\$607,818	\$871,935
Total cost	\$357,964	\$566,825	\$252,547	\$416,908	\$287,738	\$469,064	\$898,249	\$1,452,797

Notes:

- 1 We estimate lost revenue of the relocating vessels to be between 5 and 10 percent of the total landing value.
2. We estimate gear moving costs to take between 3 and 6 days at \$1,000/day.

2.4.3. Relative Ranking of Alternatives

As noted above, it is not feasible at present to estimate the economic benefits attributable to each of the regulatory alternatives that NMFS is considering. It is possible, however, to develop a relative ranking of the alternatives with respect to potential benefits, based on the estimated reduction of each alternative on the entanglement risk posed to right whales by commercial trap/pot buoy lines.

The biological impacts analysis presented in the associated EA relies primarily on NMFS' DST to examine how the regulatory alternatives might reduce the possibility of mortality and serious injury between right whales and trap/pot gear. The model integrates information on buoy line

density, line strength, and whale sightings to provide indicators of the potential for mortality and serious injury to occur as a result of entanglement. Risk reduction includes the chance of encounter estimated using co-occurrence and the relative severity of an encounter through the use of line strength. Biological impacts are characterized with respect to the percentage reduction in the overall risk reduction that each alternative would achieve. See FEIS Volume II Appendices 3.1 and 5.1 for more details on how risk reduction is derived.

Table 7 summarizes the compliance costs related to the estimated change in risk reduction under each Alternative relative to the No Action Alternative (Alternative 1). The DST model suggests the risk of right whale mortality or serious injury would be reduced by approximately 13.2 to 16.6 percent under Alternative 2, which has an estimated total compliance costs of \$338,819 to \$608,361 for about 30 vessels. For every unit of gear threat reduction, the cost of Alternative 2 is estimated at \$22,740 to \$40,830.

Alternative 3 achieved better risk reduction than Alternative 2, with an estimated 22.3 to 37.4 percent decrease in right whale entanglement mortality or serious injury. This alternative would increase the likelihood of reducing right whale entanglement risk. However, the total compliance costs associated with risk reduction in Alternatives 3 are substantially higher, ranging from \$898,249 to \$1,452,797; or \$30,092 to \$48,670 for each unit of risk reduction. That is, each risk reduction unit of Alternative 3 would cost about 19 to 32 percent more than Alternative 2.

Table 7: A summary of total compliance costs related to right whale gear threat reduction (2021 dollars)

	Alternative 2	Alternative 3
Affected vessels	26-31	53-66
Risk reduction	13.2 -16.6 %	22.3 -37.4 %
Total compliance costs	\$338,819 - \$608,361	\$898,249 - \$1,452,797
Costs for each unit of risk threat reduction	\$22,740 - \$40,830	\$30,092 - \$48,670

Notes: 1. Risk reduction ranges are based on fully relocated gear and fully removed gear.

2. To calculate the cost per unit of risk reduction, we used the middle point of the risk reduction ranges, as the compliance costs assumed that half of the vessels would relocate and half would stop fishing.

2.4.4. Uncertainties

A few assumptions are made for this analysis.

The first one is that we assumed half of the vessels that had previously fished in the wedge area would suspend their fishing activities, and the other half would relocate their traps to northern waters. The proposed restricted area is located in federal waters and surrounded by the existing MRA. During the April 2022 emergency closure, it was likely difficult for vessels in the southern portion of the restricted area to redistribute their traps outside the northern or eastern

boundaries, given the cost of operation and expected landings in April.¹ Therefore, in our analysis, we split the anticipated reaction of vessels between relocating and suspending fishing.

We also used VTR data in the calculation of the number of vessels and landing value, which may have limitations. We are aware that VTR are self-reported data and the catch and location data are limited in accuracy for some vessels. However, the geographic information and gear configuration data could not be found in any other data sources consistently for lobster and Jonah crab trap/pot fisheries. Therefore, we decided to use the data from recent years (2017-2021) after careful review and the removal of outliers.

Furthermore, due to the incompleteness of federal lobster VTR coverage, we divided the VTR landing value by the percent of VTR vessel coverage in order to estimate the total number of vessels fishing in this area. NMFS federal permit data show that from 2017 to 2021, about 41 percent of Massachusetts federal lobster vessels in LMA 1 did not have VTR requirement, which means the landing value from VTR data needed to be divided by 59 percent.

As previously noted, the inability to quantify and value the benefits of potential temporary changes to the Plan prohibits the use of BCA to identify the regulatory alternative that would provide the greatest net benefit. Instead, Table 5 summarizes the estimated cost of complying with each regulatory alternative combined with the risk reduction estimated for each alternative.

2.5. Results of Regulatory Impact Analysis

The purpose of Executive Order (E.O.) 12866 is to enhance planning and coordination with respect to new and existing regulations. This E.O. requires the Office of Management and Budget to review regulatory programs that are considered to be “significant.” E.O. 12866 requires a review of proposed regulations to determine whether or not the expected effects would be significant, where a significant action is any regulatory action that may:

- Have an annual effect on the economy of \$100 million or more, or adversely affect in a material way the economy, a sector of the economy, productivity, jobs, the environment, public health or safety, or State, local, or tribal governments or communities;
- Create a serious inconsistency or otherwise interfere with an action taken or planned by another agency;
- Materially alter the budgetary impact of entitlements, grants, user fees, or loan programs or the rights and obligations of recipients thereof; or
- Raise novel legal or policy issues arising out of legal mandates, priorities of the President, of the principles set forth in the Executive Order.

In deciding whether and how to regulate, agencies should assess all costs and benefits of available regulatory alternatives, including the alternative of not regulating. Costs and benefits shall be understood to include both quantifiable measures (to the fullest extent that these can be usefully estimated) and qualitative measures of costs and benefits that are difficult to quantify, but nevertheless essential to consider.

¹ Per personal communication with MA DMF on Jan 12, 2023.

Given the information presented in this document and the analysis contained in the associated Environmental Assessment, it is hereby determined that the proposed action will not significantly impact the quality of the human environment as described above and in the Environmental Assessment. In addition, all beneficial and adverse impacts of the proposed action have been analyzed to reach the conclusion of no significant impacts. NMFS has considered the cost information presented above and believes that Alternative 2 (Preferred) offers the best option for achieving the goal of reducing acute entanglement risk adjacent to the MRA in 2023. Alternative 1 remains status quo and though it does not incur a cost, it leaves right whales at risk. Compared to Alternative 2, Alternative 3 provides more protection, but the cost per unit of risk reduction is 19 to 32 percent higher. Based on these considerations, NMFS has identified Alternative 2 (Preferred) as its proposed approach to achieving the goals of the Plan.

3. References

- NMFS. 2002. Workshop on the effects of fishing gear on marine habitats off the Northeastern United States October 23-25, 2001 Boston, Massachusetts. National Marine Fisheries Service, Northeast Fisheries Science Center, Woods Hole, Massachusetts.
- NMFS. 2021. June. Final Environmental Impact Statement (FEIS), Regulatory Impact Review, and Final regulatory Flexibility Analysis for Amending the Atlantic Large Whale Take Reduction Plan: Risk Reduction Rule Volume I.
https://www.greateratlantic.fisheries.noaa.gov/public/nema/aprd/2021FEIS_Volume%20I.pdf
- U.S. Bureau of Economic Analysis (BEA). 2022. “Table 1.1.4. Price Indexes for Gross Domestic Product”. [accessed Feb 10, 2022]. Available from:
<https://apps.bea.gov/iTable/iTable.cfm?reqid=19&step=2#reqid=19&step=2&isuri=1&1921=survey>.